ORIGINAL ARTICLE

RNA-binding Protein UNR Promotes Glioma Cell Migration and Regulates the Expression of Ribosomal Protein L9

Ningyu Tian, Yingjiao Qi, Yan Hu, Bin Yin, Jiangang Yuan,
Boqin Qiang, Xiaozhong Peng*, Wei Han*

State Key Laboratory of Medical Molecular Biology, Department of Molecular Biology and Biochemistry, Institute of Basic Medical Sciences Chinese Academy of Medical Sciences & School of Basic Medicine Peking Union Medical College, Beijing 100005, China

Key words: UNR; glioma; migration; ribosomal protein L9; vimentin

Objective To investigate the role of RNA binding protein—upstream-of-N-Ras (UNR) in the development of glioma and its molecular mechanism.

Methods First, bioinformatics analysis of CGGA database was performed to detect UNR expression level and prognosis of patients with glioma. Western blot and real-time PCR were used to detect UNR expression level in glioma cell lines and tissues. Next, UNR siRNAs were transfected in glioma cells, and MTS assay and scratch wound-healing assay were used to detect changes in cell proliferation and migration. Then, the candidate UNR target mRNAs were identified by analyzing the sequencing data of UNR iCLIP-seq, RNA sequencing and ribosome profiling databases of human melanoma. RNA immunoprecipitation and biotin pull-down assays were used to identify the UNR target mRNAs in glioma cells. Finally, western blot was used to detect the effect of UNR knockdown on ribosomal protein L9 (RPL9) and RPL9 protein expression level in glioma cell lines. RPL9 siRNA was transfected in A172 and T98G and the expression of vimentin in the cells was detected with western blot.

Results Bioinformatics analysis showed that UNR mRNA expression level was significantly higher in high-grade glioma [Grade II ($n=126$), Grade III ($n=51$), Grade IV ($n=128$), $P<0.001$]. UNR high expression levels were associated with poor prognosis ($P=0.0177$). UNR had high expression level in glioma cell lines and patient samples compared with normal cell lines and normal brain samples ($P<0.01$). Knockdown of UNR inhibited glioma cells migration ($P<0.05$), but did not inhibit glioma cells growth in three glioma cell lines. UNR binded the 3' untranslated region (UTR) of PTEN and RPL9 mRNAs. RPL9 protein was significantly highly expressed in most glioma cell lines ($n=9$) and knockdown of UNR resulted in a downregulation of RPL9 protein expression.

Received for publication March 14, 2018. Published online June 1, 2018.

*Corresponding author Xiaozhong Peng Tel: 86-10-69156434, E-mail: pengxiaozhong@pumc.edu.cn; Wei Han Tel: 86-10-69156434, E-mail: hanwei2012@ibms.pumc.edu.cn

Supported by the CAMS Innovation Fund for Medical Sciences (CIFMS, 2016-I2M-1-001).

Cited this article as: Ningyu Tian, Yingjiao Qi, Han Hu, Bin Yin, Jiangang Yuan, Boqin Qiang, Xiaozhang Peng, Wei Han. RNA-binding protein UNR promotes glioma cell migration and regulates the expression of ribosomal protein L9. Chin Med Sci J 2018; 33(3): 143-151.

Epithelial-mesenchymal transition (EMT)-related marker—vimentin was positively regulated by RPL9.

Conclusions UNR could bind to the 3'UTR of PTEN and RPL9 in glioma cell lines, therefore promoting glioma cell migration and regulating the expression of RPL9. Here, we establish a link between UNR and RPL9 protein, which will provide new ideas for the further study of glioma.

RNA-BINDING proteins (RBPs) are particularly important because they can bind hundreds of mRNAs to modulate nearly every post-transcriptional step during the regulation of gene expression.¹ RBPs have diverse biological functions during the process of cell physiology; however, some RBPs have been identified as critical modulators in the development and progression of cancer.² Our lab reported that a RBP, namely PCBP2 has a high expression level in human glioma cell lines and tissues, and that knockdown of PCBP2 inhibits the growth of glioma *in vitro* and *in vivo*. It could thus be seen as a potential drug therapy target for glioma.³ A great many dysregulated RBPs are associated with many cancers; however, we still know little about their roles and molecular mechanisms during cancer development and progression.

Upstream-of-N-Ras (UNR), also known as CSDE1 in mammals, is a kind of RBP that contains five copies of the cold shock domain; it exists widely in organisms. It is located in p13.2 region of chromosome 1, upstream of the *N-ras* oncogene.⁴ UNR has diverse functions in the regulation of gene expression and mainly regulates mRNA translation and stability. It has been reported to be an activator of cap-independent translation initiation in the internal ribosome entry site (IRES) of HRV-2⁵ and poliovirus type 1.⁶ Interestingly, some studies have shown that UNR can bind to the 5'-untranslated region (5'UTR) of other UNR molecules⁷⁻⁸ and poly A-binding protein 1 (PABP1) mRNAs⁹ to act as an inhibitor of translation with its protein partners. Additionally, UNR was found to regulate many of its targets at the level of translation elongation/termination. In summary, whether UNR acts as a regulator depends on its targets or cell types.¹⁰⁻¹¹

UNR participates in diverse cell physiological processes, such as precerebellar neuron migration¹² and maintaining human embryonic stem cells in an undifferentiated state.¹³ UNR is also related to some diseases, such as Diamond-Blackfan anemia,¹⁴ and has genetic associations with autism.¹⁵

However, whether UNR plays roles in human cancer development and progression is still unclear. Recently, a study showed that UNR has a high expres-

sion level in human melanoma and promotes invasion and metastasis.¹⁶ Another study showed that low UNR expression was significantly associated with shorter patient survival and may thus be a potential prognostic biomarker for pancreatic ductal adenocarcinoma.¹⁷ Combining these previous results, we supposed UNR may have potential roles in the progression of other cancers. In our study, we evaluated the potential role of UNR in glioma cell migration and preliminarily investigated its molecular mechanism.

MATERIALS AND METHODS

Patient samples and cell lines

Fresh glioma samples were obtained from Beijing Tiantan Hospital and Beijing Sanbo Brain Hospital.¹⁸ Human glioma cell lines (T98G, A172, U118MG, LN18 and H4) were purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA) and HS683, SF126, SF763, SF767, TJ905, SHG44 and CCF-STTG1 were purchased from the Cell Center of Peking Union Medical College. Three human normal astrocyte cell lines (HA, HA-sp and HA-c) were purchased from ScienCell (Carlsbad, CA, USA). The primary glioma cell line, N3 was identified and donated by Tao Jiang's lab from Beijing Tiantan Hospital.

Cell culture

The U118MG, T98G, SF126, SF763 and SF767 cells were cultured in modified Eagle's medium (MEM), and the A172, LN18, H4, HS683, TJ905 and N3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM). SHG44 and CCF-STTG1 cells were cultured in RPM1-1640 medium. All media were supplemented with 10% foetal bovine serum (FBS; HyClone, Logan, UT, USA), 100 U/ml penicillin (Life Science) and 100 U/ml streptomycin (Life Science). The normal human astrocyte cells (HA, HA-c and HA-sp) were cultured in astrocyte medium (ScienCell). All cells were cultured in a culture chamber with 5% CO₂ at 37°C.

Bioinformatics analysis of CGGA database

To gain insight into the potential role of UNR in glioma

progression, we analyzed the UNR mRNA expression level in 305 glioma patient samples from the Chinese Glioma Genome Atlas (CGGA, <http://www.cgca.org.cn>) and the effect of UNR expression level on glioma prognosis by using Kaplan-Meier analysis.

siRNAs and cell infection

The cDNA target sequences of siRNAs for UNR have been described previously.¹⁹⁻²⁰ The three siRNAs (si1#: GGACAGAAATGGUAAAGAA; si2#: GAGAUAAAGGUUGAAUUUAG; si3#: CCAUAUAGAAGUUCUGUC) could specifically target UNR. Control siRNA: UUCUCCGAACGUGUCACGU. A172, HS683 and LN18, which have relatively high UNR expression levels, were used to detect the UNR protein level after 48 hours of siRNA-mediated knockdown with western blot.

RPL9 siRNA (#s226955, Thermo Scientific, Austin, TX, USA) was used. A172 and T98G were used to detect the RPL9 protein level after 48 hours of siRNA-mediated knockdown with western blot.

MTS assay

Logarithmic growth phase cells were cultured in 96-well plate with the density of 3000 cells/well in 100 μ l medium. 0, 24, 48, 72 and 96 hours after adherence, additional 20 μ l MTS reagent (500 μ g/ml, #G1112, Promega, Madison, USA) was put into the wells of 96-well plate and cells incubated for 2 hours at 37°C for a minimum. Absorbance values were determined at 490 nm using a microplate reader (Multiskan Spectrum, Thermo Fisher, Austin, TX, USA). Each experiment was conducted in triplicate.

Scratch wound-healing assay

For scratch wound-healing assays, when the infected glioma cells reached 80%-90% confluence, we used a 200 μ l pipette tip to create a wound and washed the detached cells twice. Cells were imaged at regular time intervals after scratching, and the migration distance was calculated by measuring the width of the wound with ImageJ software. We got the vertical coordinate values of statistical results by analyzing the width of the wound compared with the initial width of wound area which was set as 100%.

Identification of candidate UNR target mRNAs

Recently, Laurence Wurth and his colleagues revealed the role of UNR in human melanoma. They used a series of high-throughput sequencing technologies

including iCLIP-seq, RNA sequencing and ribosome profiling to unveil CSDE1 targets.¹⁶ GO analysis of UNR targets showed an enrichment in categories related to cell adhesion and the extracellular matrix. To investigate the mechanism of UNR function in glioma, we took advantage of their sequencing data to identify UNR targets. The ArrayExpress accession numbers for the RNA iCLIP-seq, iCLIP-seq, and ribosome profiling experiments in human melanoma databases are E-MTAB-3805, E-MTAB-3818, and E-MTAB-3815, respectively. By analyzing the sequencing data of UNR iCLIP-seq, RNA sequencing and ribosome profiling databases, we totally found 28 direct UNR target mRNA overlaps that changed at both the mRNA and translational levels. Excluding histone and UNR mRNAs, we performed the RNA-binding protein immunoprecipitation experiment and biotin pull-down assay to verify whether UNR can bind to candidate mRNAs, i.e., IER3, PTEN, RPL22L1, RPL9, TIMP2 and TMEM47, in glioma cell lines. We chose GUSB and PABPC1 mRNAs as negative and positive controls, respectively, according to the published data.¹⁶

RNA immunoprecipitation (RIP)

We used the RiboCluster Profiler RIP-Assay Kit (011FA, MBL, Nagoya, Japan) for RNA immunoprecipitation according to the manufacturer's instructions. In brief, we incubated the antibody anti-UNR (A5941, ABclonal, Wuhan, China) and protein A agarose beads overnight in 4°C. Next, prepare new beads and incubate it with the cell lysates with rotation for 1 hour at 4°C to pre-clear the cell lysates. Then, incubate the antibody-immobilized protein A agarose beads with cell lysates after pre-clearing with rotation for 3 hours at 4°C. Finally, wash the beads and isolate the RNA. The real-time PCR was used to identify the targeted mRNA.

Real-time PCR

Real-time RT-PCR was carried out using SYBR Premix Ex Taq Master Mix and a 2-Step kit (TaKaRa, Dalian, China). Total RNA was isolated using TRIzol reagent according to the manufacturer's instructions (Life Technologies). Total RNA of 2.0 μ g was used for cDNA synthesis (Transgene, Beijing, China) and 1/10 of the cDNA volume was used for quantitative PCR (TaKaRa, Dalian, China). The PCR amplification was carried out using a CFX96 touch system (BioRad, Hercules, CA, USA) according to the manufacturer's instructions. The Ct values were normalized to human glyceral-

dehydro-3-phosphate dehydrogenase (GAPDH) gene. The $\Delta\Delta C_t$ method was used to determine the relative expression level of the target genes. All samples were run in triplicate in each experiment. All primers were synthesized by Life Technologies (Beijing, China) and TSINGKE (Beijing, China), and the sequences are listed in Table 1.

Biotin pull-down assay

We inserted the sequence into the pGEM-3zf vector containing a T7 promoter. The GUSB and PABPC1 3'UTRs were used as negative and positive controls, respectively. The LN18 cell lysates pulled down by the probes were analyzed by western blot using an antibody against UNR (A5941, ABclonal, Wuhan, China). The biotin-labeled sense RNA probes were synthesized *in vitro* using T7 RNA polymerase (TaKaRa). Cytoplasmic cell extracts were isolated from LN18 cells using cytoplasmic protein extraction reagents (Thermo Scientific, Austin, TX, USA). RNA affinity capture was conducted with streptavidin-sepharose beads as described previously.³ All primers were synthesized by Life Technologies (Beijing, China) and TSINGKE (Beijing, China), and the sequences are listed in Table 2.

Bioinformatics analysis for RPL9 expression level

We took advantage of the website GEPIA (<http://gepia>).

cancer-pku.cn/index.html), a newly developed interactive web server for analyzing RPL9 RNA sequencing expression based on 9736 tumors and 8587 normal samples from the TCGA and GTEx databases.²¹

Western blot

Whole-cell extracts were obtained by lysing cells in TNTE buffer (50 mmol/L Tris, pH 7.4, 150 mmol/L NaCl, 1 mmol/L EDTA, 10 mmol/L sodium pyrophosphate, 0.5% Triton X-100, 1 mmol/L sodium vanadate, and 25 mmol/L sodium fluoride) containing protease inhibitors (5 μ g/ml PMSF, 0.5 μ g/ml leupeptin, 0.7 μ g/ml pepstatin, and 0.5 μ g/ml aprotinin). The detailed western blot procedures have been described previously.²²

The protein samples were analyzed using polyclonal rabbit anti-UNR (A5941, ABclonal); rabbit anti-RPL9 (ab182556; Abcam, Cambridge, MA, USA); mouse anti- β -actin (A5441, Sigma-Aldrich, St. Louis, USA) and rabbit anti-vimentin (bs-0756R, BIOSS, Beijing, China) antibodies.

Statistical analysis

Data were expressed as the means $\pm$ SD unless otherwise indicated and were analyzed with SPSS 14.0 software (SPSS, Chicago, IL, USA) and GraphPad Prism. Student's *t*-test was used for statistical analysis. *P* < 0.05 was considered statistically significant.

Table 1. Sequences of PCR amplification

Name	Forward (5'-3')	Reverse (5'-3')
GUSB	CAAGCATGAGGATGCGGACA	GCTGGTACGGAAGCGTTGG
PABPC1	CAGGCTCACCTCACTAACCAG	GGTAGGGTTGATTACAGGGT
IER3	CAGCCGACGGGTCTCTAC	GATCTGGCAGAAGACGATGGT
PTEN	ACAAAGCCAACCGATACTTTTCTC	GAAGTTGAAGTCTAGCCTCTGG
RPL22L1	GCAATTTCTACGGGAGAAGGTT	ACTCGAAGCCAATCACGAAGA
RPL9	GCACAGTTATCGTAAGGGC	TTACCCACCATTTGTCAACC
TIMP2	FGCCAAAGCGGTCAGTGAGAA	GGAGGGGGCCGTGTAGATAA
TMEM47	CAGCAGCGATTGGCAGATTG	TCCACGCGAGATAGAAATCAAAC

GUSB: glucuronidase beta; PABPC1: poly(A) binding protein cytoplasmic 1; IER3: immediate early response 3; PTEN: phosphatase and tensin homolog; RPL22L1: ribosomal protein L22 like 1; RPL9: ribosomal protein L9; TIMP2: tissue inhibitor of metalloproteinase 2; TMEM47: transmembrane protein 47.

Table 2. Sequences for pGEM-3zf plasmids construction

Name	Forward (5'-3')	Reverse (5'-3')
GUSB 3'UTR	TCTGATGTTCACTGAAGAGTACCAG	TTCAGTAGCCACTTTTCATGCC
PABPC1 3'UTR	TTGATCAGGGACCATGAAAAG	TTCAGTGTAAAGGAATTTTATTAAAGC
PTEN 3'UTR (Fragment A)	GTGGAGGCTATCAACAAAGAATG	GGAGGGGGCCGTGTAGATAA
RPL9 3'UTR	GATCTAAGAGTTACCTGGCTACAGAAA	TTTTTAAGAAGAAGGTCCAAATCAATA

3'UTR: 3' untranslated region.

RESULTS

UNR is upregulated in glioma cell lines and glioma tissues

Bioinformatics analysis showed that the UNR mRNA expression level was significantly higher in high-grade glioma (Fig. 1A). Kaplan-Meier curves showed that a high expression level of UNR correlated with poor survival of patients according to the CGGA database (Fig. 1B, $P=0.0177$). Further, real-time PCR and western blot showed that UNR expression level was up-regulated in glioma tissue samples (Figs. 1C, 1D). Next, we detected the protein expression level using 9 glioma cell lines along with 3 normal human astrocyte cell lines, which were set as the normal control. The results showed that UNR protein had significantly higher expression levels in a variety of glioma cell lines (Fig. 1E).

Knockdown of UNR expression inhibits glioma cell migration but does not inhibit glioma cell growth

In A172, the three siRNAs all worked well, where-

as only si1#, si2# in HS683 and si1#, si3# in LN18 worked well compared with the control siRNA (Fig. 2A).

MTS assay results showed that knockdown of UNR for 0 to 96 hours did not significantly inhibit glioma cell growth (Fig. 2B). To further confirm the role of UNR in glioma cell lines, we performed the wound-healing assay. Images were captured; HS683 and LN18 were observed from 0 to 72 hours, and A172 was observed from 0 to 36 hours. Strikingly, the results demonstrated that knock-down of UNR dramatically inhibited the cell migration associated with closing an artificial wound created over a confluent monolayer ($P<0.05$, Figs. 2C, 2D).

UNR binds the 3'UTR of biotin-labeled candidate mRNAs

RNA immunoprecipitation assay confirmed the enrichment of PTEN, RPL9 and RPL22L1 mRNAs in LN18 by UNR antibody (Fig. 3A). To further verify the result, a biotin pull-down assay was applied. The iCLIP-seq data were used to identify a potential binding sequence between UNR and the candidate mRNAs' 3' untranslated

Figure 1. Upstream-of-N-Ras (UNR) is upregulated in glioma cell lines and tissues compared with normal human astrocytes and brain tissues.

A. UNR mRNA expression level in whole genome gene profiling of glioma samples from the Chinese Glioma Genome Atlas (CGGA, <http://www.cgga.org.cn>); Grade II ($n=126$), Grade III ($n=51$), Grade IV ($n=128$) ($*P<0.05$, $**P<0.005$, $***P<0.001$, Student's t -test). B. Kaplan-Meier curves showing high expression levels of UNR correlated with poor patient survival according to the CGGA database. C. Real-time PCR analysis of the relative UNR mRNA expression level in normal human brain ($n=5$), Grade II ($n=11$), Grade III ($n=20$), and Grade IV ($n=32$) glioma tissues ($*P<0.05$, $***P<0.001$, Student's t -test). D. Relative UNR protein levels in normal brain ($n=7$), Grade II ($n=12$), Grade III ($n=24$), and Grade IV ($n=47$) glioma tissues analyzed by western blot ($*P<0.05$, $**P<0.01$, Student's t -test). E. Western blot showing the relative UNR protein expression level in 3 normal human astrocyte cell lines and 9 glioma cell lines.

Figure 2. Knockdown of UNR inhibits glioma cell migration.

A. Western blot analysis of UNR expression in 3 glioma cell lines (A172, LN18 and HS683) transiently transfected with the control siRNA or UNR siRNA for 48 hours. β -Actin was used as a loading control. B. MTS assay on 3 glioma cell lines after transfection with the control siRNA or UNR siRNA as above. The data have no statistical significance compared with the control siRNA. C, D. Wound-healing assays in HS683 and LN18 cells were performed after infection with control siRNA and UNR-siRNA for 0-72 hours. The assays in A172 cells were performed after infection with control siRNA and UNR-siRNA for 0-36 hours. Pictures were taken during this period. The graph represents the width of the remaining open wound calculated in relation to time 0 (* $P < 0.05$ compared with control siRNA, student's t -test). All samples were treated in triplicate in each experiment. Each assay was repeated three times.

region (3'UTR); the red fragments represent the iCLIP-seq sequence, and the black fragments represent the probe sequences that were designed between 100 bp and 500 bp for the biotin pull-down assay to protect their secondary binding structures (Fig. 3B). The resulted revealed UNR was not detected in either the no-probe group or the GUSB 3'UTR pull-down complexes. In contrast, the PABPC1 3'UTR, PTEN 3'UTR fragment A and RPL9 3'UTR showed strong binding with UNR (Fig. 3C). In our study, we focused on whether RPL9 expression could be regulated by UNR in glioma. UNR regulation of PTEN expression has been studied in hu-

man melanoma¹⁶ which needs further study in glioma in the future.

RPL9 has a high expression level in glioma and knockdown of UNR expression inhibits the expression of RPL9

We noticed that knockdown of UNR resulted in a down-regulation of RPL9 protein expression in the A172, LN18 and HS683 glioma cell lines (Fig. 4A). GEPIA website analysis showed RPL9 has a higher expression level in glioma than in normal tissues (Fig. 4B). Further, we detected the protein expression levels of

Figure 3. UNR binds 3'UTR of biotin-labeled candidate mRNAs.

A. Real-time PCR of the candidate mRNAs by RNA binding protein immunoprecipitation of UNR antibodies in LN18 cell lysates. B. Schematic of the indicated candidate mRNAs. The red fragments represent the iCLIP-seq sequence, and the black fragments represent the probe sequences that were designed between 100 bp and 500 bp for the biotin pull-down assay to protect their secondary binding structures. C. Biotin pull-down analysis of complexes formed *in vitro* using biotin-labelled candidate mRNA segments (shown in B) in LN18 cell lysates.

RPL9 by using 9 glioma cell lines, including a primary glioma cell line, N3, and a normal human astrocyte cell line HA. The results showed that the RPL9 protein was significantly highly expressed in most glioma cell lines, which mostly coincided with UNR (Fig. 4C). Meanwhile, knockdown of RPL9 inhibits the expression of vimentin, an epithelial-mesenchymal transition marker, which could maintain cellular integrity and modulate the motility of cancer cells (Fig. 4D).

DISCUSSION

In our research, we found that UNR was overexpressed in glioma and that high expression levels were associated with poor prognosis. We revealed the role of UNR in glioma cell migration, which may contribute to the clinical treatment of glioma. We also took advantage of sequencing data from human melanoma and verified that UNR can bind to the 3'UTR of PTEN and to RPL9 in glioma. Moreover, RPL9, which can be positively regulated by UNR, has a high expression level in glioma. Knockdown of RPL9 inhibits the expression of

the epithelial-mesenchymal transition marker vimentin, which suggests it may have an important role in glioma migration and invasion.

Although RBPs have been associated with the progression of cancer, UNR has not been well studied in this area. Some researchers have shown that UNR knockdown induces apoptosis in human hepatoma cells, but the detailed mechanism is unclear.²⁰ Recently, UNR was reported to have a high expression levels in human melanoma and to promote invasion and metastasis. Another study has shown that low UNR expression was associated with shorter patient survival.¹⁷ Based on these results, we hypothesize that UNR may have entirely different roles, depending on the cancer type.

Gliomas are the most common type of primary malignant tumors in the central nervous system. It has features such as high lethality, a low cure rate and easy recurrence.²³ Thus, studying the essential carcinogenesis factors and their molecular mechanism during the development and progression of cancer can have great clinical impacts. In our previous study, we discovered that UNR has a higher expression level in

Figure 4. Downregulation of UNR inhibits the expression of RPL9 and RPL9 has high expression levels in gliomas.

A. Knockdown of UNR inhibits the expression of RPL9 in the glioma cell lines assayed by western blot. B. RPL9 has a higher expression level in gliomas than in normal tissues in the TCGA and GTEx projects by using the GEPIA website (GBM: glioblastoma multiform, LGG: low grade glioma). Red: glioma patients. Gray: normal patients. C. Western blot showing the RPL9 protein levels in normal human astrocyte and glioma cell lines. D. Western blot analysis of RPL9 and vimentin expression in A172 and T98G cells transiently transfected with the control siRNA (SiNC) or RPL9 siRNA for 48 hours.

high-grade glioma patients than in normal brain tissues. This gives us a hint that UNR may participate in the malignant progression of glioma. However, we only proved UNR can promote glioma cell migration *in vitro*, which should be further proven *in vivo*.

RPL9 was identified as a Bax suppressor that promotes cell survival in yeast and is overexpressed in colorectal cancer. In addition, knockdown of RPL9 expression inhibits colorectal carcinoma growth. However, there are few reports about its role in other types of cancers. Luckily, we found that it is overexpressed in glioma and is a downstream target gene of UNR. A series of assays provided evidence that UNR can bind to the RPL9 3'UTR in glioma cells. According to the known mechanisms of UNR as an mRNA regulator, it may maintain the stability of mRNA 3'UTR or regulate mRNA at the level of translation elongation/termination, which should be investigated further. Moreover, knockdown of RPL9 inhibits the expression of the vimentin, which provides a research clue for us.

In conclusion, UNR could bind to the 3'UTR of PTEN and RPL9 in glioma cell lines, therefore promot-

ing glioma cell migration and regulating the expression of RPL9. This will provide a new idea for the further study of glioma.

Conflicts of interests statement

The authors declare no conflict of interests.

Acknowledgments

We are grateful to Professor H.Jacquemin-Sablon from INSERM, E362, Bordeaux, France for providing us UNR plasmids and useful discussions on UNR. We also thank Professor F.Gebauer from the Barcelona Institute of Science and Technology for suggesting the UNR siRNA sequence.

REFERENCES

1. Marchese D, de Groot NS, Lorenzo Gotor N, et al. Advances in the characterization of RNA-binding proteins. *Wiley Interdiscip Rev RNA* 2016; 7(6):793-810. doi: 10.1002/wrna.1378.
2. Campos-Melo D, Droppelmann CA, Volkening K, et al.

- RNA-binding proteins as molecular links between cancer and neurodegeneration. *Biogerontology* 2014; 15(6):587-610. doi: 10.1007/s10522-014-9531-2.
3. Han W, Xin Z, Zhao Z, et al. RNA-binding protein PCBP2 modulates glioma growth by regulating FHL3. *J Clin Invest* 2013; 123(5):2103-18. doi: 10.1172/JCI61820.
 4. Jeffers M, Paciucci R, Pellicer A. Characterization of UNR: a gene closely linked to N-ras. *Nucleic Acids Res* 1990; 18(16):4891-9.
 5. Hunt SL, Hsuan JJ, Totty N, et al. UNR, a cellular cytoplasmic RNA-binding protein with five cold-shock domains, is required for internal initiation of translation of human rhinovirus RNA. *Genes Dev* 1999; 13(4):437-48.
 6. Boussadia O, Niepmann M, Creancier L, et al. UNR is required *in vivo* for efficient initiation of translation from the internal ribosome entry sites of both rhinovirus and poliovirus. *J Virol* 2003; 77(6):3353-9.
 7. Cornelis S, Tinton SA, Schepens B, et al. UNR translation can be driven by an IRES element that is negatively regulated by polypyrimidine tract binding protein. *Nucleic Acids Res* 2005; 33(10):3095-108.
 8. Dormoy-Raclet V, Markovits J, Jacquemin-Sablon A, et al. Regulation of UNR expression by 5'- and 3'-untranslated regions of its mRNA through modulation of stability and IRES mediated translation. *RNA Biol* 2005; 2(3):e27-35.
 9. Patel GP, Ma S, Bag J. The autoregulatory translational control element of poly(A)-binding protein mRNA forms a heteromeric ribonucleoprotein complex. *Nucleic Acids Res* 2005; 33(22):7074-89.
 10. Elatmani H, Dormoy-Raclet V, Dubus P, et al. The RNA-binding protein UNR prevents mouse embryonic stem cells differentiation toward the primitive endoderm lineage. *Stem Cells* 2011; 29(10):1504-16. doi: 10.1002/stem.712.
 11. Dinur M, Kilav R, Sela-Brown A, et al. *In vitro* evidence that upstream of N-ras participates in the regulation of parathyroid hormone messenger ribonucleic acid stability. *Mol Endocrinol* 2006; 20(7):1652-60.
 12. Kobayashi H, Kawauchi D, Hashimoto Y, et al. The control of precerebellar neuron migration by RNA-binding protein Csd1. *Neuroscience* 2013; 253:292-303. doi: 10.1016/j.neuroscience.2013.08.055.
 13. Ju Lee H, Bartsch D, Xiao C, et al. A post-transcriptional program coordinated by CSDE1 prevents intrinsic neural differentiation of human embryonic stem cells. *Nat Commun* 2017; 8(1):1456. doi: 10.1038/s41467-017-01744-5.
 14. Horos R, Ijspeert H, Pospisilova D, et al. Ribosomal deficiencies in Diamond-Blackfan anemia impair translation of transcripts essential for differentiation of murine and human erythroblasts. *Blood* 2012; 119(1):262-72. doi: 10.1182/blood-2011-06-358200.
 15. Xia K, Guo H, Hu Z, et al. Common genetic variants on 1p13.2 associate with risk of autism. *Mol Psychiatry* 2014; 19(11):1212-9. doi: 10.1038/mp.2013.146.
 16. Wurth L, Papasaikas P, Olmeda D, et al. UNR/CSDE1 drives a post-transcriptional program to promote melanoma invasion and metastasis. *Cancer Cell* 2016; 30(5):694-707. doi: 10.1016/j.ccell.2016.10.004.
 17. Martinez-Useros J, Georgiev-Hristov T, Fernandez-Acenero MJ, et al. UNR/CDSE1 expression as prognosis biomarker in resectable pancreatic ductal adenocarcinoma patients: a proof-of-concept. *PLoS One* 2017; 12(8):e0182044. doi: 10.1371/journal.pone.0182044.
 18. Hu PS, Xia QS, Wu F, et al. NSPc1 promotes cancer stem cell self-renewal by repressing the synthesis of all-trans retinoic acid *via* targeting RDH16 in malignant glioma. *Oncogene* 2017; 36(33):4706-18. doi: 10.1038/onc.2017.34.
 19. Ray S, Anderson EC. Stimulation of translation by human Unr requires cold shock domains 2 and 4, and correlates with poly(A) binding protein interaction. *Sci Rep* 2016; 6:22461. doi: 10.1038/srep22461.
 20. Dormoy-Raclet V, Markovits J, Malato Y, et al. UNR, a cytoplasmic RNA-binding protein with cold-shock domains, is involved in control of apoptosis in ES and HuH7 cells. *Oncogene* 2007; 26(18):2595-605.
 21. Tang Z, Li C, Kang B, et al. GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res* 2017; 45(W1):W98-W102. doi: 10.1093/nar/gkx247.
 22. Zhang Y, Chao T, Li R, et al. MicroRNA-128 inhibits glioma cells proliferation by targeting transcription factor E2F3a. *J Mol Med (Berl)* 2009; 87(1):43-51. doi: 10.1007/s00109-008-0403-6.
 23. Legler JM, Ries LA, Smith MA, et al. Cancer surveillance series [corrected]: brain and other central nervous system cancers: recent trends in incidence and mortality. *J Natl Cancer Inst* 1999; 91(16):1382-90.

RNA 结合蛋白 UNR 促进人神经胶质瘤细胞的迁移并调控核糖体蛋白 L9 的表达

田宁宇, 齐颖娇, 胡艳, 阴彬, 袁建刚, 强伯勤, 韩为*, 彭小忠*

中国医学科学院基础医学研究所 北京协和医学院基础学院,
国家医学分子生物重点实验室, 北京 100005

关键词: UNR; 神经胶质瘤; 迁移; 核糖体蛋白 L9; 波形蛋白

目的 研究 RNA 结合蛋白 UNR 在神经胶质瘤发生发展中的作用以及其发挥作用的分子机制。

方法 首先, 利用生物信息学方法分析 CGGA 数据库中 UNR 的表达水平及其表达水平高低与病人预后的关系。采用 western blot 和 real-time PCR 方法分析胶质瘤病人组织样本和细胞株中 UNR 的表达水平。其次, 在胶质瘤细胞株 A172, HS683, LN18 中转染 siRNA 敲低 UNR, 采用 MTS 和划痕实验检测肿瘤细胞增殖能力以及迁移能力的变化。再次, 通过分析已发布的黑色素瘤 UNR iCLIP-seq、RNA-seq 和核糖体印迹测序数据库筛选出 UNR 在胶质瘤细胞中可能结合的靶 mRNA。利用 RNA 免疫沉淀和生物素标记的 RNA pull down 实验确认 UNR 可结合核糖体蛋白 L9 (RPL9) 的 mRNA。最后, 通过 western blot 检测 RPL9 在胶质瘤细胞株中的表达水平, 并且检测敲低 UNR 以后 RPL9 蛋白水平的变化。在神经胶质瘤细胞株 A172, T98G 中转染 siRNA 敲低 RPL9, 检测和肿瘤侵袭迁移有关的上皮向间充质转化有关的标志物—波形蛋白的表达变化。

结果 生物信息学分析 UNR mRNA 表达水平, 发现其在恶性程度更高的胶质瘤样本中表达较高 ($P < 0.001$)。高表达 UNR 的患者预后较差 ($P = 0.0177$)。与正常细胞株以及正常脑组织样本相比, UNR 在 9 株胶质瘤细胞以及患者组织样本中具有较高的表达水平 ($P < 0.01$)。敲低 UNR 抑制肿瘤细胞的迁移能力 ($P < 0.05$), 但不影响其增殖能力。UNR 可以结合在 PTEN mRNA 和 RPL9 mRNA 的 3' 非翻译区。敲低 UNR 以后 RPL9 的表达水平下调。RPL9 在 9 株胶质瘤细胞中具有高表达的趋势, RPL9 可以正向调控波形蛋白的表达。

结论 本研究揭示了 RNA 结合蛋白 UNR 在胶质瘤细胞迁移方面的功能, 并且发现 UNR 可以结合 PTEN 和 RPL9 mRNA 的 3' 非翻译区并调控 RPL9 的表达。这为我们更进一步研究胶质瘤的发生发展提供了新思路。