ORIGINAL ARTICLE

Physicians' Perception of Palliative Care Consultation Service in a Major General Hospital in China

Xuan Qu, Nan Jiang, Nan Ge, Xiaohong Ning*

Department of Geriatrics, Peking Union Medical College Hospital,
Chinese Academy of Medical Sciences & Peking Union
Medical College, Beijing 100730, China

Key words: palliative care; consultation; medical service

Objectives The in-hospital palliative care consultation (PCC) is emerging as a routine service in some medical center in China. The current study evaluated how physicians in primary care team and consultation team perceive the PCC service for the purpose of investigating the effectiveness of this consultation model in a general hospital.

Methods In-hospital palliative care consultations have been carried out at Peking Union Medical College Hospital by a dedicated consultation team, and 37 consultations were completed in 2016. A questionnaire was designed for physicians in terms of its benefits to patients, their family as well as the primary care team. Physicians who applied for consultation in 2016 formally (requested from the department other than the Geriatrics) and informally (by rotating residents and unemployed visiting doctors in geriatric department) were invited to participate in the survey by scanning a two dimensional code on social networking platform.

Results There were 103 physicians participated in the survey, including primary care physicians from the department of Internal Medicine ($n=8$), Gynaecology ($n=16$) and Surgery ($n=13$), rotating residents ($n=30$), visiting doctors ($n=16$) in Geriatric department, and PCC team members ($n=20$). 94.0% of the non-PCC physicians agreed that PCC relieved the suffering of patients; 89.2% thought PCC improved the quality of patients' life; there were 91.6%, 95.2%, 90.4% physicians who felt it relieved the anxiety of patients, of family members and of care providers, respectively. There were 96.4% physicians who felt it could ease the tension in physician-patient relationship; 97.6% felt it lower the risk for medical negligence, and 96.4% of doctors who applied for PPC felt satisfied with PCC service in terms of process and achieving objectives of consultation. More primary-team physician agree "PCC service helps the physicians better understand palliative care" than PCC members (97.6% vs. 80%, $P<0.05$), while both were interested in learning more on palliative medicine (100% vs. 96.4%, $P>0.05$).

Received: September 18, 2018; accepted: October 22, 2018.

*Corresponding author Tel: 86-10-69154065; E-mail: ningxh1973@foxmail.com

Fund supported by the Educational Reform Project of Peking Union Medical College (2015zlgc012).

Cited this article as Qu X, Jiang N, Ge N, et al. Physicians' perception of palliative care consultation service in a major general hospital in China. Chin Med Sci J 2018; 33(4): 228-233. doi:10.24920/003515.

Conclusion Palliative care consultation service in a general hospital is efficacious and acclaimed. The primary care physicians and the PCC members hold positive attitudes to the benefits that the PCC services bring to patients, family members, and physicians themselves. PCC for terminal patients in a general hospital may serve as a good model for promotion of palliative care in China.

WITH the aging of the population, an ever-increasing number of patients with severe or end-stage illnesses are witnessed. The advance of life-support technologies have prolonged lives of severely diseased patients, while the quality of life and the suffering of patients and their family are often relatively neglected. Palliative care provides approaches that improve the life quality of patients and help their families to get through problems that are associated with life-threatening illness through prevention, early identification, impeccable assessment, and managements of pain, physical, psychosocial and spiritual problems. The goal is to help terminal patient have peace, comfort and dignity. The palliative care uses team approach, with extensive joint efforts of primary-team physicians, palliative care specialists, nurses, social workers and volunteers.¹

Palliative care has been widely accepted in western countries as well as in Chinese Hong Kong and Chinese Taiwan. In mainland China, the cognition and practice of palliative care is still at preliminary stage. The palliative care services in mainland China mainly include specialized palliative care centers, consultation in general hospitals, and home-based caring service, etc.

Peking Union Medical College Hospital (PUMCH), as one of the top acknowledged general hospitals in mainland China, has developed their palliative care team in recent years and provide specialized palliative care consultation (PCC) service for all inpatients. Upon request by primary care physician in the hospital, PCC team members visit the patient and his/her family, participate in the physician-patient communication, provide pain-relief, symptom control and bereavement care. This survey was conducted aiming at learning how primary care physicians and the consulting physicians perceive the benefits that PCC services bring to patient, patient's family and physicians themselves.

MATERIALS AND METHODS

Questionnaire design

To understand the perspectives and opinions of primary care team and PCC team on the effectiveness

of PCC service, we designed a questionnaire survey including 18 questions (**Table 1**) through Wenjuan Xing, a commercial customized questionnaire platform (Ranxing Inc. Changsha, Hunan, China) (<http://www.sojump.com>). Participants open the questionnaire through scanning two dimensional code using their mobile phone and answer the questions by checking a single answer from the five options: strongly disagree, disagree, uncertain, agree and strongly agree.

Survey performance and targeted participants

From January to December in 2016, our PCC team have done 37 in-hospital consultations by requests from the primary care physicians in the department of internal medicine ($n=8$), gynaecology ($n=16$), and surgery ($n=13$). We sent an invitation of survey through WeChat to the physicians who applied for consultation. As residents on rotation and non-employed visiting

Table 1. The contents of the questionnaire

Does the patient benefit from PCC service?
Q1. Pain and symptom relieved
Q2. Decreased anxiety of the patient
Q3. Improved life quality
Q4. Better understanding of death
Q5. Increased patients' adherence to treatment
Do patients' families benefit from PCC service?
Q6. Decreased anxiety of family members
Q7. Better understand the circumstances
Q8. Knows what they can do for the patient
Do primary physicians benefit from PCC service?
Q9. Decreased anxiety
Q10. Improved understanding of palliative care
Q11. Increased confidence
Q12. Better cope with future patients
Q13. Relieved occupational burnout
Q14. Improved physician-patient relationship
Q15. Decreased risk for medical negligence
Overall attitude to the PCC service?
Q16. The objectives of consultation accomplished
Q17. Satisfied with the process of service
Q18. Interested in learning more on palliative medicine
PCC: palliative care consultation.

physicians in geriatrics department provide end-stage patients care and ask for a lot palliative consultations informally, they were also invited to participate in the survey as primary care physicians.

The PCC team consists of members from various disciplines, including 12 physicians, 4 nurses, 1 pharmacist, 1 physical therapist, 1 nutritionist, and 1 social worker. They all participated the survey.

Statistic analysis

For each question, answer of agree or strongly agree were defined as a positive answer. Enumeration data were described as percentage. The agree rates for each question were compared between the PCC team and non-PCC team using Chi-square test. Statistical analysis was performed by using SPSS (version 23.0), and statistical significance was considered when *P* value was less than 0.05.

RESULTS

Participants

Totally 103 professionals completed the questionnaires,

including 37(35.9%) physicians from the primary teams who requested PCC service formally, 46 (44.7%) physicians who requested PCC informally (30 rotating residents and 16 visiting physicians), and 20(19.4%) professionals in PCC team. Among all the participants, 46 (44.7%) have working experience of 1-3 years, 16(15.5%) have experience of 4-6 years, 15 (14.6%) have experience of 7-9 years, and 26 (25.2%) have worked for over 10 years.

Perceptives of the primary team for palliative care consultation

As shown in **Table 2**, most non-PCC team physicians agreed that PCC consultation helped to relieve patients' symptoms, increase patients' adherence to the treatment, decrease patients' anxiety. They also thought that PCC service helped to improve patients' quality of life and the relationship between healthcare providers and patients, and reduce risk for medical negligence.

Attitudes of PCC members to consultation service

PCC members also thought they helped to relieve patient's symptoms and improve doctor-patient commu-

Table 2. Physicians' perception of the effectiveness of PCC service between PCC members and non-PCC physicians (*n*=103)

Questions in the survey	Agree in total (<i>n</i> =103) <i>n</i> (%)	Agree in PCC team members (<i>n</i> =20) <i>n</i> (%)	Agree in non-PCC phy- sicians (<i>n</i> =83) <i>n</i> (%)	χ^2	<i>P</i> value
Does the patient benefit from PCC service?					
Pain and symptom relieved	96(93.2)	18(90.0)	78(94.0)	0.402	0.409
Decreased anxiety of the patient	94(91.3)	18(90.0)	76(91.6)	0.05	0.555
Improved life quality	92(89.3)	18(90.0)	74(89.2)	0.012	0.638
Better understanding of death	93(90.3)	18(90.0)	75(90.4)	0.002	0.618
Increased patients' adherence to treatment	98(95.1)	19(95.0)	79(95.2)	0.001	0.668
Do patients' families benefit from PCC service?					
Decreased anxiety of the family members	96(93.2)	17(85.0)	79(95.2)	2.637	0.131
Better understand the circumstances	94(91.3)	19(95.0)	75(90.4)	0.435	0.445
Knows what they can do for the patient	100(97.1)	19(95.0)	81(97.6)	0.382	0.48
Do primary physicians benefit from PCC service?					
Decreased anxiety	92(89.3)	17(85.0)	75(90.4)	0.486	0.362
Improved understanding of palliative care	97(94.2)	16(80.0)	81(97.6)	9.09	0.012
Increased confidence	93(90.3)	17(85.0)	76(91.6)	0.793	0.301
Better cope with future patients	98(95.1)	18(90.0)	80(96.4)	1.423	0.249
Relieved occupational burnout	92(89.3)	17(85.0)	75(90.4)	0.486	0.362
Improved physician-patient relationship	99(96.1)	19(95.0)	80(96.4)	0.083	0.584
Decreased risk for medical negligence	99(96.1)	18(90.0)	81(97.6)	2.488	0.169
Overall attitude?					
Objectives of consultation accomplished	99(96.1)	19(95.0)	80(96.4)	0.083	0.584
Satisfied with the process of service	98(95.1)	18(90.0)	80(96.4)	1.423	0.429
Interested to learn more on palliative medicine	100(97.1)	20(100)	80(96.4)	0.745	0.52

nication greatly. Interestingly, for the question "PCC helped the primary team get a better understanding of palliative care", 80% of the PCC team members hold positive attitude, compared to 97.6% of non-PCC physicians agree the statement ($\chi^2=9.09$, $P=0.012$), but both were interested in learning more on palliative medicine (100% vs. 96.4%, $\chi^2=0.745$ $P=0.52$). For the other questions, majority of PCC members and non-PCC physicians hold the positive attitudes, with no significant difference in the agree rates between two groups.

DISCUSSION

The aging population and the increased prevalence of multiple chronic diseases lead to large unmet need of palliative care around the world. The prevalent incidence of chronic diseases, such as aging and cancer, increase the demand for palliative care in the world.² Service of palliative consultation was firstly established at St. Thomas' Hospital in 1976, and quickly became popular in the western countries.³ Up to 2003, 1027 hospitals in the US have established palliative consultation service.⁴ However, in China, less than 1% of population has access to palliative care service.⁵

For patients with advanced cancer who have reached the end of their lives and can not tolerate surgery or chemotherapy, or patients with chronic diseases who have undergone standardized treatment but the condition is still progressing, traditional radical treatments such as tracheal intubation may allow patients to die in pain. The family is also suffering. Lacking systemic knowledge of palliative care usually renders the primary team physicians fail to control pain and maintain the dignity of patients in the inevitable dying process.^{6,7}

With the popularization of palliative medical concepts and the development of professional consultation services, primary team doctors can request palliative medical consultations to help to relieve patient's symptoms and improve communication between doctors and patients. At the same time, consultation put great attention to the convening of family meetings, where patient's will for the treatment, the caring approaches and whether to inform the patient of the situation can be discussed among the family members, primary medical staff, and PCC members. During the consultation, physical discomforts such as pain, dyspnea, nausea and vomiting, bowel obstruction, delirium, etc.,

as well as psychological and spiritual issues were fully cared for. Difficulties in patient management were discussed during consultation within the team, and the outcomes were always actively followed up to build experiences and improve ability of team member. The patient-centered care has shown promising results in controlling symptoms and improving the quality of life.⁸

In the 1990s, palliative care consultation teams (PCCTs) comprising a multidisciplinary group of health care providers were established in many Western countries, with the aim of maximizing quality of life for patients and their families facing life-threatening illness. Since then, the number of hospital-based PCCTs has increased markedly in various countries, including the United Kingdom, the United States, Canada, and Australia, and in all cases these teams play important roles in the overall health care system.^{9,10} Many systematic reviews,¹¹ randomized control trials^{12,13} and cohort studies^{14,15} have also reported on the efficacy and activity of PCCTs.

We found that primary team physicians thought highly of the effectiveness of palliative care consultation services in terms of benefit to the patients, their families, and benefit to the primary team physicians. They felt satisfied with the outcome of consultation. Previous studies have shown the effectiveness of PCC service.^{12,16} According to a systemic review by Higginson et al., hospital-based palliative teams was found to be effective in reducing length of stay, improving symptoms control and enhancing communication. Palliative care consultations have significantly lead to improvements in quality of life and quality of care.¹⁷

This study showed a high satisfaction rate in general, however, in daily medical practices, we found the cooperation between the primary team and the PCC team was not always as smooth as expected. Some of the treatment methods proposed by the PCC consultation members were not fully accepted by the primary care team and occasionally they failed in effective communication. Timely information exchanges are needed for PCC team to better understand the difficulties that primary care team need helps to deal with; meanwhile, the concept and knowledge of palliative care need to be disseminated more widely and deeply in clinical professionals. By doing so, better terminal patient care could be achieved on a regular base.

Our study showed that most professionals in survey hold positive attitudes to the palliative consultaion

in hospital, and were willing to receive training on palliative care. However, as a systematic discipline, major medical schools in China have not set up detente medical course and systematic training programs for medical students. So it is urgent to implant palliative medicine in the existing medical education system. Since palliative care is underdeveloped in China, patients and their family members are unfamiliar with it, the PCC consultation actually serve as a vivid case-based teaching for not only primary care physicians, but also patient's family members. Additionally, our PCC team members actively give lectures on palliative medicine for physicians in hospital, and they are also the key persons who educate publics on various medias. We believe the expertise of PCC team should play a crucial part in improving awareness of HPC of the whole society.

The results of survey in physicians indicate the benefits that patients, family and primary care team get from palliative care consultation. The effectiveness of palliative care consultation in a teaching hospital like PUMCH may promote the development of palliative care cross the China. Future actions should be taken for early interventions of palliative care in order to reduce ICU admissions, hospital re-admissions, and healthcare costs.¹⁸

There exists limitation in the study. The participants in survey were professionals who had been reached by the palliative care service, either as a bystander or a care provider, which can cause bias. The sample size and the representiveness should be considered when interpreting the results. Future studies should be carried out with increased number of participants in an extended targeted population in hospital, including physicians who have never applied for PCC as well as patients and their families, which will help us to take appropriate measures to improve the quality of PCC service in the hospital.

To conclude, taking good care of beginning and the end of life is related to the value orientation of medicine and the civilization of society. It is also an important part of the national social security system. Palliative care consultation service in a general hospital is efficacious and acclaimed, which can improve the quality of patient care, disseminate the concepts of palliative care, and boost the PCC providers' confidence. More efforts should be put to foster better accessibility, which calls for joint actions of the medical societies, the government and the general publics.

REFERENCE

1. Kang L, Zhu ML, Liu XH. Hospice and palliative care. *Chin J Geriatr* 2012; 31(6):538-40. doi:10.3760/cma.j.issn.0254-9026.2012.06.030.
2. Cook D, Rucker G. Dying with dignity in the intensive care unit. *N Engl J Med* 2014; 370(26):2506-14. doi:10.1056/NEJMr1208795.
3. Bates T, Hoy AM, Clarke OG, et al. The St Thomas Hospital terminal care support team. A new concept of hospice care. *Lancet* 1981; 1(8231):1201-3.
4. Morrison RS, Maroney-Galin C, Kralovec PD, et al. The growth of palliative care programs in United States hospitals. *J Palliat Med* 2005; 8(6):1127-34. doi:10.1089/jpm.2005.8.1127.
5. Cao Jijun, Yan Weiqi. The population enjoying palliative care in China is less than 1%. *Guangming Daily*, 2015-11-30: A6. http://epaper.gmw.cn/gmrb/html/2015-11/30/nw.D110000gmrb_20151130_1-06.htm?div=-1. Accessed December 15, 2018.
6. Smith TJ, Temin S, Alesi ER, et al. American Society of Clinical Oncology provisional clinical opinion: the integration of palliative care into standard oncology care. *J Clin Oncol* 2012; 30(8):880-7. doi:10.1200/JCO.2011.38.5161.
7. Shin J, Temel J. Integrating palliative care: when and how. *Curr Opin Pulm Med* 2013; 19(4):344-9. doi:10.1097/MCP.0b013e3283620e76.
8. Peppercorn JM, Smith TJ, Helft PR, et al. American Society of Clinical Oncology statement: toward individualized care for patients with advanced cancer. *J Clin Oncol* 2011; 29(6):755-60. doi:10.1200/JCO.2010.33.1744.
9. Jack B, Hillier V, Williams A, Oldham J. Hospital based palliative care teams improve the symptoms of cancer patients. *Palliat Med* 2003; 17(6):498-502. doi:10.1191/0269216303pm794oa.
10. Pan CX, Morrison RS, Meier DE, et al. How prevalent are hospital-based palliative care programs? Status report and future directions. *J Palliat Med* 2001; 4(3):315-24. doi:10.1089/109662101753123922.
11. Firn J, Preston N, Walshe C. What are the views of hospital-based generalist palliative care professionals on what facilitates or hinders collaboration with in-patient specialist palliative care teams? A systematically constructed narrative synthesis. *Palliat Med* 2016; 30(3):240-56. doi:10.1177/0269216315615483.
12. Hanks GW, Robbins M, Sharp D, et al. The imPaCT study: a randomized controlled trial to evaluate a hos-

- pital palliative care team. *Br J Cancer* 2002; 87(7): 733-9. doi:10.1038/sj.bjc.6600522.
13. Rabow MW, Dibble SL, Pantilat SZ, et al. The comprehensive care team: a controlled trial of outpatient palliative medicine consultation. *Arch Intern Med* 2004; 164(1):83-91. doi:10.1001/archinte.164.1.83.
14. May P, Garrido MM, Cassel JB, et al. Cost analysis of a prospective multi-site cohort study of palliative care consultation teams for adults with advanced cancer: Where do cost-savings come from? *Palliat Med* 2017; 31(4):378-86. doi:10.1177/0269216317690098.
15. Vinant P, Joffin I, Serresse L, et al. Integration and activity of hospital-based palliative care consultation teams: the INSIGHT multicentric cohort study. *BMC Palliat Care* 2017; 16(1):36. doi:10.1186/s12904-017-0209-9.
16. Higginson IJ, Finlay I, Goodwin DM, et al. Do hospital-based palliative teams improve care for patients or families at the end of life? *J Pain Symptom Manage* 2002; 23(2):96-106. doi:10.1016/S0885-3924(01)00406-7.
17. Rubens M, Ramamoorthy V, Saxena A, et al. Palliative care consultation trends among hospitalized patients with advanced cancer in the United States, 2005 to 2014. *Am J Hospice Palliative Med* 2018; 1049909118809975. doi:10.1177/1049909118809975.
18. Blue B, Vegunta R, Rodin M, et al. (July 20, 2018) Impact of an inpatient palliative care consultation in terminally ill cancer patients. *Cureus* 10(7):e3016. doi:10.7759/cureus.3016.