ORIGINAL ARTICLE

Effect of Dihydrotestosterone on Costimulatory Molecules in a Mouse Model of Graves' Disease

Lianye Liu^{1,2}, Bingyin Shi¹, Fengyi Zhao¹, Peng Hou¹,
Shu Liu¹, Xiaomei Liu^{1,3}, Liping Wu^{1*}

¹Department of Endocrinology, The First Affiliated Hospital of
Xi'an Jiaotong University, Xi'an 710061, China

²Department of Endocrinology, Weinan Central Hospital,
Weinan, Shaanxi 714000, China

³Department of Endocrinology, Nanjing First Hospital,
Nanjing Medical University, Nanjing 210001, China

Key words: Graves' disease; 5 α -dihydrotestosterone; costimulatory molecules; inhibitory molecules

Objective Graves' disease is the most common autoimmune thyroid disease and its prevalence and clinical manifestations are disparate between females and males. Costimulatory molecules play an essential role in regulating autoimmune responses. The objective of this study was to determine if expression of inhibitory molecules was correlated with treatment by dihydrotestosterone (DHT) in an *in vivo* BALB/c mouse model of experimental autoimmune Graves' disease.

Methods Female BALB/c mice were immunized three times with thyroid stimulating hormone receptor A-subunit encoded by adenovirus to establish a Graves' disease model. Three different doses of DHT or a matching placebo were administered by implantation of slow-release pellets a week before the first immunization. Four weeks after the third immunization, the mice were euthanatized, and then the spleen and thymus were removed. Total thyroxine and free thyroxine levels in serum of mice were detected using a radioimmunoassay kit. Real-time polymerase chain reaction was performed to estimate the expression of costimulatory molecules in lymphocytes from the spleen and thymus. Flow cytometry was used to analyze the percentage of CD4⁺ T cells in splenic lymphocytes. Quantitative data were compared with unpaired *t*-tests. Correlation between two variables was analyzed using Analysis of Variance.

Results Treatment with DHT can dramatically reduce total thyroxine and free thyroxine levels. Higher expression of programmed death-1 was found in the spleen of Graves' disease mice receiving 5 mg of DHT

treatment (0.635 ± 0.296 vs. 0.327 ± 0.212 ; $t=2.714$, $P=0.014$), similarly, T-cell immunoglobulin domain and mucin domain 3 (TIM-3) in both the spleen (1.004 ± 0.338 vs. 0.646 ± 0.314 ; $t=2.205$, $P=0.022$) and the thymus (0.263 ± 0.127 vs. 0.120 ± 0.076 ; $t=3.221$, $P=0.004$) also increased after 5 mg of DHT treatment compared with the parallel placebo model mice. Moreover, the percentage of CD4⁺T cells declined in the splenic lymphocytes of Graves' disease mice treated with 5 mg of DHT ($19.90\% \pm 3.985\%$ vs. $24.05\% \pm 2.587\%$; $t=2.804$, $P=0.012$). A significant negative association was observed between expression of TIM-3 in the spleen and serum levels of total thyroxine ($r=-0.7106$, $P=0.014$) as well as free thyroxine ($r=-0.6542$, $P=0.029$).

Conclusion This study demonstrates that DHT can ameliorate experimental autoimmune Graves' disease, which may occur by up-regulating expression of programmed death-1 and TIM-3 and inhibiting development of CD4⁺T cells.

GRAVES' disease (GD) is the most common autoimmune thyroid disease and displays significant gender-related differences in prevalence and clinical feature. It has been established that GD is a multifactorial disease caused by a complex interaction between genetic and environmental factors, which results in the loss of immune tolerance to thyroid antigens, leading to the initiation of immune reactions against the thyroid.^[1] GD arises from circulating antibodies that mimic the action of thyroid stimulating hormone (TSH) by binding to and activating the TSH receptor (TSHR), causing increased synthesis and release of thyroid hormones and subsequent hyperthyroidism.^[1]

The course of autoimmune disorders involves immune cells such as T lymphocytes, B lymphocytes and antigen presenting cells.^[2] The full activation of T cells resulting in proliferation and cytokine secretion requires at least two distinct signals. The first signal is antigen-specific, while the second costimulatory signal is pivotal for full activation of the naïve T lymphocyte.^[3] Costimulatory molecules also play a role in the pathogenesis of GD.^[4] The immune system possesses various mechanisms to inhibit harmful activation of immune cells. One mechanism is to maintain a balance between stimulatory and inhibitory receptors on lymphocytes.^[5] In the absence of the second costimulatory signal, some T lymphocytes will become unresponsive to antigen stimulation.^[3] Costimulatory molecules are distinctly classified into stimulatory and inhibitory receptors, depending on the effect of the signal on T cell activation.

The CD28 family contains four related members: CD28, cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), inducible T cell costimulatory (ICOS) and programmed death-1 (PD-1).^[3] CD28 is a stimulatory receptor for T cell proliferation, survival and produc-

tion of interleukin-2 (IL-2).^[6] A study has shown a remarkable increase in the percentage of lymphocytes expressing CD28 in the peripheral blood of children, adolescents and adults^[7] with hyperthyroidism, which may indicate that this abnormality is associated with development of autoimmune disorders. CTLA-4 is a negative regulator of T cell activation. It is expressed by activated T cells and is essential for maintaining tolerance to self-antigens in autoimmune disorders.^[8] ICOS is similar to CD28 in function.^[9] ICOS gene is expressed in resting T cells and the expression increases following T cell activation induced by costimulatory CD28.^[10] PD-1, a member of CD28/B7 superfamily of inhibitory receptors, is expressed on activated CD4⁺T cells, CD8⁺T cells and antigen presenting cells to modulate T cell tolerance.^[11] PD-1 exerts an inhibitory signal on activated T cells similar to CTLA-4.^[12] B and T lymphocyte attenuator (BTLA) is also an inhibitory receptor controlling activation of T cells and has effects similar to CTLA-4 and PD-1.^[13] The ligand for BTLA is herpes virus-entry mediator—a tumor necrosis factor receptor family protein, and the interaction between BTLA and herpes virus-entry mediator inhibits activation of T lymphocytes.^[14]

T cell immunoglobulin domain and mucin domain 3 (TIM-3), a member of the TIM family, was first identified as a surface molecule expressed on CD4⁺T helper type 1 (Th1) and on CD8⁺T cytotoxic type 1 (Tc1) cells.^[15] The TIM family includes eight members (TIM1 through TIM8) in mice, but only TIM1, TIM-3 and TIM4 are expressed in humans. The lectin galectin 9 (Gal9) was then discovered as a TIM-3 ligand. The interaction of Gal9 and TIM-3 on Th1 cells has been shown to contribute to cell death.^[16] Thus, TIM-3 came to be revealed as a negative regulatory molecule important for inhibiting immune responses driven by Th1 and Tc1.

CD154 (CD40 ligand, CD40L) is another surface

protein expressed predominantly by CD4⁺ T cells.^[17] The interaction between CD154 and CD40 stimulates B cell proliferation and secretion of immunoglobulins.^[18] CD154 has been shown to be an essential regulator of immune inflammatory responses in autoimmune diseases, and thus increases the severity of autoimmune diseases.^[19] Expression of OX40 (CD134), a member of the tumor necrosis factor receptor superfamily, is restricted to T cells.^[20] Interaction of OX40/OX40 ligand (OX40L) plays a key role in CD4⁺ Th1 cell responses^[20] and proliferation of memory T cells.^[21]

In many autoimmune diseases, sex hormones play a significant role. Numerous autoimmune disorders including thyroiditis, rheumatoid arthritis, systemic sclerosis and systemic lupus erythematosus affect more women than men.^[22]

Studies have revealed that androgen can provide protection against autoimmune thyroiditis.^[23, 24] Observers have found that GD is caused by both Th1 and Th2 immune responses,^[25] and may be related by similar autoimmune processes with thyroiditis.^[26] In our previous study, we used dihydrotestosterone (DHT) to pretreat female BALB/c mice with experimental autoimmune Graves' disease (EAGD) induced by adenovirus Ad-TSHR289. And the severity of EAGD was ameliorated after DHT treatment.^[27]

To further understand the influence of androgen on costimulatory molecules in T cells, we utilized mice with EAGD, which is an established model of hyperthyroidism.^[28] Induced by immunization with Ad-TSHR289 expressed in adenovirus, EAGD is characterized by the production of autoantibodies against TSHR and elevated levels of pro-inflammatory cytokines.^[29] We evaluated the effect of DHT on the expression of four stimulatory molecules (CD28, ICOS, OX40, and CD154) and four inhibitory molecules (CTLA-4, BTLA, PD-1, and TIM-3) in lymphocytes from the spleen and thymus of the mice with EAGD, using quantitative real-time PCR. We also investigated the percentage of CD4⁺ T cells in splenocytes by flow cytometry.

MATERIALS AND METHODS

Mice and treatment

Ninety-six healthy female 6–8-week-old BALB/c mice (weighting 18.38±0.22 g) housed in specific pathogen-free conditions were purchased from the Laboratory Animal Center (Xi'an Jiaotong University School of Medicine, China). All mice were maintained under

constant temperature (23°C) and humidity (60%) and 12-h-light and 12-h-dark cycles. Water and standard lab chow were available ad libitum. The 96 mice were randomly divided into eight groups, 12 in each group, 7 groups of which were immunized with adenovirus Ad-TSHR289 to establish EAGD model. This experiment was approved by the Animal Care and Use Committee of Xi'an Jiaotong University.

Ninety-day release pellets of DHT at doses of 1.5, 5 and 15 mg and matching placebos were purchased from Innovative Research of America (Sarasota, FL, USA). For six immunized groups, a week before immunization, a DHT-containing pellet or a placebo pellet was implanted under the skin of the neck^[30] of the mouse who was anesthetized with 10% ketamine (0.03–0.05 ml/10 g).

Immunization of mice with Ad-TSHR289 and sampling

Ad-TSHR289 was constructed as previously described.^[28] In brief, Ad-TSHR289 was propagated in HEK293 cells and purified by ion-exchange column chromatography. Immunization with Ad-TSHR289 was performed according to the method described previously.^[31] Each mouse in the immunized groups was injected intramuscularly with Ad-TSHR289 in 50 µl of phosphate buffered saline containing 10⁹ virus particles. The control mice were injected with null adenovirus. Immunizations were performed three times at three-week intervals.

Four weeks after the third immunization, blood was collected to measure thyroid function and autoantibodies, and after the mice were euthanatized, thyroid tissues, spleen and thymus were removed. The thyroid glands were fixed in 10% formalin for histological examination. Part of each spleen and thymus was rapidly frozen in liquid nitrogen and processed for RNA extraction and real-time polymerase chain reaction (PCR) as described below. The remaining portion of each spleen was used for analyzing percentage of CD4⁺ T cells by flow cytometry.

Thyroid function and autoantibodies

Total thyroxine (TT4) and free thyroxine (FT4) levels in undiluted serum of mice (50 µl) were detected using a radioimmunoassay kit (Beijing Atomic Hi-Tech Co. Ltd., Beijing, China), according to the manufacturer's protocols.

The levels of thyrotrophin receptor antibodies in undiluted serum (100 µl) and diluted serum (50 µl and 35 µl) were measured using a commercial kit (Medi-

pan GmbH, Berlin, Germany) based on the manufacturer's protocols, and thyroid stimulating hormone (TSH)-binding inhibition (TBI) was analyzed using the formula=[1-(TSH binding in the presence of test serum-nonspecific binding)/(TSH binding in the presence of normal serum-nonspecific binding)] $\times$ 100.^[28]

Thyroid histology

Thyroid tissues were fixed with 10% formalin in phosphate buffered saline. Tissues were embedded in paraffin and 5 mm thick sections were prepared and stained with hematoxylin and eosin.

RNA extraction and quantitative real-time PCR

Total RNA was extracted from each spleen and thymus using RNAiso Plus (TaKaRa Biotechnology Co. LTD, Dalian, China), according to the manufacturer's instructions. Reverse transcription (RT) reactions were performed using a PrimeScript RT reagent kit (TaKaRa). RNA transcripts were primed for transcription using both oligo-dT primers (TaKaRa) at a final concentration of 25 pmol/L and six random primers at 50 pmol/L. Real-time PCR reactions were performed on a CFX96 Real-Time System (C1000 Thermal cycler, Bio-Rad Laboratories, Hercules, CA, USA). All primers were synthesized by TaKaRa, and primer sequences for amplifying the following molecules, such as CTLA-4,^[32] BTLA,^[33] PD-1,^[34] TIM-3,^[35] CD28,^[36] ICOS,^[37] OX40,^[38] CD154,^[39] and GAPDH,^[36] are listed in **Table 1**. Amplification conditions were 95°C for 30 s, followed by 40 cycles of 5 s at 95°C, 30 s at 60°C, and 5 s at 60°C.

Preparation of cell suspension and flow cytometry

The resected spleens were gently dispersed into a sin-

gle-cell suspension through a nylon mesh. After red blood cells were lysed with a red blood cell lysis buffer (Tiangen Biotech Co. Ltd., Beijing, China), a single-cell suspension was prepared in RPMI-1640 (HyClone, Logan, USA) containing 10% heat-inactivated fetal calf serum (Hangzhou Sijiqing Biological Engineering Materials Co. Ltd., Hangzhou, China), at a final concentration of 1×10^7 cells/ml. Splenic cells from mice were immunostained with FITC-conjugated anti-mouse CD4 using a mouse regulatory T cell staining kit (eBioscience, San Diego, USA), following the manufacturer's protocol. The samples were analyzed using a FACSCalibur flow cytometer (BD, Franklin Lakes, USA) and the results were analyzed.

Statistical analysis

Quantitative data were presented as the means $\pm$ standard deviation (SD), and compared with unpaired *t*-tests. Correlation between two variables was analyzed using Analysis of Variance (ANOVA). Qualitative data were expressed as percent. A *P*-value <0.05 was considered to be statistically significant. Statistical analyses were performed using Graph Pad Prism software program 6 (GraphPad Software, Inc. La Jolla, Cal., USA).

RESULTS

Effect of DHT on TBI and thyroid function

Twelve mice died of hyperthyroidism during the experiment (**Table 2**).

As shown in **Table 2**, treatment with DHT can dramatically reduce TT4 and FT4 levels. A reduction for TBI obtained in 100 μ l of undiluted serum was also found in the DHT treated mice.

Table 1. Primer sequences for PCR amplification

Genes	Forward primers	Reverse primers	Fragment size (bp)
CTLA-4	5'-TGGGTTCAAACACATCTCAAGGC-3'	5'-TGTCGTGGCACAGACCTCAG-3'	349
BTLA	5'-GCCATTCAAGTAACCATCCATGTG-3'	5'-GCAAGGTGTAAAGCAGCCAAGTC-3'	150
PD-1	5'-CCGCTTCCAGATCATACAG-3'	5'-CTCTGGCCTCTGACATACTTG-3'	329
TIM-3	5'-AGCCCATGTGGAAATTTTG-3'	5'-CTCCAAGAACCCTAACCACG-3'	95
CD28	5'-GTGGTTGCTGGAGTCCGTT-3'	5'-GCTGGTAAGGCTTTCGAGTG-3'	151
ICOS	5'-GTCAAACAACAGCGTCTC-3'	5'-TGCAAATATCTCCACTAA-3'	129
OX40	5'-GTAGACCAGGCACCCAACC-3'	5'-GGCCAGACTGTGGTGGATTGG-3'	268
CD154	5'-ACGTTGTAAGCGAAGCCAAC-3'	5'-TATCCTTTCTTGCCCACTG-3'	61
GAPDH	5'-CATGACCACAGTCCATGCCAT-3'	5'-TCAGATCCACGACGACACATT-3'	215

CTLA-4: cytotoxic T-lymphocyte-associated protein 4; BTLA: B and T lymphocyte attenuator; PD-1: programmed death-1; TIM-3: T cell immunoglobulin domain and mucin domain 3; ICOS: inducible T cell costimulatory.

Table 2. TBI and serum levels of thyroid hormones in mice[§]

Groups	<i>n</i>	TBI ^a (%)	TT4 (nmol/L)	FT4 (pmol/L)
Control group	12	0.67±6.91	38.22±10.18	4.13±2.02
EAGD group	10	80.30±11.31	159.33±76.58	18.43±11.21
<i>t</i> value		20.31	5.446	4.354
<i>P</i> value		<0.0001	<0.0001	0.0003
Placebo _{1.5 mg DHT}	10	79.10±11.13	130.37±62.03	15.57±12.26
1.5 mg DHT group	11	70.36±10.01	75.29±35.78	11.09±5.02
<i>t</i> value		1.894	2.524	1.116
<i>P</i> value		0.07	0.02	0.28
Placebo _{5 mg DHT}	10	81.50±7.75	154.31±71.69	16.69±12.24
5 mg DHT group	11	70.00±20.09	78.12±32.18	6.831±3.02
<i>t</i> value		1.696	4.962	2.554
<i>P</i> value		0.11	0.01	0.02
Placebo _{15 mg DHT}	9	80.11±11.24	136.81±72.20	17.89±12.16
15 mg DHT group	11	69.82±12.81	75.80±53.15	6.90±4.38
<i>t</i> value		1.972	2.176	2.799
<i>P</i> value		0.06	0.04	0.01

§: Plus-minus values are means±SD.

TBI^a: thyroid stimulating hormone-binding inhibition measured in 100 µl of undiluted serum; TT4: total thyroxine; FT4: free thyroxine; DHT: dihydrotestosterone. Control group: normal mice injected with a null adenovirus; EAGD group: mice immunized with Ad-TSHR289; Placebo groups: mice immunized with Ad-TSHR289 and treated with matching placebo; DHT groups: mice immunized with Ad-TSHR289 and treated with DHT.

Effect of DHT on thyroid histology

Compared to the control mice, all hyperthyroid BALB/c mice had enlarged thyroid glands and larger thyroid follicles that intruded into follicular lumen. No lymphocytes infiltrating into the thyroid were observed. Compared with the placebo mice, the DHT treated mice had smaller thyroid follicles and fewer thyroid follicular cells (**Figure 1**).

Effects of DHT on mRNA expressions of costimulatory molecule genes in the spleen and the thymus

In the spleen, real-time PCR results showed compared to the controls the mRNA expressions of PD-1 ($t=2.624$, $P=0.035$) and TIM-3 ($t=2.669$, $P=0.015$) in the immunized mice were significantly decreased, and expression of CD154 increased ($t=2.473$, $P=0.023$) (**Figure 2A**). There were no significant differences were observed in the expression of CTLA-4, BTLA, CD28, ICOS and OX40 in the spleen between the immunized mice and controls (**Figure 2A**).

We then examined changes in the expression of costimulatory molecules in the spleen of mice after treatment with DHT. The results showed the increase in PD-1 ($t=2.714$, $P=0.014$) as well as TIM-3 mRNA ex-

pressions ($t=2.205$, $P=0.022$) in the spleen was greater after 5 mg of DHT treatment than that of the placebo groups. Finally, we did not find any significant differences in mRNA levels of CTLA-4, BTLA, CD28, ICOS, OX40 and CD154 of the spleen between 5 mg DHT treated

Figure 1. Thyroid histological pictures of female BALB/c mice with experimental autoimmune Graves' disease.

Histological examination shows compared with the mice pretreated with 5 α -DHT, the mice in the placebo group have larger thyroid follicles with epithelial cells intruding into the follicular lumen, and more follicular cells. HE staining.

mice and placebo treated mice (**Figure 2A**).

In the thymus, a significant decrease was detected in the expression of TIM-3 mRNA in the immunized BALB/c mice compared to the control mice ($t=2.293$, $P=0.033$, **Figure 2B**). In contrast, a significant increase was observed in the expression of ICOS mRNA ($t=4.831$, $P=0.0008$), but no significant changes were seen in PD-1, CTLA-4, BTLA, CD28, OX40, and CD154 expressions after immunization ($P>0.05$, **Figure 2B**). 5 mg ($t=3.221$, $P=0.004$, **Figure 2B**) and 15 mg DHT (0.210 ± 0.058 vs. 0.132 ± 0.049 ; $t=3.190$, $P=0.005$)

had a significant effect on TIM-3 mRNA expression. No significant differences were found in expression of PD-1, CTLA-4, and BTLA mRNAs between the 5 mg DHT group and the placebo group. With regard to the stimulating molecules, no significant changes were observed in CD28, ICOS, OX40 and CD154 after treatment with 5 mg DHT in the thymus.

Effects of different doses of DHT on mRNA expressions of costimulatory molecule genes

The effects of three doses of DHT on mRNA expression

Figure 2. Comparisons of mRNA expressions of costimulatory molecule genes in the spleen (A) and the thymus (B).

Six to eight weeks old female BALB/c mice were injected with 10^9 particles of Ad-TSHR289 diluted in 50 ml PBS for 3 times at one-week interval (EAGD group). Meanwhile, BALB/c mice receiving null adenovirus injection served as the Control (Con). A week before immunization, mouse was implanted with a pellet containing 5 mg DHT (DHT group) or matching placebo (Placebo group). The spleen and the thymus were obtained respectively 4 weeks after the third immunization. The extracted total RNAs from the spleen and the thymus were used to analyze mRNA expressions of costimulatory molecule genes with quantitative real-time PCR assay. Data are expressed as the means $\pm$ SD. Con ($n=12$), EAGD ($n=10$), Placebo ($n=10$), DHT ($n=11$). *** $P<0.001$, ** $P<0.01$, * $P<0.05$.

levels of costimulatory molecules in the spleen and thymus were not exactly the same and not dose-dependent. The increase in PD-1 and TIM-3 mRNA expressions in the spleen of model mice pretreated with 5 mg of DHT was greater, with a remarkable increase in TIM-3 mRNA of the mice exposed to a lower dosage of DHT (1.5 mg). But no effect of 15 mg DHT was observed on the expressions of PD-1, CTLA-4, TIM-3, and BTLA mRNA. In the thymus, both 5 mg and 15 mg DHT had an effective influence on TIM-3 expression. No significant differences were found in expression of four inhibitory costimulatory molecules after 1.5 mg DHT treatment. As to the stimulating molecules, no available changes were observed in CD28, ICOS, OX40 and CD154 after treatment with three doses of DHT in both the spleen and thymus.

Effect of DHT on percentage of CD4⁺ T cells in the splenic lymphocytes

Flow cytometry showed that the percentage of CD4⁺ T cells was dramatically less in the immunized mice supplemented with 5 mg DHT than that of the placebo EAGD mice ($t=2.804$, $P=0.012$; **Figure 3**).

Correlation between TBI/thyroid function and expression of costimulatory molecules in EAGD mice treated with 5 mg DHT

Linear regression analysis revealed excellent correlation between serum TT4 and FT4 levels in the EAGD mice treated with 5 mg of DHT ($r=0.9494$, $P<0.0001$).

TBI was negatively associated with expression of CTLA-4 ($r=-0.7962$, $P=0.003$) and BTLA ($r=-0.7944$, $P=0.004$) in the thymus (**Figure 4**). Correlations of TBI with PD-1 ($r=-0.2356$, $P>0.05$) and TIM-3 ($r=0.2392$, $P>0.05$) in the thymus were not statistically significant (**Figure 4**).

Serum TT4 level was negatively associated with expression of TIM-3 in the spleen ($r=-0.7106$, $P=0.014$; **Figure 5**). In addition, a prominent negative association was found between serum level of FT4 and expression of TIM-3 in the spleen ($r=-0.6542$, $P=0.029$; **Figure 5**).

DISCUSSION

The costimulatory signal is required for full activation of T lymphocytes.^[40] Multiple studies indicate that a deficiency or mutation of inhibitory molecules contributes to the development of autoimmune diseases in

Figure 3. Percentage of CD4⁺ cells in the splenic lymphocytes of mice analyzed by flow cytometry.

Placebo ($n=10$): Ad-TSHR289 immunized mice treated with matching placebo; DHT ($n=11$): Ad-TSHR289 immunized mice treated with 5 mg of DHT. Data are expressed as mean $\pm$ SD. * $P<0.05$.

mice as well as humans, suggesting that the negative signals from inhibitory receptors play an important role in inhibiting autoimmunity. Inhibitory molecules including CTLA-4, BTLA, PD-1 and TIM-3 can prevent immune responses to autoantigens. In the present study, we explored mRNA expressions of these inhibitory receptors as well as the stimulatory molecules CD28, ICOS, OX40, and CD154.

Our results showed a significant decrease in the expression of PD-1 in the spleen and TIM-3 in both the spleen and thymus after immunization; in contrast, a significant increase was seen in the expression of CD154 in the spleen and ICOS in the thymus compared to the control mice. Polymorphisms in the ICOS and ICOSL genes have been proved to be associated with the pathogenesis of autoimmune thyroid diseases.^[41] Recently, another study has shown that CD154 (CD40 ligand) can up-regulate the expression of vascular cell adhesion molecule 1 and E-selectin, and may play a role in patients with Graves' orbitopathy.^[42]

We found that expression of PD-1 was up-regulated in the spleen, while TIM-3 was up-regulated in both the spleen and thymus of the mice treated with 5 mg

Figure 4. Correlations between mRNA expressions of *CTLA-4* (A), *BTLA* (B), *PD-1* (C), and *TIM-3* (D) in the thymus and TBI analyzed by linear regression.

Figure 5. Correlations between TIM-3 mRNA expression in the spleen (A, B) and the thymus (C, D) and TT4/FT4 in serum analyzed by linear regression.

of DHT. We also discovered that the expression of TIM-3 not only increased in the spleen of EAGD mice after treatment with 1.5 mg of DHT but also in the thymus of EAGD mice treated with 15 mg of DHT. The effect of DHT on BALB/c mice model was not dose dependent. The influence of the dose of DHT 5 mg was more remarkable. PD-1 and its ligands (PD-L1 and PD-L2) regulate suppressive T cell activation that mediates mechanisms of tolerance and immune homeostasis. Ligation of PD-1 as a secondary signal to T cells in com-

bination with T cell receptor signaling contributes to a negative signal.^[43] TIM-3 has been reported to play a critical role in the maintenance of peripheral tolerance. A study revealed that the TIM-3 pathway blocks the development of a mouse model with multiple sclerosis by negatively regulating the production of cytokines in CD4⁺ and CD8⁺ T cells.^[44] There is also increasing evidence that TIM-3 influences chronic autoimmune diseases. Galectin-9, a TIM-3 ligand, induces Th1 cell death by interacting with TIM-3, and galectin-9-de-

ficient mice become susceptible to collagen-induced arthritis.^[45] As mentioned above, we have shown an effect of DHT on PD-1 and TIM-3. To further understand the effects of DHT on the interactions between PD-1 and PD-L1/PD-L2, as well as the interaction between TIM-3 and Galectin-9, we plan to investigate PD-L1, PD-L2 and Galectin-9 in subsequent studies. The Galectin-9 signaling pathway in particular should be investigated because of the significant negative correlations we observed between TIM-3 expression in the spleen and serum levels of FT4/TT4.

A recent study showed that the percentage of CD4⁺ T cells was significantly decreased in the testis of an experimental allergic orchitis rat model supplemented with testosterone.^[46] Consistent with this finding, our investigation showed a decrease in the percentage of CD4⁺ T cells in the spleen of BALB/c mice implanted with a pellet of 5 mg DHT. Furthermore, we observed a significant negative association between serum levels of TT4/FT4 and expression of TIM-3 in the spleen. Our results thus suggest that DHT might alleviate EAGD *via* up-regulating PD-1 and TIM-3 expression and inhibiting CD4⁺ T cell development. In our previous study, we researched the influence of DHT on cytokines in GD mice model by flow cytometry, and found the production of cytokines interferon- γ and interleukin-2 was decreased in DHT-pretreated mice.^[27] In this study, the effect of DHT on costimulatory molecules in GD mice model was explored by RT-PCR. In the next step, we will observe other mechanisms of DHT on costimulatory molecules in GD mice model using different methods, including flow cytometry.

In summary, administration of DHT significantly ameliorated the severity of GD in a female BALB/c mouse model possibly by enhancing the expression of PD-1 and TIM-3. These observations suggest that DHT may have therapeutic potential for the treatment of GD.

Conflict of Interests Statement

The authors declare no conflict of interests.

REFERENCES

- Menconi F, Marcocci C, Marino M. Diagnosis and classification of Graves' disease. *Autoimmun Rev* 2014; 4-5(13):398-402. doi: 10.1016/j.autrev.2014.01.013.
- Bossowski A, Stasiak-Barmuta A, Urban M. Relationship between CTLA-4 and CD28 molecule expression on T lymphocytes and stimulating and blocking autoantibodies to the TSH-receptor in children with Graves' disease. *Horm Res* 2005; 4(64):189-97. doi: 10.1159/000088875.
- Garapati VP, Lefranc MP. IMGT Colliers de Perles and IgSF domain standardization for T cell costimulatory activatory (CD28, ICOS) and inhibitory (CTLA-4, PDCD1 and BTLA) receptors. *Dev Comp Immunol* 2007; 10(31):1050-72. doi: 10.1016/j.dci.2007.01.008.
- Pawlak-Adamska E, Frydecka I, Bolanowski M, et al. CD28/CTLA-4/ICOS haplotypes confers susceptibility to Graves' disease and modulates clinical phenotype of disease. *Endocrine* 2017; 1(55):186-99. doi: 10.1007/s12020-016-1096-1.
- Watanabe N, Nakajima H. Coinhibitory molecules in autoimmune diseases. *Clin Dev Immunol* 2012; 2012:269756. doi: 10.1155/2012/269756.
- Turka LA, Ledbetter JA, Lee K, et al. CD28 is an inducible T cell surface antigen that transduces a proliferative signal in CD3⁺ mature thymocytes. *J Immunol* 1990; 5(144):1646-53.
- Matsuoka N, Eguchi K, Kawakami A, et al. Lack of B7-1/BB1 and B7-2/B70 expression on thyrocytes of patients with Graves' disease. Delivery of costimulatory signals from bystander professional antigen-presenting cells. *J Clin Endocrinol Metab* 1996; 11(81):4137-43. doi: 10.1210/jcem.81.11.8923872.
- Eagar TN, Karandikar NJ, Bluestone JA, et al. The role of CTLA-4 in induction and maintenance of peripheral T cell tolerance. *Eur J Immunol* 2002; 4(32):972-81. doi: 10.1002/1521-4141(200204)32:4<972::AID-IMMU972>3.0.CO;2-M.
- Hutloff A, Dittrich AM, Beier KC, et al. ICOS is an inducible T-cell co-stimulator structurally and functionally related to CD28. *Nature* 1999; 6716(397):263-6. doi: 10.1038/16717.
- Wassink L, Vieira PL, Smits HH, et al. ICOS expression by activated human Th cells is enhanced by IL-12 and IL-23: increased ICOS expression enhances the effector function of both Th1 and Th2 cells. *J Immunol* 2004; 3(173):1779-86. doi: 10.4049/jimmunol.173.3.1779.
- Ishida Y, Agata Y, Shibahara K, et al. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J* 1992; 11(11):3887-95.
- Carter LL, Carreno BM. Cytotoxic T-lymphocyte antigen-4 and programmed death-1 function as negative regulators of lymphocyte activation. *Immunol Res*

- 2003; 1(28):49-59. doi: 10.1385/IR:28:1:49.
13. Watanabe N, Gavrieli M, Sedy JR, et al. BTLA is a lymphocyte inhibitory receptor with similarities to CTLA-4 and PD-1. *Nat Immunol* 2003; 7(4):670-9. doi: 10.1038/ni944.
14. Sedy JR, Gavrieli M, Potter KG, et al. B and T lymphocyte attenuator regulates T cell activation through interaction with herpesvirus entry mediator. *Nat Immunol* 2005; 1(6):90-8. doi: 10.1038/ni1144.
15. Monney L, Sabatos CA, Gaglia JL, et al. Th1-specific cell surface protein Tim-3 regulates macrophage activation and severity of an autoimmune disease. *Nature* 2002; 415(6871):536-41. doi: 10.1038/415536a.
16. Zhu C, Anderson AC, Schubart A, et al. The Tim-3 ligand galectin-9 negatively regulates T helper type 1 immunity. *Nat Immunol* 2005; 12(6):1245-52. doi: 10.1038/ni1271.
17. Locksley RM, Killeen N, Lenardo MJ. The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell* 2001; 104(4):487-501.
18. Hollenbaugh D, Grosmaire LS, Kullas CD, et al. The human T cell antigen gp39, a member of the TNF gene family, is a ligand for the CD40 receptor: expression of a soluble form of gp39 with B cell co-stimulatory activity. *EMBO J* 1992; 11(11):4313-21.
19. Hassan GS, Merhi Y, Mourad WM. CD154 and its receptors in inflammatory vascular pathologies. *Trends Immunol* 2009; 4(30):165-72. doi: 10.1016/j.it.2009.01.004.
20. Watts TH. TNF/TNFR family members in costimulation of T cell responses. *Annu Rev Immunol* 2005; 23:23-68. doi: 10.1146/annurev.immunol.23.021704.115839.
21. Gramaglia I, Jember A, Pippig SD, et al. The OX40 co-stimulatory receptor determines the development of CD4 memory by regulating primary clonal expansion. *J Immunol* 2000; 165(6):3043-50.
22. Lumachi F, Basso SM, Orlando R. Cytokines, thyroid diseases and thyroid cancer. *Cytokine* 2010; 50(3):229-33. doi: 10.1016/j.cyto.2010.03.005.
23. Ansar AS, Young PR, Penhale WJ. Beneficial effect of testosterone in the treatment of chronic autoimmune thyroiditis in rats. *J Immunol* 1986; 136(1):143-7.
24. Fassler R, Dietrich H, Kromer G, et al. The role of testosterone in spontaneous autoimmune thyroiditis of Obese strain (OS) chickens. *J Autoimmun* 1988; 1(1):97-108.
25. Paschke R, Schuppert F, Taton M, et al. Intrathyroidal cytokine gene expression profiles in autoimmune thyroiditis. *J Endocrinol* 1994; 141(2):309-15.
26. Endo T, Kobayashi T. Immunization with thyroglobulin induces Graves'-like disease in mice. *J Endocrinol* 2009; 167(2):217-22. doi: 10.1677/JOE-09-0089.
27. Liu L, Wu L, Gao A, et al. The influence of dihydrotestosterone on the development of Graves' disease in female BALB/c mice. *Thyroid* 2016; 26(3):449-57. doi: 10.1089/thy.2015.0620.
28. Chen CR, Pichurin P, Nagayama Y, et al. The thyrotropin receptor autoantigen in Graves disease is the culprit as well as the victim. *J Clin Invest* 2003; 111(12):1897-904. doi: 10.1172/JCI17069.
29. Nagayama Y, Mizuguchi H, Hayakawa T, et al. Prevention of autoantibody-mediated Graves'-like hyperthyroidism in mice with IL-4, a Th2 cytokine. *J Immunol* 2003; 170(7):3522-7.
30. Mostafa S, Seamon V, Azzarolo AM. Influence of sex hormones and genetic predisposition in Sjogren's syndrome: a new clue to the immunopathogenesis of dry eye disease. *Exp Eye Res* 2012; 96(1):88-97. doi: 10.1016/j.exer.2011.12.016.
31. Wu L, Xun L, Yang J, et al. Induction of murine neonatal tolerance against Graves' disease using recombinant adenovirus expressing the TSH receptor A-subunit. *Endocrinology* 2011; 152(3):1165-71. doi: 10.1210/en.2010-0737.
32. Yessoufou A, Ple A, Moutairou K, et al. Docosahexaenoic acid reduces suppressive and migratory functions of CD4+CD25+ regulatory T-cells. *J Lipid Res* 2009; 50(12):2377-88. doi: 10.1194/jlr.M900101-JLR200.
33. Han LF, Qiu WM, Hu C, et al. Therapeutic effects of B and T lymphocyte attenuator extracellular domain and heat shock protein 70 antigen peptide on cervical cancer in mouse model. *Zhonghua Fu Chan Ke Za Zhi* 2010; 45(8):603-8.
34. Bishop KD, Harris JE, Mordes JP, et al. Depletion of the programmed death-1 receptor completely reverses established clonal anergy in CD4(+) T lymphocytes via an interleukin-2-dependent mechanism. *Cell Immunol* 2009; 256(1-2):86-91. doi: 10.1016/j.cellimm.2009.01.008.
35. Lech M, Susanti HE, Rommele C, et al. Quantitative expression of C-type lectin receptors in humans and mice. *Int J Mol Sci* 2012; 13(8):10113-31. doi: 10.3390/ijms130810113.
36. Teague TK, Tan C, Marino JH, et al. CD28 expression redefines thymocyte development during the pre-T to DP transition. *Int Immunol* 2010; 22(5):387-97. doi: 10.1093/intimm/dxq020.

37. Zang YS, Fang Z, Liu YA, et al. Repressor of GATA-3 can negatively regulate the expression of T cell cytokines through modulation on inducible costimulator. *Chin Med J (Engl)* 2012; 12(125):2188-94.
38. Tone Y, Kojima Y, Furuuchi K, et al. OX40 gene expression is up-regulated by chromatin remodeling in its promoter region containing Sp1/Sp3, YY1, and NF-kappa B binding sites. *J Immunol* 2007; 3(179):1760-7.
39. Vavassori S, Shi Y, Chen CC, et al. *In vivo* post-transcriptional regulation of CD154 in mouse CD4+ T cells. *Eur J Immunol* 2009; 8(39):2224-32. doi: 10.1002/eji.200839163.
40. Frauwirth KA, Thompson CB. Activation and inhibition of lymphocytes by costimulation. *J Clin Invest* 2002; 3(109):295-9. doi: 10.1172/JCI14941.
41. Yoshie N, Watanabe M, Inoue N, et al. Association of polymorphisms in the ICOS and ICOSL genes with the pathogenesis of autoimmune thyroid diseases. *Endocr J* 2016; 1(63):61-8. doi: 10.1507/endocrj.EJ15-0435.
42. Wang H, Zhu LS, Cheng JW, et al. CD40 ligand induces expression of vascular cell adhesion molecule 1 and E-selectin in orbital fibroblasts from patients with Graves' orbitopathy. *Graefes Arch Clin Exp Ophthalmol* 2015; 4(253):573-82. doi: 10.1007/s00417-014-2902-1.
43. Dong C, Nurieva RI, Prasad DV. Immune regulation by novel costimulatory molecules. *Immunol Res* 2003; 1(28):39-48. doi: 10.1385/IR:28:1:39.
44. Kaneyama T, Tomiki H, Tsugane S, et al. The TIM-3 pathway ameliorates Theiler's murine encephalomyelitis virus-induced demyelinating disease. *Int Immunol* 2014; 7(26):369-81. doi: 10.1093/intimm/dxt056.
45. Seki M, Oomizu S, Sakata KM, et al. Galectin-9 suppresses the generation of Th17, promotes the induction of regulatory T cells, and regulates experimental autoimmune arthritis. *Clin Immunol* 2008; 1(127):78-88. doi: 10.1016/j.clim.2008.01.006.
46. Fijak M, Schneider E, Klug J, et al. Testosterone replacement effectively inhibits the development of experimental autoimmune orchitis in rats: evidence for a direct role of testosterone on regulatory T cell expansion. *J Immunol* 2011; 9(186):5162-72. doi: 10.4049/jimmunol.1001958.