ORIGINAL ARTICLE

Linguistic Characteristics of Mandarin-Speaking Huntington's Disease Patients

Liling Dong, Caiyan Liu, Chenhui Mao, Shanshan Chu,
Jie Li, Xinying Huang, Jing Gao*

Department of Neurology, Peking Union Medical College Hospital,
Chinese Academy of Medical Sciences & Peking Union
Medical College, Beijing 100730, China

Key words: Huntington's disease; Alzheimer's disease; transcortical motor aphasia

Objective Linguistic problem is common in Huntington's disease (HD) patients. It has been studied before in native speakers of alphabetic languages, such as English. As a hieroglyphic language, Chinese differs from alphabetic languages in terms of phonology, morphology, semantics and syntax. We aimed to investigate the linguistic characteristics of manifest HD in native speakers of Mandarin. Meanwhile, we expected to explore the linguistic differences associated with cortical or subcortical pathology.

Methods Five HD patients and five Alzheimer's disease (AD) patients matched in age, gender, disease course and educational level were enrolled. All the participants were Mandarin native speakers. All finished history inquiry, physical examination, basic test, genetic test and neuropsychological assessment. Language evaluation was performed by Aphasia Battery of Chinese.

Results HD patients had a mean disease course of 5.4 ± 2.97 (range, 2-10) years. They showed a linguistic disorder close to transcortical motor aphasia. They exhibited prominent phonological impairment, as well as slight semantic and syntactic abnormality. Tonic errors were found in speech. Character structural errors and substitutions were detected in writing. In comparison, AD patients showed a more severe linguistic impairment, characterized by global aphasia with more semantic errors.

Conclusion Mandarin-speaking HD patients have a transcortical motor aphasia-like disturbance with prominent phonological impairment, whereas AD patients have a more severe global aphasia with salient semantic impairment.

Received October 12, 2019. Published online June 30, 2020.

*Corresponding author Tel: 86-10-69156371, Fax: 86-10-69156371, E-mail: gj107@163.com

This work was supported by grants from CAMS Innovation Fund for Medical Sciences (No. 2016-I2M-1-004), National Natural Science Foundation of China (No. 81550021), 13th Five-year National Key Research and Development Program of China (No. 2016YFC1306300), and the strategic priority research program (pilot study) "Biological basis of aging and therapeutic strategies" of the Chinese Academy of Sciences (XDPB10).

HUNTINGTON'S disease (HD) is an autosomal dominant neurodegenerative disorder due to an expansion of CAG repeats in the exon 1 of huntington (HTT) gene.^[1] It is characterized by motor disorder, cognitive decline and psychiatric symptom.^[2]

Linguistic problem is common in HD patients. It has been studied before in native speakers of alphabetic languages, such as English, French, Spanish, German, Portuguese and so on.^[3] Chinese is a kind of hieroglyphic language, which is completely different from alphabetic languages in terms of phonology, morphology, semantics and syntax. In this paper, we aimed to investigate the linguistic characteristics of Mandarin-speaking HD patients. Literature review indicates a complex and bidirectional interaction between cognitive and linguistic processes.^[4-7] As a result, we also illustrated their cognitive characteristics.

HD and Alzheimer's disease (AD) are representatives of subcortical and cortical dementia, respectively. HD is generally characterized by progressive degeneration of neostriatum, including caudate nucleus and putamen.^[8] However, the major pathology of AD is progressive deterioration of medial temporal, frontal and parietal cortices.^[9] In order to understand the linguistic differences associated with subcortical or cortical pathology, comparisons were made between HD and AD.

PATIENTS AND METHODS

Patients and enrollment

Five HD patients, together with five AD patients matched in age, gender, disease course and educational level, were recruited from the dementia and leukoencephalopathy clinic of Peking Union Medical College Hospital (PUMCH). All the participants were Mandarin native speakers. HD patients fulfilled the diagnostic criteria for HD issued in 1986.^[10] They all had an expansion of CAG repeats in HTT gene, ranging from 38 to 41. AD patients met the diagnostic criteria for probable AD dementia with evidence of AD pathophysiological process, according to the recommendations of National Institute on Aging and the Alzheimer's Association (NIA-AA) released in 2011.^[11] All the participants gave informed consent. And this study was approved by the ethical committee of PUMCH.

All the individuals finished history inquiry, physical examination, basic test, genetic test and neuropsychological assessment.

Basic and genetic tests

Basic test consisted of blood routine test, hepatic and renal function, thyroid function, homocysteine, folic acid, cobalamin, blood ammonia, rapid plasma reagin test, immune function, antibodies to Hu, Yo and Ri, electroencephalography and brain magnetic resonance imaging (MRI). In addition, five AD patients and one HD patient had cerebrospinal fluid test of A β 1-42, t-tau and p-tau. Five HD patients received CAG repeats measurement, and five AD patients took whole genome sequencing.

Neuropsychological test

Neuropsychological test consisted of screening test, multi-domain assessment and language evaluation. The latter was performed by Aphasia Battery of Chinese (ABC).^[12]

The screening test included mini-mental state exam (MMSE), Montreal cognitive assessment (Moca),^[13] activities of daily living (ADLs) and hospital anxiety and depression scale (HADS). Five HD patients received extra neuropsychiatric inventory (NPI).

The multi-domain battery involved four domains, such as executive, visuospatial, memory, reasoning and calculation domains. It is comprised of the following tests: digital symbol substitution task (DSST), trail making test part A (TMT-A), clock drawing, single and serial gesture imitation, figure copying, block design, Rey complex figure, paired associative learning (PAL), episodic memory, auditory verbal learning test (AVLT), similarity test and calculation.

Statistical analysis

Statistical analysis was carried out by SPSS (version 18.0). The demographic and neuropsychological data were displayed as mean $\pm$ standard deviation. A simple scatterplot was created to give an intuitive description of linguistic differences between HD and AD patients.

RESULTS

Clinical profile

HD patients had a mean disease course of 5.4 $\pm$ 2.97 (range, 2-10) years and a mean educational level of 10.4 $\pm$ 3.85 (range, 6-15) years. Two patients started with choreic movement of head or limbs. The cognitive, linguistic and psychiatric symptoms occurred within 3 to 8 years afterwards. Among the other 3 patients, motor disorder and neuropsychological symp-

toms occurred almost simultaneously. (**Tables 1, 2**)

AD patients had a mean disease course of 5.0 ± 1.87 (range, 3-8) years and a mean educational level of 13.4 ± 1.95 (range, 11-16) years. The initial symptom was memory loss. The linguistic symptom occurred 0 to 3 years afterwards. (**Tables 1, 2**)

Brain MRI

As demonstrated in **Table 2** and **Figure 1**, among the five HD patients, the axial MRI revealed an prominent atrophy of the head of caudate nucleus (HCN), with an enlarged anterior horn of lateral ventricle. However, the medially convex outline of HCN remained. Atrophy of frontal, temporal, parietal and occipital cortices was also detected. Compared with HD patients, AD patients showed no HCN atrophy and less cortical atrophy.

Cognitive profile

HD patients showed low scores in DSST, TMT-A, clock drawing, gesture imitation, as well as in similarity test and calculation, which suggested impairment in executive and reasoning domains. The scores in AVLT, recall of Rey complex figure, paired associative learning and episodic memory were even lower, suggesting more severe impairment in memory domain. In visuospatial domain, although they could finish block design test, the time spent was longer than expected. Comparatively, AD patients showed a more severe cognitive impairment, involving all the four domains. (**Tables 1 and 3**)

Linguistic profile

The linguistic feature in HD patients was close to transcortical motor aphasia (TCMA), called TCMA-like disturbance. Oral comprehension and repetition was slightly affected at the level of long complex sentence. Color naming was flawless, whereas picture naming, object naming and responsive naming were a bit affected. They showed a slight deficit in verbal category fluency test of animal and vegetable. They could finish the serial speech tasks well, such as counting up to 21 and reciting 12 Chinese zodiac animals. Reading was relatively intact. However, reading comprehension paralleled oral comprehension and was slightly impaired at the level of long complex sentence. (**Figure 2**)

HD patients showed a non-fluent spontaneous speech with decreased verbal output and grammatical complexity. Phonological error and speech dysprosody were obvious. The former was mainly manifested

Table 1. Demographic and neuropsychological data of HD and AD patients[§] (n=5)

Items	HD	AD
Gender (M/F)	3/2	3/2
Age on admission [§] (y/o)	61.0±6.82	55.4±5.32
Age of onset [§] (y/o)	55.6±9.21	50.4±5.59
Disease course [§] (yrs)	5.4±2.97	5.0±1.87
Educational level [§] (yrs)	10.4±3.85	13.4±1.95
MMSE [§] (score)	24.0±1.22	11.8±7.95
Moca [§] (score)	19.8±3.83	8.2±5.76
ADLs [§] (score)	23.8±2.49	32.0±16.05
HADS-A [§] (score)	1.0±1.41	4.2±3.03
HADS-D [§] (score)	1.8±2.49	4.2±4.02
NPI-frequency [§] (score)	5.0±5.48	-
NPI-severity [§] (score)	4.4±5.13	-
NPI-distress [§] (score)	5.6±6.73	-
DSST [§] (score)	20.2±4.87	12.0±19.65
TMT-A [§] (s)	121±23.82	171.4±95.78
Clock drawing [§] (score)	2.0±0.71	1.4±0.89
Single gesture [§] (score)	5.2±1.64	3.8±1.92
Serial gesture [§] (score)	1.4±0.89	0.4±0.55
Figure copying [§] (score)	9.8±0.45	6.6±4.22
Block design [§] (score)	4.7±2.51	2.3±3.90
Rey-copy (score)	17.0±0.00	13.0±7.31
Rey-recall [§] (score)	7.0±5.34	2.6±3.97
PAL-sum [§] (score)	5.0±1.77	0.8±1.79
PAL-N1 [§] (score)	1.2±0.57	0.2±0.45
PAL-N2 [§] (score)	1.9±0.42	0.3±0.67
PAL-N3 [§] (score)	1.9±1.08	0.3±0.67
Episodic memory [§] (score)	4.9±1.78	1.7±1.72
AVLT-I [§] (score)	13.0±6.28	5.4±3.78
AVLT-N1 [§] (score)	3.0±2.00	0.4±0.55
AVLT-N2 [§] (score)	4.2±1.92	2.6±2.07
AVLT-N3 [§] (score)	5.8±2.68	2.4±1.82
AVLT-N4 [§] (score)	3.0±2.83	0.2±0.45
AVLT-N5 [§] (score)	2.4±2.30	0
AVLT-N6 [§] (score)	1.4±1.67	0
AVLT-N7 [§] (score)	12.4±1.67	7.2±6.57
Similarity [§] (score)	12.2±6.38	3.0±3.67
Calculation [§] (score)	7.0±1.22	2.2±1.64

§: Plus-minus values are means±SD. HD: Huntington's disease; AD: Alzheimer's disease; M/F: male/female; y/o: years old; MMSE: mini mental state examination; Moca: Montreal cognitive assessment; ADLs: activities of daily living; HADS-A: anxiety section of Hospital Anxiety and Depression Scale; HADS-D: depression section of Hospital Anxiety and Depression Scale; NPI: neuropsychiatric inventory; DSST: digital symbol substitution task; TMT-A: trail making test part A; PAL: paired associative learning; AVLT: Auditory Verbal Learning Test; AVLT-I: sum of three immediate recalls; AVLT-N1: 1st immediate recall; AVLT-N2: 2nd immediate recall; AVLT-N3: 3rd immediate recall; AVLT-N4: short delayed recall; AVLT-N5: long delayed recall; AVLT-N6: cued recall; AVLT-N7: recognition.

Table 2. Clinical data of HD and AD patients

Case	Gender	Age (yrs)	Disease course (yrs)	Educational level (yrs)	CAG repeats	CSF test (pg/ml)		Brain MRI* (Grade)				
						A β	t-tau	HCN atrophy	FL atrophy	TL atrophy	PL atrophy	OL atrophy
1	Male	57	4	7	39			+	2	2	2	2
2	Male	67	2	6	41	346	50	+	2	1	2	1
3	Female	64	5	15	38			+	2	2	1	2
4	Male	51	10	13	40			+	2	1	2	1
5	Female	66	6	11	41			+	1	1	1	1
6	Female	47	4	14		213	306	-	1	1	1	0
7	Male	58	5	11		267	356	-	1	1	1	1
8	Male	61	5	14		122	425	-	1	2	1	0
9	Male	54	8	16		182	620	-	1	1	1	0
10	Female	57	3	12		140	437	-	1	2	1	0

Case 1-5: HD patients; Case 6-10: AD patients; CSF: cerebrospinal fluid; HCN: head of caudate nucleus; FL: frontal lobe; TL: temporal lobe; PL: parietal lobe; OL: occipital lobe.

*Cortical atrophy rating is based on the four-step Pasquier scale for global cortical atrophy.^[14] Grade 0: normal; Grade 1: widened sulci; Grade 2: widened sulci with volume loss of the gyri; Grade 3: knife blade atrophy.

Figure 1. Brain MRI of HD and AD patients.

Pictures A-B are from a 57-year-old male HD patient, who has a 4-year disease course and a 7-year education. Brain MRI demonstrates HCN atrophy, accompanied by moderate atrophy of frontal, parietal, temporal and occipital cortices. Pictures C-D are from a 58-year-old male AD patient, who has a 5-year disease course and an 11-year education. Brain MRI shows slight atrophy of frontal, parietal, temporal and occipital cortices, with HCN unaffected.

as the initial consonant substitution. YANG and DONG were pronounced as RANG and ZHONG, respectively. Pa-Ta-Ka became Ka-Ka-Ka. The latter was characterized by aperiodicity and irregular pitch fluctuation. Tonic errors also existed. The third tone was pronounced as the first tone. (**Figure 2**)

HD patients showed a worse writing impairment than other linguistic deficits. In spontaneous writing, one wrote nothing, one wrote two phrases, and only three wrote sentences. Most sentences had an intact grammatical structure. Some were lack of subjects or objects. In addition, there were plenty of character structural errors and character substitutions. The former included stroke addition, subtraction or substitution. The latter referred to character substitution by those with similar phoneme, grapheme or semanteme. (**Figures 2, 3**)

In comparison, AD patients showed a more severe form of global aphasia. Both oral and reading comprehension were affected at the word or phrase level (**Figure 2**). They showed more naming errors, as well as more verbal category fluency difficulties. They exhibited fluent aphasia, which was characterized by an effortless and empty speech. The speech prosody was almost normal (**Figure 2**). As shown in **Figures 2 and 3**, in spontaneous writing, two wrote nothing and one only wrote her name. The other two stacked some nouns or pronouns, with lack of verbs or functional words. Besides, they showed more character substitu-

Table 3. Neuropsychological data and drug treatment of HD and AD patients

Case	Screening test (score)				NPI (score)			Cognitive domain* (grade)				Drug treatment	
	MMSE	Moca	ADLs	HADS-A/D	Frequency	Severity	Distress	Executive	Visuospatial	Memory	Reasoning	Dose	Duration
1	22	17	20	2/0	2	2	2	2	1	2	2	-	-
2	24	17	24	0/0	0	0	0	1	0	2	2	-	-
3	25	21	24	3/4	3	2	3	1	1	2	1	-	-
4	24	18	27	0/5	14	13	17	2	1	3	1	-	-
5	25	26	24	0/0	6	5	6	1	1	1	1	DHT 5 mg/d	6 years
6	23	17	22	6/6	-	-	-	1	0	3	3	DHT 10 mg/d+ 1 month MEM 20 mg/d	
7	5	3	23	6/4	-	-	-	3	2	3	3	DHT 10 mg/d	2 months
8	5	3	60	0/0	-	-	-	3	3	3	3	DHT 5 mg/d	1 month
9	9	9	31	7/10	-	-	-	2	2	3	3	MEM 20 mg/d	3 months
10	17	9	24	2/1	-	-	-	2	1	3	3	MEM 15 mg/d	1 month

Case 1-5: HD patients; Case 6-10: AD patients; HADS-D/A: depression/anxiety section of Hospital Anxiety and Depression Scale; DHT: donepezil hydrochloride tablet; MEM: memantine hydrochloride tablets.

*Rating criteria for cognitive domain: Grade 0: normal; most tests had a score above -1 SD of the mean; Grade 1: slight abnormality; most tests had a score within -1 to -1.5 SD of the mean; Grade 2: moderate abnormality; most tests had a score within -1.5 to -2.5 SD of the mean; Grade 3: severe abnormality; most tests had a score below -2.5 SD of the mean.

Figure 2. Linguistic characteristics of HD and AD patients. This is a simple scatterplot created in SPSS Statistics. It displays the linguistic scores of HD (group 1) and AD (group 2) patients in 11 aspects (Item 1-11). 1. oral comprehension; 2. repetition; 3. oral expression; 4. verbal category fluency; 5. naming; 6. reading; 7. reading comprehension; 8. copying; 9. dictation; 10. writing on picture; 11. spontaneous writing. HD patients show a poor language output with relatively spared input and conduction (both oral and written language). AD patients show a more severe form of global aphasia with written language affected the most.

tions by those with relevant semanteme. For example, one patient wrote SLEEP when she saw a BED. Another answered SUN when asked "WHAT COLOUR IS THE SUNNY SKY".

DISCUSSION

With regard to cognitive profile, HD patients present with executive, reasoning and memory impairment. They can finish DSST, TMT-A and block design tests with a high accuracy but a low speed. This indicates the contribution of decreased psychomotor speed in executive dysfunction. However, even if enough time is given, they all show difficulty in gesture imitation. Therefore, other executive resource should exist in addition to the decreased psychomotor speed. After removing the possible influence from decreased psychomotor speed, the visuospatial function is relatively spared. In AVLT, they do not benefit from semantic cueing. The recognition of AVLT and Rey complex figure is also poor. These indicate that besides memory retrieval deficit due to executive dysfunction, there should be an encoding or storage problem.

As for linguistic profile, HD patients suffer from phonological impairment, as well as slight semantic and syntactic abnormality. These are in accordance with the previous findings.^[15, 16] The speech dyspros-

Figure 3. Writing examples of HD and AD patients.

Pictures A-C are from a 67-year-old HD patient, who has a 2-year disease course and 6-year education. Pictures D-E are from a 47-year-old AD patient, who has a 4-year disease course and 14-year education. Their cognitive screening scores are close, whereas the writings are quite different.

The correct answers for dictation (A, D) are COMB (梳子), KEY (钥匙), SLEEP (睡觉), RUN AWAY (跑走), FIVE-POINTED STAR (五星), THE SPRING WIND BLOWS AND THE TREES TURN GREEN (春风吹绿了树叶). The correct answers for writing on picture (B, E) are CHAIR (椅子), BED (床), FLOWER (花), TRIANGLE (三角形), CIRCLE (圆形), RED (红色), YELLOW (黄色), SMOKING (抽烟), DRINKING WATER (喝水), FALLING DOWN (摔倒).

In spontaneous writing (C), the HD patient meant to write "I HAVE HYPERACTIVITY DISORDER. AFTER SEEING A DOCTOR IN THE HOSPITAL, MY CONDITION IMPROVED GREATLY. I AM VERY GRATEFUL TO THE DOCTOR IN THE HOSPITAL". (我有多动病, 在医院看后, 病情大大好转, 我很感谢医院医生). The basic structure of the sentences remains. Strictly speaking, the last two sentences are lack of subjects. Additionally, he shows plenty of character-formation errors and character substitutions. Some characters exhibit stroke omission (角, 院) or substitution (匙, 睡, 在, 感). Some are replaced by others with similar phoneme (形→行, 图形→钉, 匙→钅, 大→太, 谢→请) or semanteme (绿→蓝, 病→医).

The AD patient can only write her name in spontaneous writing. She shows some character substitutions by those with relevant semanteme [床(BED)→睡觉(SLEEP), 花(FLOWER)→书(BOOK), 抽烟(SMOKING)→南京(NANJING) (a cigarette brand in China)].

dy might be related to voluntary motor disorder in HD. They have a slight deficit in comprehension, repetition, naming and verbal category fluency. These might be related to slight semantic or syntactic impairment, or retrieval deficit due to executive dysfunction.

Compared with native speakers of alphabetic languages, Mandarin-speaking HD patients have their own linguistic features. In writing, they show character structural errors and character substitutions, instead of letter omissions.^[3] In addition, Chinese is a typical tonal language with four tones. The same phoneme with different tones indicates different meanings. HD patients show tonic errors in speech. The third tone is pronounced as the first tone.

In comparison, AD patients present with a worse cognitive and linguistic impairment. The executive, visuospatial, memory and reasoning domains are all affected. They do not have phonological impairment as

HD patients. However, they seem to have more loss of semantic knowledge. They show a worse performance in comprehension, naming and verbal category fluency. And they have more character substitutions of semantic category in both oral and written language.

These cognitive and linguistic differences between HD and AD patients might be related to their different pathological changes. The neurodegeneration in HD starts from caudate and putamen. Gradually, it extends to the frontostriatal connections between striatum and cingulate, precentral, prefrontal cortices.^[17] The frontostriatal network is implicated in motor-language coupling, syntactic processing, executive function, working memory, emotional processing, etc.^[18, 19] Unlike HD, AD is generally characterized by progressive degeneration of medial temporal, frontal and parietal cortices.^[9] These cerebral cortices are involved in a wide range of cognitive and linguistic activities. For example, the

frontal and temporal cortices are implicated in semantic processing.^[20]

Brain MRI demonstrates cortical atrophy in both HD and AD patients. However, the severity of cortical atrophy is not parallel to their cognitive and linguistic behaviors. Compared with AD patients, HD patients show more severe cortical atrophy, but less cognitive and language impairment. These might be associated with their different genetic bases. HTT CAG-repeat expansion may influence cortical neurodevelopment. However, the influence might be independent of neurodegeneration.^[21]

In conclusion, Mandarin-speaking HD patients have a TCMA-like disturbance with prominent phonological impairment, whereas AD patients have a global aphasia with salient semantic impairment. The limitation of this paper is that the sample size is too small. We cannot stratify the subjects according to the disease course or drug treatment. Next, we expect to accumulate more cases and have a longitudinal follow-up.

Conflict of Interests Statement

The authors have no conflict of interests to report.

REFERENCES

1. The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 1993; 72(6):971-83. doi: 10.1016/0092-8674(93)90585-e.
2. Roos RAC. Huntington's disease: a clinical review. *Orphanet J Rare Dis* 2010; 5(1):40. doi: 10.1186/1750-1172-5-40.
3. Gagnon M, Barrette J, Macoir J. Language disorders in Huntington disease: a systematic literature review. *Cogn Behav Neurol* 2018; 31(4):179-92. doi: 10.1097/WNN.000000000000171.
4. Berninger V, Abbott R, Cook CR, et al. Relationships of attention and executive functions to oral language, reading, and writing skills and systems in middle childhood and early adolescence. *J Learn Disabil* 2017; 50(4):434-49. doi: 10.1177/0022219415617167.
5. Minkina I, Rosenberg S, Kalinyak-Fliszar M, et al. Short-term memory and aphasia: from theory to treatment. *Semin Speech Lang* 2017; 38(1):17-28. doi: 10.1055/s-0036-1597261.
6. Diehl JJ, Frost SJ, Sherman G, et al. Neural correlates of language and non-language visuospatial processing in adolescents with reading disability. *Neuroimage* 2014; 101:653-66. doi: 10.1016/j.neuroimage.2014.07.029.
7. Krzemien M, Jemel B, Maillart C. Analogical reasoning in children with specific language impairment: evidence from a scene analogy task. *Clin Linguist Phon* 2017; 31(7-9):573-88. doi: 10.1080/02699206.2017.1302509.
8. Vonsattel JP, Myers RH, Stevens TJ, et al. Neuropathological classification of Huntington's disease. *J Neuropathol Exp Neurol* 1985; 44(6):559-77. doi: 10.1097/00005072-198511000-00003.
9. Braak H, Braak E. Neuropathological staging of Alzheimer related changes. *Acta Neuropathol (Berl)* 1991; 82(4):239-59. doi: 10.1007/bf00308809.
10. Folstein SE, Leigh RJ, Parhad IM, et al. The diagnosis of Huntington's disease. *Neurology* 1986; 36(10):1279-83. doi: 10.1212/wnl.36.10.1279.
11. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011; 7(3):263-9. doi: 10.1016/j.jalz.2011.03.005.
12. Gao SR. Aphasia. 2nd ed. Beijing: Peking University Medical Publishing House; 2006.p.568-93. Chinese.
13. Tan JP, Li N, Gao J, et al. Optimal cutoff scores for dementia and mild cognitive impairment of the Montreal Cognitive Assessment among elderly and oldest-old Chinese population. *J Alzheimers Dis* 2015; 43(4):1403-12. doi: 10.3233/JAD-141278.
14. Pasquier F, Leys D, Weerts JG, et al. Inter- and intraobserver reproducibility of cerebral atrophy assessment on MRI scans with hemispheric infarcts. *Eur Neurol* 1996; 36(5):268-72. doi: 10.1159/000117270.
15. Azambuja MJ, Radanovic M, Haddad MS, et al. Language impairment in Huntington's disease. *Arq Neuropsiquiatr* 2012; 70(6):410-5. doi: 10.1590/s0004-282x2012000600006.
16. Rusz J, Saft C, Schlegel U, et al. Phonatory dysfunction as a preclinical symptom of Huntington disease. *PLoS One* 2014; 9(11):e113412. doi: 10.1371/journal.pone.0113412.
17. Beste C, Ness V, Falkenstein M, et al. On the role of fronto-striatal neural synchronization processes for response inhibition—evidence from ERP phase-synchronization analyses in pre-manifest Huntington's disease gene mutation carriers. *Neuropsychologia* 2011; 49(12):3484-93. doi: 10.1016/j.neuropsychologia.2011.08.024.

18. Birba A, García-Cordero I, Kozono G, et al. Losing ground: frontostriatal atrophy disrupts language embodiment in Parkinson's and Huntington's disease. *Neurosci Biobehav Rev* 2017; 80:673-87. doi: 10.1016/j.neubiorev.2017.07.011.
19. De Diego-Balaguer R, Couette M, Dolbeau G, et al. Striatal degeneration impairs language learning: evidence from Huntington's disease. *Brain* 2008;131(Pt 11):2870-81. doi: 10.1093/brain/awn242.
20. Wawrzyniak M, Hoffstaedter F, Klingbeil J, et al. Front-to-temporal interactions are functionally relevant for semantic control in language processing. *PLoS One* 2017; 12(5):e0177753. doi: 10.1371/journal.pone.0177753.
21. Kubera KM, Schmitgen MM, Hirjak D, et al. Cortical neurodevelopment in pre-manifest Huntington's disease. *Neuroimage Clin* 2019; 23:101913. doi: 10.1016/j.nicl.2019.101913.