ORIGINAL ARTICLE

Characteristics of Chinese Families in Which Children and Both Parents Are Diagnosed with Malignant Tumors: A Retrospective Study

Ju Liu, Zhijian Xu*, Xiaofeng Bi, Ping Sun, Jiaqin Huang

Department of Cancer Prevention, National Cancer Center / National Clinical Research Center for Cancer / Cancer Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100021, China

Key words: family history; parents; daughter; son; early detection; malignant tumor

Objective To characterize Chinese families in which both parents and at least one child are diagnosed with malignant diseases and provide reference for cancer screening or early detection in people whose both parents are diagnosed with cancer.

Methods Medical records of all clients to the center of cancer screening and prevention of the National Cancer Center/Cancer Hospital between January 2008 and February 2018 were screened to select families in which both parents and at least one child were diagnosed with malignant diseases. The cancer profiles of fathers, mothers, sons and daughters, their age distribution at diagnosis, and similarity of cancers between two generations were analyzed. The proportions of each cancer in males and females of the cohort were compared with corresponding data from the National Cancer Center Registry of China (NCCRC) in 2013.

Results Totally 135 families were identified from records of 33 200 clients. Proportion of lung cancer in fathers (40/135, 29.6%) and in mothers (38/135, 28.1%) were higher than the national data (23.9% in males and 14.9% in females, respectively). The proportion of breast cancer in daughters (35/109, 32.1%) was higher than that of mothers (14/135, 10.4%) and the national data (17.1%). In 71 father-son pairs of cancer, 46.5% (33/71) were of the same systematic disease, and 16.9% (12/71) were of the same cancer. These two indexes were 31.2% ($n=34$) and 10.1% ($n=11$), respectively in the 109 father-daughter pairs of cancer, 36.6% ($n=26$) and 8.5% ($n=6$) respectively in the 71 mother-son pairs of cancer, and 31.2% ($n=34$) and 20.2% ($n=20$) respectively in the 109 mother-daughter pairs of cancer. Sons were more likely to suffer from cancers originated from the same system as father's cancer than daughters ($\chi^2=4.299$, $P<0.05$), and daughters were more likely to suffer from the same cancer as their mother's cancer than sons ($\chi^2=4.506$, $P<0.05$). The age (mean $\pm$ standard deviation)

Received June 3, 2020; accepted September 15, 2020.

*Corresponding author Zhijian Xu, E-mail: xuzj678@aliyun.com

This work was supported by the National Key R&D Program of China (2017YFC1308700) and Peking Union Medical College Discipline Development Project (201920200303).

of the daughters (52.4 ± 12.7) and the sons (59.4 ± 10.9) at diagnosis were significantly younger than the fathers (65.5 ± 12.2) and the mothers (65.7 ± 12.5) (all $P < 0.001$).

Conclusions For people whose both parents are diagnosed as cancer, screening or early detection examinations should cover a full range of cancers rather than the cancers their father and mother have suffered, or cancers stemmed from the same system as their parent's cancers. We suggest screening or early detection program for these special population start earlier than that for the general population, with emphasis on cancers derived from digestive system for males and women-specific cancers, i.e., breast cancer, ovarian cancer, cervical cancer and uterine cancer for females.

ACCORDING to China Statistical Year 2019 published by National Bureau of Statistics of China, cancer has become the leading cause of death in Chinese of urban areas. This leads to an increasing number of people who has a family history of cancer in the general population. The fact that 90% men whose father was diagnosed with prostate cancer were positively inclined to undergo screening for prostate cancer^[1] revealed their concerns about the possible inheriting and the significant psychological effects caused by the first-degree relatives being diagnosed with malignant diseases. For those whose parents are both diagnosed with cancer, psychological stress, concerns about cancer screening and early detection are doubled.

Children share similar environmental risk factors of cancer with parents before grown up. Moreover, the lifestyles of parents have a significant influence on their children. Therefore, parents and children have the highest hereditary and environmental similarities as far as cancer risk factors are concerned. Previous studies have revealed that the familial history of parents suffering from cancer, such as cancers of breast, colon, and prostate, constitutes a risk factor for children suffering the same cancer.^[2-4] A research based on the Swedish Cancer Registry data from 1958 to 1996 showed that there is an increased risk of cancers in 21 out of 23 cancer sites for offspring with familiar cancer history of either mother or father.^[5] If both parents had cancer, the risk of their children increased by 1.39 times for sons and 1.34 times for daughters.^[6]

Unlike the Swedish population-based study, we focused on families in which children and both parents were diagnosed with cancer. Analyses of the characteristics of this special group provide information about whether the disease profile of children is related to their parents and their gender. This information would be beneficial to formulate screening strategies and facilitate the early detection for Chinese people with

family history of cancer in both parents.

PATIENTS AND METHODS

Study population

We retrospectively examined the database of all clients of the Cancer Screening and Prevention center, the National Cancer Center/Cancer Hospital between January 2008 and February 2018 for cancer screening or early detection. Doctors performed epidemiological survey on the family history of cancer and filled in information form for each client with consent prior to physical examinations. Data were reviewed in order to select families in which father, mother and at least 1 child were diagnosed with primary malignancies pathologically. Other inclusion criteria: parents were the biological mother or father of the diagnosed children, and they all lived together at least till children's age of 18 years old. The excluding criteria include: 1) primary cancer site of any family member was unknown; 2) age of any family member with cancer at diagnosis was unavailable; 3) families with diagnosed hereditary syndrome. The study was approved by the institutional review board, written informed consents were waived.

Data analyses

Proportions of malignant diseases in the subgroup of fathers, mothers, sons, and daughters in this study were compared with the published national data of NCCRC^[7] in terms of the common cancers, the digestive tract cancers (cancer of esophageal, stomach, colon, and rectum), digestive system cancers (cancer of stomach, esophageal, liver, pancreas, and colon), women specific cancers (breast cancer, ovarian, uterus and cervical cancer).

Continuous data were expressed as mean $\pm$ standard deviation (SD). Categorical variables were presented as number and percentage. Chi-square test was used to determine the similarity of malignant diseases

between parents and children. The age differences at diagnosis between parents and children, daughters and sons in the research population were analyzed using independent *t*-test. Statistical significance was defined as $P < 0.05$ bilaterally. Data analysis was conducted using SPSS (Version 24.0, IBM, USA)

RESULTS

General information of study population

There were totally 33 200 individuals came to the Center of Cancer Screening and Prevention between January 2008 and February 2018, and 135 families were identified and included in this study. Among them, 105 families (77.8%) had 1 child diagnosed with malignancy, and 30 families (22.2%) had 2 or more children diagnosed with malignancies. Of 180 children with malignancies in the research population, 71 (39.4%) were sons and 109 (60.6%) were daughters. Of the 105 families in which only 1 child was diagnosed with malignancies, there were 37 sons (35.2%) and 68 daughters (64.8%). While among the 30 families with 2 or more children affected with malignancies, malignancies were suffered by 2 sons in 6 families, 2 daughters in 6 families, 1 son and 1 daughter in 9 families, 2 daughters and 1 son in 2 families, 3 daughters in 2 families, 4 sons in 1 family, 2 sons and 2 daughters in 2 families, 1 son and 3 daughters in 1 family, and 2 sons and 3 daughters in 1 family.

Proportion of malignant diseases

Of 135 families with parents and child/children suffering from malignancies, lung cancer, ranked as the top one common cancer, was diagnosed in 29.6% fathers ($n=40$), which was higher than that of the NCCRC data in males (23.9%); while lung cancer was found in 19.7% ($n=14$) sons, which was lower than that of the NCCRC data. The digestive system cancers stand as the top 5 most common cancers in both fathers and sons (**Table 1**).

There were 28.1% ($n=38$) mothers diagnosed as lung cancer, and this proportion was much higher than that determined by the NCCRC data (14.9%); on the contrast, only 10.4% ($n=14$) mothers were diagnosed with breast cancer, which was much lower than that determined by the NCCRC data (17.1%). While the proportion of breast cancer in 109 daughters were 32.1%, which was much higher than that in mothers (10.4%) and the national data (17.1%). The colorec-

tal cancers were found in 14.7% daughters ($n=16$), which was comparable to that of mothers, but higher than that of the NCCRC data (9.1%). Thyroid cancers were found in 9.2% daughters ($n=10$), which was also higher than that in mothers (2.2%) and that of NCCRC data (6.6%). The proportions of other cancers in women in the study population were similar to that of the NCCRC data on women.

In total, there were 46 daughters suffering from female-specific cancers (breast, uterine, cervical, and ovarian cancers), constituting 42.2% of all the malignancies in daughters; while when combining female-specific cancers with lung cancers together, it constituted 54.1% of all malignancies in daughters (**Table 2**).

The proportions of the mothers with digestive tract cancers (33.3%) and digestive system cancers (38.5%) were higher than that of the NCCRC data (21.9% and 27.7%, respectively). A similar tendency was also found among the sons (39.4% vs. 33.9%, and 60.5% vs. 49.4%, respectively). The combined proportion of digestive system and lung cancer accounts for 66.7% of all cancers in subgroup of mothers, which is higher than the NCCRC data on women (42.6%).

Similarity of cancers in parents and among children

Among the 71 pairs of father-son who both suffered malignancies, 33 (46.5%) pairs suffered malignant diseases of the same system, including digestive system disease (72.7%, $n=24$), respiratory system disease (24.2%, $n=8$), and reproductive system disease (3.0%, $n=1$). Besides, 12 father-son pairs (16.9%) shared exactly the same malignant disease, including lung cancer ($n=6$), gastric cancer ($n=4$), hepatic cancer ($n=1$), and esophageal cancer ($n=1$). Among these 12 cases, digestive tract diseases accounted for 41.7% ($n=5$), while digestive system diseases accounted for 50.0% ($n=6$).

Among the 109 father-daughter pairs, 34 (31.2%) pairs suffered the malignant diseases of the same system, including digestive system (61.8%, $n=21$), reproductive system (17.6%, $n=6$), respiratory system (14.7%, $n=5$), and endocrine system (5.9%, $n=2$). In 9 families with fathers diagnosed as prostate cancer, 6 daughters (66.7%) were breast cancer patients. Besides, there were 11 (10.1%) father-daughter pairs who suffered exactly the same malignant diseases,

Table 1. Malignancies and proportion ranking in subgroups of fathers and sons of Chinese families with child/children and both parents diagnosed with malignancies

Rank	Fathers (n=135)		Sons (n=71)		NCCRC data of men ^[7]	
	Disease	n (%)	Disease	n (%)	Disease	%
1	Lung cancer	40 (29.6)	Lung cancer	14 (19.7)	Lung cancer	23.9
2	Gastric cancer	28 (20.7)	Gastric cancer	13 (18.3)	Gastric cancer	14.6
3	Esophageal cancer	15 (11.1)	Hepatic cancer	10 (14.1)	Hepatic cancer	13.1
4	Hepatic cancer	13 (9.6)	Colorectal cancer	10 (14.1)	Colorectal cancer	9.8
5	Colorectal cancer	9 (6.7)	Esophageal cancer	5 (7.0)	Esophageal cancer	9.6
6	Prostate cancer	5 (3.7)	Pancreatic cancer	5 (7.0)	Prostate cancer	2.9
7	Bladder cancer	5 (3.7)	Prostate cancer	2 (2.8)	Bladder cancer	2.9
8	Pancreatic cancer	5 (3.7)	Lymphoma	2 (2.8)	Pancreatic cancer	2.5
9	Nasopharyngeal cancer	3 (2.2)	Laryngeal cancer	2 (2.8)	Lymphoma	2.3
10	Laryngeal cancer	2 (1.5)	Brain tumor	2 (2.8)	Brain and central nervous system tumor	2.3
11	Cholangiocarcinoma	2 (1.5)	Nasopharyngeal cancer	1 (1.4)		
12	Lymphoma	2 (1.5)	Thyroid cancer	1 (1.4)		
13	Thyroid cancer	1 (0.7)	Bladder cancer	1 (1.4)		
14	Brain tumor	1 (0.7)	Melanoma	1 (1.4)		
15	Kidney cancer	1 (0.7)	Leiomyosarcoma	1 (1.4)		
16	Gastrointestinal stromal tumor	1 (0.7)	Penis cancer	1 (1.4)		
17	Oral cancer	1 (0.7)				
18	Penis cancer	1 (0.7)				

NCCRC, National Cancer Center Registry of China.

Table 2. Malignancies and the proportion ranking in subgroups of mothers and daughters in Chinese families with child/children and both parents diagnosed with malignancies

Rank	Mothers (n=135)		Daughters (n=109)		NCCRC data of women ^[7]	
	Disease	n(%)	Disease	n(%)	Disease	%
1	Lung cancer	38(28.1)	Breast cancer	35(32.1)	Breast cancer	17.1
2	Gastric cancer	17(12.6)	Colorectal cancer	16(14.7)	Lung cancer	14.9
3	Colorectal cancer	16(11.9)	Lung cancer	13(11.9)	Colorectal cancer	9.1
4	Breast cancer	14(10.4)	Thyroid cancer	10(9.2)	Gastric cancer	7.9
5	Esophageal cancer	12(8.9)	Gastric cancer	6(5.5)	Thyroid cancer	6.6
6	Hepatic cancer	5(3.7)	Hepatic cancer	5(4.6)	Cervical cancer	6.2
7	Cervical cancer	5(3.7)	Ovarian cancer	5(4.6)	Hepatic cancer	5.8
8	Kidney cancer	4(3.0)	Esophageal cancer	3(2.7)	Esophageal cancer	4.9
9	Uterine cancer	4(3.0)	Uterine cancer	3(2.7)	Uterine cancer	3.8
10	Thyroid cancer	3(2.2)	Cervical cancer	3(2.7)	Ovarian cancer	3.1
11	Ovarian cancer	3(2.2)	Pancreatic cancer	3(2.7)		
12	Leukemia	3(2.2)	Lymphoma	2(1.8)		
13	Pancreatic cancer	2(1.5)	Thymoma	1(0.9)		
14	Cholangiocarcinoma	2(1.5)	Brain tumor	1(0.9)		
15	Bladder cancer	2(1.5)	Nasopharyngeal cancer	1(0.9)		
16	Lymphoma	1(0.7)	Melanoma	1(0.9)		
17	Ureter cancer	1(0.7)	Mesothelioma	1(0.9)		
18	Pharyngeal cancer	1(0.7)				
19	Osteosarcoma	1(0.7)				
20	Glioma of brain	1(0.7)				

including lung cancer ($n=4$), thyroid cancer ($n=2$), esophageal cancer ($n=3$), gastric cancer ($n=1$), and pancreatic cancer ($n=1$). Malignancies of digestive tract and digestive system accounted for 36.4% ($n=4$) and 45.5% ($n=5$), respectively of the cancers suffered by both father and daughter.

Among the 71 mother-son pairs, 26 (36.6%) pairs suffered malignant diseases of the same system, including 22 (84.6%) digestive system diseases, 3 (11.5%) respiratory system diseases, and 1 (3.8%) reproductive system disease. There were 6 (8.5%) mother-son pairs suffered exactly the same malignant diseases: lung cancer ($n=2$), esophageal cancer ($n=2$), colorectal cancer ($n=1$), and hepatic cancer ($n=1$). In addition, malignancies of digestive tract and digestive system accounted for 50.0% ($n=3$) and 66.7% ($n=4$), respectively of cancers suffered by both mother and son.

Among 109 mother-daughter pairs, 34 (31.2%) had malignant diseases of the same system, including 17 (50.0%) digestive system diseases, 10 (29.4%) reproductive system diseases, and 7 (20.6%) respiratory system diseases. There were 22 (20.2%) mother-daughter pairs suffered exactly the same malignant diseases: lung cancer ($n=7$), breast cancer ($n=6$), colorectal cancer ($n=4$), liver cancer ($n=1$), uterine cancer ($n=1$), cervical cancer ($n=1$), esophageal cancer ($n=1$), and gastric cancer ($n=1$). Malignancies of digestive tract and reproductive system accounted for 27.3% ($n=6$) and 36.4% ($n=8$), respectively of cancers suffered by both mother and daughter.

Father-son pairs were more likely to suffer can-

cers of the same system than were father-daughter pairs ($\chi^2=4.299$, $P=0.038$). Mother-daughter pairs were more likely to suffer from the same disease than were the mother-son pairs ($\chi^2=4.506$, $P=0.034$). There were no significant difference in suffering from the same disease between the father-son and father-daughter pairs, and no significant difference in suffering from diseases of same system between mother-son and mother-daughter pairs (**Table 3**).

Age at diagnosis

The age (years, mean $\pm$ SD) at diagnosis of the fathers was significantly older than the sons (65.5 $\pm$ 12.2 vs. 59.4 $\pm$ 10.9, $t=3.566$, $P<0.001$), and the daughters (65.5 $\pm$ 12.2 vs. 52.4 $\pm$ 12.7, $t=8.176$, $P<0.001$). Similarly, the age at diagnosis of the mothers was significantly older than the sons (65.7 $\pm$ 12.5 vs. 59.4 $\pm$ 10.9, $t=3.579$, $P<0.001$) and the daughters (65.7 $\pm$ 12.5 vs. 52.4 $\pm$ 12.7, $t=8.143$, $P<0.001$). Additionally, the age of sons at diagnosis was significantly older than that of daughters (59.4 $\pm$ 10.9 vs. 52.4 $\pm$ 12.7, $t=3.778$, $P<0.001$). The age distribution at diagnosis was presented in **Table 4**.

Among father-children pairs, 17 fathers were diagnosed with malignancies before 50 years old, with the mean age of their sons ($n=7$) and daughters ($n=10$) at diagnosis being 53.6 years and 52.7 years, respectively. Among mother-children pairs, 23 mothers were diagnosed before 50 years old, with the mean age of their sons ($n=9$) and daughters ($n=14$) being 56.0 years and 50.6 years at diagnosis, respectively.

Table 3. Daughters and sons diagnosed with the same cancer or cancers from the same system with fathers or mothers in 135 families that both parents suffering malignancies.

Parents with cancer	Cancers of same system, n (%)		Same cancer, n (%)	
	Son($n=71$)	Daughter($n=109$)	Son ($n=71$)	Daughter ($n=109$)
Father ($n=135$)	33 (46.5) ^a	34 (31.2)	12 (16.9)	11 (10.1)
Mother ($n=135$)	26 (36.6)	34 (31.2)	6 (8.5)	22 (20.2) ^b

^a $P<0.05$ compared with the father-daughter pairs; ^b $P<0.05$ compared with the mother-son pairs.

Table 4. The distribution of age at diagnosis of fathers, mothers and children who were diagnosed with malignancies in 135 Chinese families [n (%)]

Sub-group	<30 yrs	30-39 yrs	40-49 yrs	50-59 yrs	60-69 yrs	70-79 yrs	≥ 80 yrs
Fathers ($n=135$)	0	3(2.2)	12(8.9)	24(17.8)	43(31.8)	38(28.1)	15(11.1)
Mothers ($n=135$)	0	4(3.0)	9(6.7)	22(16.3)	38(28.1)	44(32.6)	18(13.3)
Sons ($n=71$)	0	1(1.4)	10(14.1)	24(33.8)	21(29.6)	12(16.9)	3(4.2)
Daughters ($n=109$)	2(1.8)	9(8.3)	40(36.7)	30(27.5)	15(13.8)	11(10.1)	2(1.8)

DISCUSSION

Our study found in cancer patients with both parents also diagnosed with malignant diseases, lung cancer in their parents had a higher proportion than that of national data on lung cancers. Breast cancer in the daughters had a much higher proportion than in the mothers and that of national data. The probability of children developing the same malignant disease or a disease from the same system with their parents was relatively low. Ages of the fathers and mothers at diagnosis were comparable, but their son or daughter with malignancy were generally diagnosed at a younger age than their parents.

Our study comprised 109 daughters and 71 sons who suffered from malignancies. We found daughters tended to develop malignant diseases 7 years earlier than did the sons. These findings corresponded to a previous study which showed that in people whose mother was diagnosed with malignant diseases, more daughters than sons suffered from cancer.^[8] Therefore, when both parents have been diagnosed with cancers, their daughters may be at a higher risk than their sons. However, in a Swedish study, it was suggested that sons had a 1.39 times higher risk while daughters had a 1.34 times higher risk.^[6] The fact that more daughters with cancer than sons with cancer in this study can be partly explained by that the age-standardized incidence rate of women increases by 2.2% per year.^[7] Daughters in our study seem at higher risk than sons. This phenomenon needs further investigation.

Because the age of the subjects in our study ranged from 25-80 years old, the time at diagnosis for a family member ranged from 1930s to 1990s, which may overlap between different family members. For example, children of one family may be diagnosed at a time earlier than parents of another family. Therefore, we chose the nearest national data for comparison.

The proportion of breast cancer in mothers (10.4%) ranked fourth, and was lower than that in daughters (32.1%) and that determined by the national data (17.1%) where breast cancer ranked as the most common cancer. The higher proportion of breast cancer among daughters than mothers may be associated with the increasing incidence of breast cancer in Chinese women in the past a few decades.^[7] Conversely, the increased proportion of breast cancer corresponded to the findings of a previous study that

showed the incidence of breast cancer in daughters with either mother or father diagnosed with malignancy was higher than that of the general women population; the proportion of breast cancer was 32.7% in daughters whose maternal parent was diagnosed with cancers, and 27.7% whose agnate parent was diagnosed with cancers.^[8, 9]

The average age of fathers and mothers at diagnosis were almost the same; however, we found their child/children were diagnosed with cancer at significantly younger age, while daughters suffered cancers at an even younger age than sons. The age difference at which parents and children being diagnosed showed that environmental factors may play important roles in the carcinogenesis within a family. Fathers and mothers shared similar environmental risk factors with each other for a longer time than with their children; as children grew up, they may be exposed to different environment with different risk factors, their ages at diagnosis of malignancies differed from one another and from their parents. Besides, the advances in examination techniques and the increasing awareness of early detection of cancers cause more cancers detected in population, which might also contribute to the age differences of diagnoses between the two generations.

Compared to the NCCRC data, the proportion of lung cancer in fathers (29.6%) was relatively high, while that in sons was relatively low. In contrast, the proportion of lung cancer in mothers was twice as high as that of the national data (28.1% vs. 14.9%),^[7] and ranked the first, while lung cancer (11.9%) ranked the third in daughters. Both sons and daughters had a lower proportion of lung cancer, while both fathers and mothers had a higher proportion of lung cancer, when compared with the national data. Environmental factors associated with lung cancer, as reported, include smoking, passive smoking, air pollution, exposure to carcinogenic chemicals, and alcohol consumption.^[10] The smoking rate of Chinese women was only 3.4%, which is much lower than that of men (62.4%).^[11] Generally, parents live with each other for much longer time than with their children. We infer that higher proportion of lung cancer in mothers might be related to passive smoking. The prevalence of smoking was reported to decrease slightly during the period from 1991 to 2011,^[12] which might be associated with lower proportions of lung cancer in sons and daughters than in their parents.

In the studied families, 72.7% (24/33) father-son

pairs, 84.6% (22/26) mother-son pairs, 61.8% (21/34) father-daughter pairs, and 50.0% (17/34) mother-daughter pairs suffered cancers from digestive system. For gastric cancer, *Helicobacter pylori* infection and insufficient nutritional intake have been confirmed as major risk factors.^[13, 14] More fruit and vegetable intake has been shown to be a protective factor for esophageal and pancreatic cancers.^[15, 16] The risk factors of pancreatic cancer also include smoking and alcohol,^[17] while virus infection was the main causative factor of hepatic cancer among Chinese.^[18] Colon cancer is related to diet and nutritional factors.^[19] Most of the risk factors for digestive system cancers are related to lifestyle factors, such as diet, infectious factors, and alcohol consumption.

The proportion of digestive system cancers in mothers were much higher than the national data; among fathers and daughters, it was also slightly higher than the national data. As digestive system cancers are mainly caused by nonhereditary, especially lifestyle factors, diet habits of parents influence their children greatly. Lifestyle, especially of a mother, may put a great impact on the development of digestive system cancers within a family.

On the other hand, people whose both parents are diagnosed with malignancies concern much more about their risk for cancer, and the strategy of cancer screening and/or early detection. Since smoking, diet, and alcohol consumption are the main modifiable risk factors for the incidence of malignant diseases within a family, we suggest that individuals whose both parents are diagnosed with malignant diseases strive to lead a very healthy life, quit smoking and drinking, eat more vegetables and fruits, prevent from infections of virus and *Helicobacter pylori*, and pursue treatment at earlier stage of diseases.

Furthermore, regular screenings and early detection for cancers are of paramount importance for this special population. For the starting age of screening, as the result of this study, the mean age at diagnosis of sons was 6.3 years younger than the parents, and mean age of daughters at diagnosis was 13.3 years younger than their patients, 63.4% of sons were diagnosed between age of 50 and 69 years old, and more than one-third of daughters were diagnosed with cancer between age of 40 and 49 years old, we suggest that individuals with both parents suffering cancer should begin cancer screening earlier than general Chinese population.

For the targeting cancers, this special population need to take screening or early detection examinations for all common cancers, not just for the cancers of parents, or cancers originated from the same system as of their parents. Among sons, 28 (39.4%) were diagnosed with digestive tract cancer. Digestive tract cancers and lung cancer in sons preserved high proportions among father-children pairs and mother-children pairs who suffering from exactly the same cancer. The proportion of the digestive system cancer in sons accounted for 60.5%, and increase to 80.3% if combined with lung cancer. Therefore, we suggest men with family history of malignant diseases in both parents should participate screening and early detection for digestive system cancers and lung cancer.

Breast cancer occurred in almost one-third of the daughters; besides, female-specific cancers accounted for 42.2% and thyroid cancer accounted for 9.2% of all cancers among daughters. National data showed that the age-standardized incidence of lung, breast, cervix, uterine ovarian, and thyroid cancers significantly increased, whereas that of esophageal, stomach, and liver cancers showed a decreasing trend.^[7] Therefore, screening program of women with both parents diagnosed with malignancy should focus on female-specific cancers, especially breast cancer. If combining the above mentioned cancers with lung cancer and thyroid cancers, it covers 63.3% of all the malignancies; if digestive system diseases were also considered together, the combined proportion is 93.6% of all malignancies. These data provide evidence in designing cancer screening program for females in this special population. Additionally, of 9 daughters whose fathers had prostate cancer, 6 suffered from breast cancer, so females whose fathers have prostate cancer should also pay special attention to breast cancer screening. Hormone-related factors might be involved in the underlying mechanism, which need to be further investigated.

As virus infections were the risk factors for hepatic and cervical cancer, we suggest that individuals with both parents diagnosed with malignancies need to take screenings for virus infections and early detection examinations as per the disease-specific guidelines.

To conclude, our study provides some clues for cancer etiology researches, cancer screening programmes and early detection strategy in people whose both parents are diagnosed with cancers. It should be noted that as a retrospective study, the diagnostic

time of the subjects in this study scattered in several decades, in which the incidences and mortalities of cancers in population may have changed over time. The results need to be verified by studies in future.

Conflict of interests

The authors declared no conflicting interests.

REFERENCES

1. Bratt O, Kristoffersson U, Lundgren R, et al. Sons of men with prostate cancer: their attitudes regarding possible inheritance of prostate cancer, screening, and genetic testing. *Urology* 1997; 50(3):360-5. doi: 10.1016/S0090-4295(97)00250-1.
2. Tao P, Hu YY, Huang Y, et al. Risk factors of breast cancer in Asian women: a meta-analysis. *Zhonghua Liu Xing Bing Xue Za Zhi* 2011; 32(2):164-9. Chinese.
3. Johns LE, Houlston RS. A systematic review and meta-analysis of familial colorectal cancer risk. *Am J Gastroenterol* 2001; 96(10):2992-3003. doi: 10.1111/j.1572-0241.2001.04677.x.
4. Kiciński M, Vangronsveld J, Nawrot TS. An epidemiological reappraisal of the familial aggregation of prostate cancer: A meta-analysis. *PLoS One* 2011; 6(10):1-7. doi: 10.1371/journal.pone.0027130.
5. Hemminki K, Dong C, Vaittinen P. Cancer risks to spouses and offspring in the Family-Cancer Database. *Genet Epidemiol* 2001; 20(2):247-57. doi: 10.1002/1098-2272(200102)20:2<247::AID-GEPI7>3.0.CO;2-U.
6. Hemminki K, Vaittinen P. Effect of paternal and maternal cancer on cancer in the offspring: A population-based study. *Cancer Epidemiol Biomarkers Prev* 1997; 6(12):993-7.
7. Chen W, Zheng R, Zhang S, et al. Cancer incidence and mortality in China, 2013. *Cancer Lett* 2017; 401:63-71. doi: 10.1016/j.canlet.2017.04.024.
8. Liu J, Shu T, Chang S, et al. Risk of malignancy associated with a maternal family history of cancer. *Asian Pac J Cancer Prev* 2014; 15(5):2039-44. doi: 10.7314/apjcp.2014.15.5.2039.
9. Liu J, Li N, Chang S, et al. Characteristics of 240 Chinese father-child pairs with malignant disease. *Asian Pac J Cancer Prev* 2014; 14(11):6501-5. doi: 10.7314/apjcp.2013.14.11.6501.
10. Lichtenstein P, Holm NV, Verksalo PK, et al. Environmental and heritable factors in the causation of cancer—analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 2000; 343(2):78-85. doi: 10.1016/S0039-6257(00)00165-x.
11. Liu S, Zhang M, Yang L, et al. Prevalence and patterns of tobacco smoking among Chinese adult men and women: Findings of the 2010 national smoking survey. *J Epidemiol Community Health* 2017; 71(2):154-61. doi: 10.1136/jech-2016-207805.
12. Li S, Meng L, Chioloro A, et al. Trends in smoking prevalence and attributable mortality in China, 1991–2011. *Prev Med* 2016; 93(12):82-7. doi: 10.1016/j.ypmed.2016.09.027.
13. Conteduca V, Sansonno D, Lauletta G, et al. H. pylori infection and gastric cancer: state of the art (review). *Int J Oncol* 2013; 42(1):5-18. doi: 10.3892/ijo.2012.1701.
14. Li L, Ying XJ, Sun TT, et al. Overview of methodological quality of systematic reviews about gastric cancer risk and protective factors. *Asian Pac J Cancer Prev* 2012; 13(5):2069-79. doi: 10.7314/apjcp.2012.13.5.2069.
15. Berretta M, Lleshi A, Fisichella R, et al. The role of nutrition in the development of esophageal cancer: What do we know? *Front Biosci (Elite Ed)* 2012; 4:351-7. doi: 10.2741/382.
16. Wu QJ, Wu L, Zheng LQ, et al. Consumption of fruit and vegetables reduces risk of pancreatic cancer: Evidence from epidemiological studies. *Eur J Cancer Prev* 2016; 25(3):196-205. doi: 10.1097/CEJ.0000000000000171.
17. Korc M, Jeon CY, Edderkaoui M, et al. Tobacco and alcohol as risk factors for pancreatic cancer. *Best Pract Res Clin Gastroenterol* 2017; 31(5):529-36 doi: 10.1016/j.bpg.2017.09.001.
18. Yuen MF, Hou JL, Chutaputti A. Hepatocellular carcinoma in the Asia pacific region. *J Gastroenterol Hepatol* 2019; 24(3):346-53. doi: 10.1111/j.1440-1746.2009.05784.x.
19. Vargas AJ, Thompson PA. Diet and nutrient factors in colorectal cancer risk. *Nutr Clin Pract* 2012; 27(5):613-23. doi: 10.1177/0884533612454885.

135 个父母双亲和子女均患癌症的 中国家庭发病特征回顾性分析

刘炬, 徐志坚*, 毕晓峰, 孙萍, 黄佳琴

防癌体检中心, 国家癌症中心 / 国家肿瘤临床医学研究中心 /
中国医学科学院北京协和医学院肿瘤医院, 北京 100021, 中国

摘要

目的 分析父母双亲和至少一个子女均患癌症的中国家庭的癌症发病特点, 为制定父母均患癌症人员的早期诊断和筛查方案提供参考。

方法 从国家癌症中心的防癌体检中心自 2008 年 1 月至 2018 年 2 月间体检人员家族史数据库中搜集父亲、母亲及至少一位子女均患癌的家庭。分析患癌成员的癌症病种及所占比例、诊断年龄的分布特点, 子女与父亲或母亲所患癌种及癌症所来源生理系统的相似性。将父亲组、母亲组、女儿组和儿子组中不同肿瘤的发病占比与国家癌症中心肿瘤登记办公室 (NCCRC) 2013 年统计的全国数据进行比较。

结果 从 33 200 名体检人员癌症家族史数据库中搜集 135 个父母双亲与子女均患癌症的家庭纳入研究。肺癌在父亲组 (40/135, 29.6%) 和母亲组 (38/135, 28.1%) 患癌中的占比明显高于 NCCRC 数据肺癌在男性 (23.9%) 及女性 (14.9%) 患癌中的占比。乳腺癌在女儿组患癌的占比 (35/109, 32.1%) 明显高于在母亲组 (14/135, 10.4%) 和 NCCRC 数据乳腺癌在女性患癌中的占比 (17.1%)。71 对父子中, 患同种癌症的占 16.9% (12/71), 患癌来源于同一系统的占 46.5% (33/71)。这两个指标在 109 对父女中分别是 31.2% ($n=34$) 和 10.1% ($n=11$); 在 71 对母子中分别为 36.6% ($n=26$) 和 8.5% ($n=6$); 在 109 对母女中分别为 31.2% ($n=34$) 和 20.2% ($n=20$)。父子患来源于同一系统的恶性肿瘤的可能性高于父女 ($X^2=4.299$, $P<0.05$), 而母女患同一种恶性肿瘤的可能性高于母子 ($X^2=4.506$, $P<0.05$)。罹患癌症的诊断年龄均数在儿子组 [(59.4 ± 10.9) 岁] 和女儿组 [(52.4 ± 12.7) 岁] 明显低于父亲组 [(65.5 ± 12.2) 岁] 和母亲组 [(65.7 ± 12.5) 岁] (均 $P<0.001$)。

结论 父母双亲均患癌的人员在进行癌症早诊或筛查时, 应进行多种癌症的全面检查, 而不应仅检查与父亲或者母亲同样的恶性肿瘤或者来源于同一系统的恶性肿瘤, 并且筛检的开始年龄应较针对一般人群建议的开始年龄提前。男性个体需尤其重视消化系统来源恶性肿瘤的筛检, 而女性个体应重点筛查女性特有癌症 (乳腺癌、卵巢癌、宫颈癌和子宫癌), 尤其应重视乳腺癌的筛查。

关键词: 家族史; 早期诊断; 父母; 儿子; 女儿; 恶性肿瘤

基金: 国家重点研发计划重大慢性非传染性疾病防控研究专项 (2017YFC1308700); 北京协和医学院学科建设项目 (201920200303)。