ORIGINAL ARTICLE

Effect of Focused Cardiac Ultrasound in Combination with Lung Ultrasound on Critically Ill Patients: A Multicenter Observational Study in China

Hongmin Zhang¹, Lina Zhang², Lixia Liu³, Ying Zhu⁴, Wanhong Yin⁵, Wei He⁶,
Xiuling Shang⁷, Yangong Chao⁸, Liwen Lv⁹, Xiaoting Wang^{1*}, Dawei Liu¹;
Chinese Critical Ultrasound Study Group

¹Department of Critical Care Medicine, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100730, China

²Department of Critical Care Medicine, Xiangya Hospital, Central South University, Changsha 410013, China

³Department of Critical Care Medicine, The Fourth Hospital of Hebei Medical University, Shijiazhuang 050011, China

⁴Department of Critical Care Medicine, Affiliated Hangzhou First People's Hospital, Zhejiang University School of Medicine, Hangzhou 310006, China

⁵Department of Critical Care Medicine, West China Hospital, Sichuan University, Chengdu 610041, China

⁶Department of Critical Care Medicine, Beijing Tongren Hospital, Capital Medical University, Beijing 100730, China

⁷Department of Critical Care Medicine, Fujian Provincial Hospital, Fuzhou 350001, China

⁸Department of Critical Care Medicine, The First Hospital of Tsinghua University, Beijing 100016, China

⁹Department of Emergency Medicine, The People's Hospital of Guangxi Zhuang Autonomous Region, Nanning 530021, China

Key words: cardiac ultrasound; lung ultrasound; critically ill; point-of-care

Objective Focused cardiac ultrasound (FCU) and lung ultrasound (LU) are increasingly being used in critically ill patients. This study aimed to investigate the effect of FCU in combination with LU on these patients and to determine if the timing of ultrasound examination was associated with treatment change.

Methods This is a multicenter cross-sectional observational study. Consecutive patients admitted to the intensive care unit (ICU) were screened for enrollment. FCU and LU were performed within the first 24 h, and treatment change was proposed by the performer based on the ultrasound results and other clinical conditions.

Results Among the 992 patients included, 502 were examined within 6 h of ICU admission (early phase group), and 490 were examined after 6 h of admission (later phase group). The early phase group and the later phase group had similar proportions of treatment change (48.8% vs. 49.0%, $\chi^2=0.003$, $P=0.956$). In the multivariable analysis, admission for respiratory failure was an independent variable associated with treatment change, with an odds ratio (OR) of 2.357 [95% confidence interval (CI): 1.284-4.326, $P=0.006$]; the timing of examination was not associated with treatment change (OR=0.725, 95%CI: 0.407-1.291, $P=0.275$).

Conclusions FCU in combination with LU, whether performed during the early phase or later phase, had a significant impact on the treatment of critically ill patients. Patients with respiratory failure were more likely to experience treatment change after the ultrasound examination.

CRITICAL care physicians can be effectively trained on point-of-care ultrasound in terms of image acquisition and interpretation.^[1, 2] The ultrasound training permits the physicians to apply the results of the imaging study to the clinical situation for purposes of immediate diagnosis and patient management.^[2, 3]

To date, most studies regarding the effect of point-of-care ultrasound in critically ill patients were conducted either on one single type of ultrasound, such as focused cardiac ultrasound (FCU) and point-of-care lung ultrasound (LU), or on a full-scale ultrasound examination from head to toe.^[4-7] It is not feasible to examine every critically ill patient with a head-to-toe approach because it is time consuming. Sekiguchi *et al.* noted that FCU was a valuable tool during resuscitation of septic shock patients.^[8] FCU is strongly recommended in any patient presenting with shock.^[9] LU is readily available at the bedside to monitor the critically ill and can be used to identify the cause of respiratory failure. In addition, it helps titrate the ventilatory support.^[10-12] Moreover, LU can also be of assistance in the management of decompensated heart failure patients by monitoring B-lines from pulmonary congestion.^[13, 14] FCU can also help in the diagnosis and management of respiratory failure patients, including patients with pulmonary embolism, acute respiratory distress syndrome (ARDS), or cardiogenic pulmonary edema.^[15-17] We hypothesized that FCU in combination with LU as a

screening protocol could lead to significant change in the management of intensive care unit (ICU) patients.

The timing of ultrasound examination varied greatly. A recent published study noted that FCU is a valuable examination tool during early resuscitation of these patients.^[8] Some prior studies found that ultrasound performed within 12 or 24 h of admission could also benefit the patients.^[5, 18] However, whether the ultrasound performed at the early resuscitation phase or after that would result in different treatment changes has not been discussed.

In this study, we aimed to investigate the effect of FCU in combination with LU on critically ill patients. We would also assess whether the timing of ultrasound examination can affect the treatment change in these patients.

PATIENTS AND METHODS

This is a cross-sectional observational study conducted in 8 Chinese university-affiliated hospital ICU from March 15 to June 15, 2018, and 5 of the 8 ICUs were surgical ICUs and 3 were general ICUs. The examiners were all credited by Chinese Critical Ultrasound Study Group (CCUSG) as trainers with at least a year or more of experience in critical care echocardiography practice.

Consecutive patients admitted to ICU from April to May 31, 2018 were screened for enrollment within

the first 24 h of admission. The exclusion criteria included inadequate echocardiographic imaging for measurement or lack of consent form.

This study was approved by the institutional committee of Peking Union Medical College Hospital (Approval No. ZS1422). Written informed consent was obtained from the patients or the next of kin.

Focused cardiac ultrasound and lung ultrasound

Ultrasound examination was completed within the first 24 h of ICU admission. The patients were in the semi left lateral position during the FCU examination and were in supine position during the LU examination. The images were recorded for offline analysis. The heart rate (HR), mean arterial pressure (MAP), norepinephrine dose, and mechanical ventilation parameters were recorded at the onset of ultrasound examination.

Structure abnormality included pericardial effusion, atrial or ventricular dilatation, ventricular hypertrophy, and obvious valvular abnormalities on 2D ultrasound.

Left ventricular ejection fraction was estimated through the eyeballing method with three categories: normal ($\geq 55\%$), moderately abnormal ($30\%-55\%$) and severely abnormal ($<30\%$).^[19] Right ventricle abnormal was defined by two signs: right ventricle enlargement and paradoxical septal motion.^[20]

Inferior vena cava (IVC) was measured from the subcostal view where the long axis was displayed. The measurement of IVC was at the end-expiration and just proximal to the junction of the hepatic veins. Dilated and fixed IVC: internal diameter >2 cm and no variation with breathing. Small and collapsed IVC: internal diameter <1 cm and obvious variation with breathing. The rest was categorized into moderate IVC.^[21, 22]

LU was performed based on the BLUE protocol.^[6] The A profile indicated anterior lung-sliding with A-lines. B-profile was associated anterior lung-sliding with lung-rockets. The C-profile indicated anterior lung consolidation regardless of size and number. The A'-profile was an A-profile with abolished lung-sliding. The B'-profile was a B-profile with abolished lung-sliding. The A/B profile was a half A-profile at one lung and a half B-profile at another. The PLAPS-profile indicated Posterolateral Alveolar and/or Pleural Syndrome.^[23]

Ultrasound performers were required to be familiar with the patients' medical history, the status of cir-

culatory and respiratory support, and the parameters regarding organ function and tissue perfusion before the ultrasound examination. After the ultrasound examination, the performers were asked to decide two aspects: 1. The ultrasonic diagnosis on each item of FCU and LU; 2. The treatment change proposal based on the patients' clinical and ultrasonic results.

Treatment change

Treatment changes included volume expansion or diuresis, vasopressor or inotrope adjustment, drainage of pericardial effusion, ventilator adjustment, drainage of pleural effusion or pneumothorax, in need of other type of ultrasound examination, in need of computerized tomography, and in need of continuous cardiac output monitoring. The treatment change proposal was made by the ultrasound performer, but the treatment application was at the discretion of the attending physician in charge of the patients.

Other parameters collected

Demographic information and the diagnosis, Acute Physiology and Chronic Health Evaluation (APACHE) II score, Sequential Organ Failure Assessment (SOFA) score, comorbidities, vasopressor and inotrope dose were collected for all patients. Each patient's HR, MAP and whether on mechanical ventilation were also collected at the time of the echo examination.

Statistical analysis

Statistical analysis was performed using the SPSS 13.0 statistical software package (SPSS Inc., Chicago, Illinois, USA). Normal distribution of the continuous variables was assessed by the Kolmogorov-Smirnov test. Continuous data were expressed as the mean $\pm$ standard deviation (SD) when normally distributed. Not normally distributed variables were expressed as median and interquartile range. Categorical variables were presented as the number and percentages. Differences among groups were assessed by Student's unpaired *t* test, the Mann-Whitney *U* test, the Kruskal-Wallis test, the Chi-squared test, or Fisher's exact test, as appropriate. We performed a logistic regression analysis to assess the independent factors of treatment change. The following variables were considered for the multivariable analysis: age, APACHE II score, SOFA score, admission for respiratory failure, admission for shock, and the timing of ultrasound examination. Two tailed $P < 0.05$ was considered significant.

RESULTS

General characteristics of the patients

A total of 1133 patients were screened for enrollment in the study, and 992 with adequate image quality were included (**Fig. 1**). The early phase group included 502 patients who were examined within 6 h of ICU admission. The remaining 490 patients who were examined between 6 h and 24 h of admission were classified into the later phase group. The general characteristics of the two groups of patients were comparable (**Table 1**). The reasons for admission to the ICU included high risk surgery, respiratory failure, shock and other rea-

sons (acute kidney injury, seizure, stroke, liver failure, trauma, and poisoning).

Results of ultrasound examination

The median duration of examination in both groups was 15 minutes. No difference was found between the two groups in terms of patients with structure abnormality, impaired left ventricular systolic function and right ventricular dysfunction. The early phase group had more patients with dilated and fixed IVC and had fewer patients with moderate IVC (34.5% vs. 26.3%, $\chi^2=7.751$, $P=0.005$ and 47.8% vs. 56.3%, $\chi^2=7.208$, $P=0.007$, respectively). No difference was found between the two

Figure 1. Flow chart showing patient selection and treatment change.

Table 1. Comparisons of features of the two groups of patients

Items	Early phase (n=502)	Later phase (n=490)	t or Z or χ^2	P
Male [n (%)]	287 (57.2)	301 (61.4)	1.862	0.172
Age (yr, mean $\pm$ SD)	61 $\pm$ 18	59 $\pm$ 17	1.849	0.064
APACHE II	15 (11, 21)	17 (11, 23)	-0.924	0.270
SOFA [score, median (interquartile range)]	6 (4, 8)	6 (4, 8)	-0.662	0.508
Diagnosis [n (%)]				
High risk surgery	183 (36.5)	187 (37.8)	0.310	0.578
Respiratory failure	130 (25.9)	152 (31.0)	3.200	0.074
Shock	84 (16.7)	66 (13.5)	2.058	0.151
Shock+Respiratory failure	39 (7.8)	29 (5.9)	1.330	0.249
Others*	118 (23.5)	114 (23.3)	0.008	0.929
HR [bpm, median (interquartile range)]	92 (77, 114)	98 (79, 110)	-0.012	0.991
MAP [mmHg, median (interquartile range)]	88 (77, 102)	88 (80, 98)	-0.273	0.785
NE infusion [n, (%)]	103 (20.5)	115 (23.5)	0.725	0.124
MV [n, (%)]	264 (52.6)	279 (56.9)	1.893	0.169

*Others: acute kidney injury, seizure, stroke, trauma, liver failure, and poisoning. APACHE: Acute Physiology and Chronic Health Evaluation; SOFA: Sequential Organ Failure Assessment; HR: heart rate; MAP: mean arterial pressure; NE: norepinephrine; MV: mechanical ventilation.

groups in terms of patients with each LU profile. (**Table 2**)

Treatment change

We did not observe differences between early phase and later phase regarding volume expansion and ino-

trope. The early phase group had more patients with norepinephrine adjustment but had not statistically significant ($P=0.055$). The proportions of patients with treatment change in the two groups were 48.8% and 49.0% ($\chi^2=0.003$, $P=0.956$) (**Table 3**).

Table 2. Comparisons of results of ultrasound examination [n (%)]

Items	Early phase ($n=502$)	Later phase ($n=490$)	Z or χ^2	P
Duration of examination [min, median (interquartile range)]	15 (11, 30)	15 (12, 30)	-1.495	0.135
FCU				
Structure abnormality*	92 (18.3)	94 (19.2)	0.120	0.730
Left ventricular systolic function				
Normal	364 (72.5)	343 (70.0)	0.763	0.382
Moderately abnormal	120 (23.9)	132 (26.9)	1.205	0.272
Severely abnormal	18 (3.6)	15 (3.1)	0.212	0.645
IVC				
Dilated and fixed	173 (34.5)	129 (26.3)	7.751	0.005
Moderate	240 (47.8)	276 (56.3)	7.208	0.007
Small and collapsed	89 (17.7)	85 (17.3)	0.025	0.874
RV dysfunction	7 (1.4)	9 (1.8)	0.306	0.580
LU				
A profile	212 (42.2)	181 (36.9)	2.903	0.088
A' or B'+lung point	8 (1.6)	11 (2.2)	0.560	0.454
B profile	35 (7.0)	29 (5.9)	0.456	0.499
Other profiles*	247 (49.2)	269 (54.9)	3.222	0.073

*Obvious abnormality: pericardial effusion, valvulopathy, atrial or ventricular dilatation, ventricular hypertrophy; Other profiles: A/B, C, A+P, B+P, A/B+P, and C+P profile.

A profile: anterior lung-sliding with A-lines; B-profile: anterior lung-sliding with lung-rockets; C-profile: anterior lung consolidation; A'-profile: an A-profile with abolished lung-sliding; B'-profile: a B-profile with abolished lung-sliding; A/B profile: a half A-profile at one lung and a half B-profile at another; P profile: posterolateral alveolar and/or pleural syndrome; FCU: focused cardiac ultrasound; IVC: inferior vena cava; RV: right ventricle; LU: lung ultrasound.

Table 3. Comparison of treatment change between the two groups [n (%)]

Items	Early phase ($n=502$)	Later phase ($n=490$)	χ^2	P
Patients with treatment change	245 (48.8)	240 (49.0)	0.003	0.956
One treatment change	153 (30.5)	149 (30.4)	0.001	0.991
More than one treatment change	92 (18.3)	91 (18.6)	0.010	0.921
Volume expansion	47 (9.4)	44 (9.0)	0.044	0.835
Negative fluid balance	166 (33.1)	139 (28.4)	2.573	0.109
Inotrope	28 (5.6)	39 (8.0)	2.233	0.135
Norepinephrine adjustment	37 (7.4)	22 (4.5)	3.679	0.055
Drainage of pericardial effusion	3 (0.6)	0	2.937	0.087
Ventilator adjustment	93 (18.5)	104 (21.2)	1.135	0.287
Drainage of pleural cavity	23 (4.6)	24 (4.9)	0.055	0.815
In need of other type of ultrasound examination	91 (18.1)	97 (19.8)	0.449	0.503
In need of CT	88 (17.5)	98 (20.0)	0.993	0.319
In need of CCO monitoring	39 (7.8)	26 (5.3)	2.456	0.117

CT: computerized tomography; CCO: continuous cardiac output.

In the multivariable analysis, admission for respiratory failure was an independent variable associated with treatment changes, with an odds ratio (OR) of 2.357 [95% confidence interval (CI): 1.284-4.326], $P=0.006$; the timing of examination was not associated with treatment change (OR=0.725, 95%CI: 0.407-1.291, $P=0.275$) (Table 4).

DISCUSSION

In the present study, we investigated the effect of FCU in combination with LU on ICU patients when performed during the first 24 h of admission. The assessment of treatment change was made by the physicians who performed the ultrasound and were also familiar with the patients' clinical conditions. After the ultrasound examination, a large proportion of patients had treatment change; the later phase group had similar proportions of treatment change compared to the early phase group.

This study found that the shock and respiratory failure patients accounted for approximately 50% of total admissions. Furthermore, 6%-8% of patients had both shock and respiratory failure. This could partly explain why the combination of FCU and LU was suitable for the management of these patients. According to a recent survey involving 142 ICUs in Europe, FCU and LU are the top two types of ultrasound that are most frequently performed.^[24] We also found that the need for other types of ultrasound was not as high.

FCU and LU can play complementary roles in the management of circulatory shock and respiratory failure patients. LU can play an important role in volume management. Left ventricle dysfunction is very common in critically ill patients and can lead to pulmonary edema.^[8, 18] LU can be used as a reliable and easy way to evaluate pulmonary congestion through assessment

of B-lines.^[25] Lee *et al.* suggests using the inferior vena cava and LU to guide fluid resuscitation.^[26] Lichtenstein, who created the BLUE protocol, also suggests that LU can be integrated into the management of fluid administration and shock management.^[27] In this study, more patients in the early examination group had dilated IVC, yet the number of patients requiring a negative balance was similar in the two groups. One explanation can be that the decision was made based on the results of both FCU and LU. FCU can also benefit the treatment of respiratory failure patients. ARDS is not a rare syndrome, as it is a widespread disease with a massive socio-economic impact.^[28] ARDS is one of the most common conditions challenging the right ventricle. The incidence of acute cor pulmonale in ARDS patients is approximately 25%, even with lung protection ventilation. Cardiac ultrasound is usually recommended for diagnosing acute cor pulmonale.^[16, 29] Therefore, the treatment change of ventilator adjustment might be made based on the result of FCU and LU rather than solely on LU.

The time spent on this protocol was also desirable for the management of critically ill patients. A prior study reported that a limited echocardiography could be completed within 10 minutes for each patient.^[5] In the present study, performers spent a median time of 15 minutes on each ultrasound examination. Thus, adding LU will not be time consuming. However, if all types of ultrasound were performed, the time spent for examination would be much longer, which would not be practical in the management of critically ill patients. Therefore, in examining ICU patients, the FCU+LU can be the main screening protocol with other types of ultrasound that can be added according to a patient's clinical condition.

In the present study, the treatment change between the early examination and later examination groups was similar. The results indicated that the timing of examination had limited effect on the decision of treatment change. The early phase resuscitation could not supplant the treatment after the early phase of the critically ill. Vincent *et al.* identified at least four phases in the treatment of shock patients.^[9] The results suggested that ultrasound performed after the early resuscitation phase could still benefit the critically ill. This is consistent with a prior study, which suggested that repeated monitoring with transthoracic echocardiography and LU changed the diagnosis of cardiac and/or respiratory disorders in 67% of post cardiac

Table 4. Results of logistic regression analysis for factors associated with treatment change

Risk factors	OR	95%CI	P
Age	0.987	0.969-1.005	0.157
APACHE II	0.997	0.991-1.003	0.379
SOFA	1.034	0.946-1.130	0.465
Respiratory failure	2.357	1.284-4.326	0.006
Shock	1.733	0.896-3.353	0.103
Timing of examination	0.725	0.407-1.291	0.275

surgery patients.^[30] Therefore, critical care ultrasound should be seen not only as a diagnostic tool, but also as a monitoring tool. Nevertheless, the ultrasound examination in this study was performed either during the early phase or after that, thus we were uncertain that we can get the similar result if we perform the ultrasound examination both in the early phase and the later phase. A future study is warranted to help elucidate this hypothesis.

There were several limitations in this study. First, it is a cross-sectional study, and thus the causal effect of FCU in combination with LU needs to be confirmed by a prospective cohort or even randomized clinical trials. Second, the decisions of treatment change were made subjectively by the clinician who performed the ultrasound. However, all the performers were certified through formal training, and had enough experience in FCU and LU. Third, this research was done as a screening approach rather than problem-oriented. Therefore, the examination timing was chosen by the performer instead of the patients' need. However, we still found as much information regarding the change of treatment, which added value to the FCU and LU examination.

In conclusion, FCU in combination with LU, whether performed during the early phase or later phase, had a significant impact on the treatment of critically ill patients. Patients with respiratory failure were more likely to experience treatment change after the ultrasound examination.

ACKNOWLEDGEMENTS

We would like to thank Prof. Fuhai Shen, from Department of Epidemiology North China University of Science and Technology, for his kind suggestions on the statistical issue. We would also like to thank Dr. Li Li, Chenghuan Hu, Meilin Ai, Qianyi Peng from Xiangya University, Dr. Daqian Zhang from the Affiliated Hospital of Guizhou Medical University, Hua Zhao, Huan Chen from Peking Union Medical College, Yang Liu, Na Liu, from Beijing Tongren Hospital, Shanmu Ai, from Daping Hospital, Minjia Wang, from Zhejiang Hospital, and Ran Zhu from the First Hospital of China Medical University for their precious effort in the data collection.

Conflicts of Interest Statement

The authors declare that they have no competing interests.

REFERENCES

- Greenstein YY, Littauer R, Narasimhan M, et al. Effectiveness of a critical care ultrasonography course. *Chest* 2017; 151(1):34-40. doi: 10.1016/j.chest.2016.08.1465.
- Dinh VA, Giri PC, Rathinavel I, et al. Impact of a 2-day critical care ultrasound course during fellowship training: a pilot study. *Crit Care Res Pract* 2015; 2015:675041. doi: 10.1155/2015/675041.
- Via G, Hussain A, Wells M, et al. International evidence-based recommendations for focused cardiac ultrasound. *J Am Soc Echocardiogr* 2014; 27(7):683.e1-683.e33. doi: 10.1016/j.echo.2014.05.001.
- Andruszkiewicz P, Sobczyk D, Gorkiewicz-Kot I, et al. Reliability of focused cardiac ultrasound by novice sonographer in preoperative anaesthetic assessment: an observational study. *Cardiovasc Ultrasound* 2015; 13:45. doi: 10.1186/s12947-015-0039-y.
- Kanji HD, McCallum J, Sirounis D, et al. Limited echocardiography-guided therapy in subacute shock is associated with change in management and improved outcomes. *J Crit Care* 2014; 29(5):700-5. doi: 10.1016/j.jcrc.2014.04.008.
- Lichtenstein D. Lung ultrasound in the critically ill. *Curr Opin Crit Care* 2014; 20(3):315-22. doi: 10.1097/MCC.0000000000000096.
- Manno E, Navarra M, Faccio L, et al. Deep impact of ultrasound in the intensive care unit: the "ICU-sound" protocol. *Anesthesiology* 2012; 117(4):801-9. doi: 10.1097/ALN.0b013e318264c621.
- Sekiguchi H, Harada Y, Villarraga HR, et al. Focused cardiac ultrasound in the early resuscitation of severe sepsis and septic shock: a prospective pilot study. *J Anesth* 2017; 31(4):487-93. doi: 10.1007/s00540-017-2312-8.
- Vincent JL, De Backer D. Circulatory shock. *N Engl J Med* 2013; 369(18):1726-34. doi: 10.1056/NEJMr1208943.
- Monastesse A, Girard F, Massicotte N, et al. Lung ultrasonography for the assessment of perioperative atelectasis: a pilot feasibility study. *Anesth Analg* 2017; 124(2):494-504. doi: 10.1213/ANE.0000000000001603.
- Theerawit P, Sutherasan Y, Ball L, et al. Respiratory monitoring in adult intensive care unit. *Expert Rev Respir Med* 2017; 11(6):453-68. doi: 10.1080/17476348.2017.1325324.
- Via G, Storti E, Gulati G, et al. Lung ultrasound in the

- ICU: from diagnostic instrument to respiratory monitoring tool. *Minerva Anestesiol* 2012; 78(11):1282-96.
13. Volpicelli G, Caramello V, Cardinale L, et al. Bedside ultrasound of the lung for the monitoring of acute decompensated heart failure. *Am J Emerg Med* 2008; 26(5):585-91. doi: 10.1016/j.ajem.2007.09.014.
 14. Martindale JL, Secko M, Kilpatrick JF, et al. Serial sonographic assessment of pulmonary edema in patients with hypertensive acute heart failure. *J Ultrasound Med* 2018; 37(2):337-45. doi: 10.1002/jum.14336.
 15. Tsou PY, Kurbedin J, Chen YS, et al. Accuracy of point-of-care focused echocardiography in predicting outcome of resuscitation in cardiac arrest patients: a systematic review and meta-analysis. *Resuscitation* 2017; 114:92-9. doi: 10.1016/j.resuscitation.2017.02.021.
 16. Repessé X, Charron C, Vieillard-Baron A. Acute cor pulmonale in ARDS: rationale for protecting the right ventricle. *Chest* 2015; 147(1):259-65. doi: 10.1378/chest.14-0877.
 17. Vignon P, Repessé X, Vieillard-Baron A, et al. Critical care ultrasonography in acute respiratory failure. *Crit Care* 2016; 20(1):228. doi: 10.1186/s13054-016-1400-8.
 18. Marcelino PA, Marum SM, Fernandes AP, et al. Routine transthoracic echocardiography in a general Intensive Care Unit: an 18 month survey in 704 patients. *Eur J Intern Med* 2009; 20(3):e37-42. doi: 10.1016/j.ejim.2008.09.015.
 19. Randazzo MR, Snoey ER, Levitt MA, et al. Accuracy of emergency physician assessment of left ventricular ejection fraction and central venous pressure using echocardiography. *Acad Emerg Med* 2003; 10(9):973-7.
 20. Vieillard-Baron A. Assessment of right ventricular function. *Curr Opin Crit Care* 2009; 15(3):254-60. doi: 10.1097/MCC.0b013e32832b70c9.
 21. Zhang Z, Xu X, Ye S, et al. Ultrasonographic measurement of the respiratory variation in the inferior vena cava diameter is predictive of fluid responsiveness in critically ill patients: systematic review and meta-analysis. *Ultrasound Med Biol* 2014; 40(5):845-53. doi: 10.1016/j.ultrasmedbio.2013.12.010.
 22. Rudski LG, Lai WW, Afilalo J, et al. Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology, and the Canadian Society of Echocardiography. *J Am Soc Echocardiogr* 2010; 23(7):685-713; quiz 786-8. doi: 10.1016/j.echo.2010.05.010.
 23. Lichtenstein DA. Lung ultrasound in the critically ill. *Ann Intensive Care* 2014; 4(1):1. doi: 10.1186/2110-5820-4-1.
 24. Zieleskiewicz L, Muller L, Lakhal K, et al. Point-of-care ultrasound in intensive care units: assessment of 1073 procedures in a multicentric, prospective, observational study. *Intensive Care Med* 2015; 41(9):1638-47. doi: 10.1007/s00134-015-3952-5.
 25. Miglioranza MH, Picano E, Badano LP, et al. Pulmonary congestion evaluated by lung ultrasound predicts decompensation in heart failure outpatients. *Int J Cardiol* 2017; 240:271-8. doi: 10.1016/j.ijcard.2017.02.150.
 26. Lee CW, Kory PD, Arntfield RT. Development of a fluid resuscitation protocol using inferior vena cava and lung ultrasound. *J Crit Care* 2016; 31(1):96-100. doi: 10.1016/j.jcrc.2015.09.016.
 27. Lichtenstein D. FALLS-protocol: lung ultrasound in hemodynamic assessment of shock. *Heart Lung Vessel* 2013; 5(3):142-7.
 28. Lewandowski K, Lewandowski M. Epidemiology of ARDS. *Minerva Anestesiol* 2006; 72(6):473-7.
 29. Krishnan S, Schmidt GA. Acute right ventricular dysfunction: real-time management with echocardiography. *Chest* 2015; 147(3):835-46. doi: 10.1378/chest.14-1335.
 30. Alsaddique A, Royse AG, Royse CF, et al. Repeated monitoring with transthoracic echocardiography and lung ultrasound after cardiac surgery: feasibility and impact on diagnosis. *J Cardiothorac Vasc Anesth* 2016; 30(2):406-12. doi: 10.1053/j.jvca.2015.08.033.

中文摘要

床旁心脏超声结合肺部超声在重症患者中的应用： 一项全国多中心观察性研究

张宏民¹，张丽娜²，刘丽霞³，朱英⁴，尹万红⁵，何伟⁶，尚秀玲⁷，晁彦公⁸，

吕立文⁹，王小亭^{1*}，刘大为¹，重症超声研究组

¹ 中国医学科学院北京协和医学院 北京协和医院，重症医学科，中国，北京，100730

² 中南大学湘雅医院，重症医学科，中国，长沙，410013

³ 河北医科大学第四医院，重症医学科，中国，石家庄，050011

⁴ 浙江大学医学院附属杭州第一人民医院，重症医学科，中国，杭州，310006

⁵ 四川大学华西医院，重症医学科，中国，成都，610041

⁶ 北京同仁医院，重症医学科，中国，北京，100730

⁷ 福建省立医院，重症医学科，中国，福建，350001

⁸ 清华大学第一附属医院，重症医学科，中国，北京，100016

⁹ 广西壮族自治区人民医院，急诊科，中国，南宁，530016

摘要

目的 床旁心脏超声和肺部超声在重症患者中的应用日益广泛。本研究旨在调查心脏超声联合肺部超声在重症患者中的作用，以及超声检查时机与治疗改变是否有关。

方法 本研究为多中心横断面观察性研究。连续收入重症医学科住院患者，在 24 h 内行床旁心脏超声和肺部超声检查，操作者根据患者的临床情况和超声检查结果做出是否需要改变治疗的判断。

结果 在收入的 992 例患者中，502 例患者（早期组）在入科 6 h 内完成超声检查，490 例患者（后期组）在入科 6 h 后完成超声检查。早期组和后期组治疗改变的比例接近（48.8% 比 49.0%， $\chi^2=0.003$ ， $P=0.956$ ）。多元分析表明，因呼吸衰竭入院是治疗改变的独立影响因素（ $OR=2.357$ ，95% 置信区间：1.284-4.326， $P=0.006$ ）；超声检查时机不是治疗改变的影响因素（ $OR=0.725$ ，95% 置信区间：0.407-1.291， $P=0.275$ ）。

结论 床旁心脏超声结合肺部超声检查能显著改变重症患者的治疗，而且 24h 内早期检查和后期检查均有作用。超声检查更容易给呼吸衰竭患者带来治疗改变。

关键词：心脏超声；肺部超声；重症患者；床旁