ORIGINAL ARTICLE

Aortic Valve Replacement for Patients with Heyde Syndrome: A Literature Review

Huili Li¹, Peishuang Lin², Yuntai Yao^{3*}

¹Department of Anesthesiology, The First Hospital of Hebei Medical University, Shijiazhuang 050000, China

²Department of Anesthesiology, Fujian Medical University Affiliated First Quanzhou Hospital, Quanzhou 362000, China

³Department of Anesthesiology, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100037, China

Key words: Heyde syndrome; gastrointestinal bleeding; aortic valve stenosis

Objective To explore the characteristics and clinical outcomes of patients with Heyde syndrome (HS) who undergo aortic valve replacement (AVR).

Methods Electronic databases including PubMed, Embase, Ovid, WANFANG, VIP and CNKI were searched to identify all case reports of HS patients undergoing AVR surgery, using different combinations of search terms “Heyde syndrome”, “gastrointestinal bleeding”, “aortic stenosis”, and “surgery”. Three authors independently extracted the clinical data including the patients’ characteristics, aortic stenosis severity, gastrointestinal bleeding sites, surgical treatments and prognosis.

Results Finally, 46 case reports with 55 patients aging from 46 to 87 years, were determined eligible and included. Of them, 1 patient had mild aortic stenosis, 1 had moderate aortic stenosis, 42 had severe aortic stenosis, and 11 were not mentioned. Gastrointestinal bleeding was detected in colon ($n=8$), jejunum ($n=6$), ileum ($n=4$), cecum ($n=3$), duodenal ($n=3$) and multiple sites ($n=8$). No specific bleeding site was identified in 23 patients. Preoperative hemoglobin level ranged from 43 to 117 g/L. All but one of 16 patients showed decreased level of high molecule weight von Willebrand factor. Of the 55 patients, 43 underwent AVR, and 12 received transcatheter AVR. Aortic valves of 14 cases were replaced by mechanical valves, and 33 cases by biological valves. All patients recovered well during the follow-up, except 5 patients. One patient who had perivalvular leakage and gastrointestinal bleeding after AVR underwent the second AVR. Two patients had recurrent gastrointestinal bleeding. Two patients died of life-threatening acute subdural hematoma and multiple organ failure, respectively.

Conclusions HS is a rare syndrome characterized by aortic stenosis and gastrointestinal bleeding. AVR is an effective treatment for HS.

HEYDE syndrome (HS) was first described in 1958 by Edward Heyde^[1] in a letter written to *The New England Journal of Medicine* as the triad of aortic stenosis (AS), acquired coagulopathy, and anemia due to bleeding from intestinal angiodysplasia. The prevalence of AS increases with age: 1.3% patients between 65 and 75 years, 2.4% between 75 and 85 years, and 4% over 85 years.^[2] AS is associated with coagulation dysfunction and bleeding tendency (bleeding occurs in about 20% of patients with AS^[3]), which has been repeatedly reported. The incidence of gastrointestinal bleeding is reported to be 1%-3%.^[4] Due to the lack of understanding of this disease by clinicians, patients with a small amount of gastrointestinal bleeding may not be further examined. And with different degrees of missed diagnosis, it is difficult to accurately determine the incidence of HS. Aortic valve replacement (AVR) seems to be an effective method for HS treatment. The current study presents a literature review of published cases reporting the clinical characteristics and prognosis of HS patients undergoing AVR.

MATERIALS AND METHODS

Study search

A review of case reports was performed to identify relevant articles published before September 9, 2020 by searching the electronic databases of PubMed, Embase, and Ovid. Chinese literatures from CNKI, VIP, and WANFANG databases were also searched. Different combinations of terms that included "Heyde syndrome", "aortic stenosis", "gastrointestinal bleeding", "aortic valve replacement", and "surgery" were used in the search strategy (**Appendix Table 1**). All relevant case reports were included. Exclusion criteria included: (1) non-English and non-Chinese articles, (2) review articles, (3) studies based on animal models, (4) duplicate publications, and (5) studies lacking outcomes of interest. Each author independently reviewed the titles and abstracts of all identified reports to exclude obviously ineligible studies. The eligibility of those remaining reports for final inclusion was determined further by examining the full text. Disagreements were resolved by discussion among three authors during the process of data abstraction.

Data abstraction

The following data from the included case reports were abstracted by three authors independently: (1) author, year, country, age and sex; (2) AS severity, bleeding site, and hemoglobin; (3) high molecular weight (HMW) von Willebrand factor (vWF), operation, and follow-up.

RESULTS

As depicted in **Fig. 1**, we identified 55 potentially qualified articles through searching databases. Finally, 46 case reports (55 patients in total) were determined eligible and included. Forty-five were written in English, and one in Chinese. Descriptive analyses of these cases are presented in **Table 1**.^[5-50] These 55 patients aged between 46 to 87 years; 36 were female and 17 were male, and 2 reports failed to provide the gender of patients. All patients had various degrees of AS and gastrointestinal bleeding. The peak pressure gradients of aortic valve of 28 cases ranged from 32 to 162 mm Hg, and the mean pressure gradients of aortic valve of 24 cases reported from 22 to 103 mm Hg. Aortic valve area of 31 patients varied from 0.4 to 1.0 cm². Gastrointestinal endoscopy confirmed that the bleeding sites of 8 patients were located in the colon, 6 in the jejunum, 4 in the ileum, 3 in the cecum, 3 in the duodenum, and 8 in multiple organs. Gastrointestinal endoscopy did not identify the specific bleeding site of 23 patients. Preoperative hemoglobin level ranged from 43 to 117 g/L. A decreased level of HMW vWF factor was detected in all^[5, 9, 10, 12, 16-19, 22, 24, 34, 35, 37, 40] but one^[36] of the 16 patients who underwent this examination. Of the 55 patients, 43 underwent surgical aortic valve replacement (SAVR) and 12 transcatheter aortic valve replacement (TAVR). A total of 14 patients received mechanical valves and 33 biological valves. All patients recovered well during the follow-up, except 5 patients who developed various complications. One patient^[15] had perivalvular leakage after AVR followed by gastrointestinal bleeding, and the second AVR with biological valve stopped the bleeding. Gastrointestinal bleeding of 2 patients recurred.^[43] Two patients^[14, 17] died of life-threatening acute subdural hematoma and multiple organ failure, respectively.

DISCUSSION

HS is characterized by the association of gastrointestinal bleeding from angiodysplasia, aortic stenosis and acquired von Willebrand deficiency. In recent years, there have been increasing numbers of reports on this rare disease. However, the mechanism, clinical features and treatment of HS remains undetermined.

The current study suggested that HS occurred more frequently in the elderly and was more common in female. It has been reported more frequently in Asia and America. Transthoracic echocardiography showed severe AS (moderate AS was occasionally seen). Most HS patients presented with anemia as the first manifestation. Intestinal angiodysplasia occurred commonly in the duodenum and the right side of the colon. However, our current study found a greater risk of multiple bleeding, so it might be inappropriate for these patients to undergo intestinal resection.

Possible mechanisms of HS are diverse, including common pathophysiology and common cardiovascular risk factors of AS and vascular dysplasia, low cardiac output and cholesterol emboli caused by AS, or intestinal hypoxia caused by atherosclerotic plaques. But HMW vWF multimer deficiency is proposed as a key mechanism of gastrointestinal bleeding. Additionally,

AS has been shown to cause an acquired form of von Willebrand syndrome type 2A occurring from the degradation of vWF multimers by the shear stress across the narrowed valve, leading to loss of large multimers involved in hemostasis and bleeding in previously latent intestinal angiodysplasias. The HMW vWF levels of HS patients were lower except for one. Gel electrophoresis is the gold standard measurement of HMW vWF level when diagnosing HS, although it is costly and resource intensive.^[51, 52]

Currently, HS treatment includes medications, colectomy and AVR. However, medications and intestinal resection are not effective and the recurrence rate is high. Segmental bowel resection, electrocoagulation, or laser photocoagulation are often ineffective.^[53, 54]

AVR could alleviate coagulopathy and anemia of HS patients.^[55] This is related to the possible mechanism of HS, where blood flow through the narrowed AS can lead to the destruction of HMW vWF, which plays an important role in the coagulation process. In the current study, 43 patients underwent SAVR, and 12 underwent TAVR; 43 underwent biological valve replacement and 14 underwent mechanical valve replacement. The patients with mechanical valve received anticoagulation after surgery, and no bleeding was found during postoperative follow-up. After mechanical

Figure 1. Flow diagram of study screening.
HS: Heyde syndrome.

Table 1. Characteristics of the included cases

Studies	Country	Age (yrs)	Sex	AS		AVA (cm ²)	Gastrointes- tinal bleed- ing site ^a	Hb (g/L)	vWF	Operation	Valve	Follow-up	
				PPG (mm Hg)	MPG (mm Hg)							Prognosis	Time
Tsuchiya <i>et al.</i> ^[5]	Japan	83	F	NR	81	0.5		NR	↓	TAVR	M	Well	2 years
Garcia <i>et al.</i> ^[6]	Brazil	64	M	60	36.5	0.6	Jejunum	NR	NR	SAVR	M	Well	
Mahboobi <i>et al.</i> ^[7]	USA	56	F	NR	NR	NR		NR	NR	SAVR	NR	Well	
Mirna <i>et al.</i> ^[8]	Australia	73	NR	98	53	0.8	Ileum	65	NR	TAVR	M	Well	
Alshuwaykh <i>et al.</i> ^[9]	USA	56	F	NR	35	0.8		67	↓	TAVR	M	Well	6 months
Iijima <i>et al.</i> ^[10]	Japan	77	M	88	NR	0.52		55	↓	SAVR	B	Well	20 months
Achamallah <i>et al.</i> ^[11]	USA	73	F	NR	NR	0.75		NR	NR	SAVR	B	Well	
Achamallah <i>et al.</i> ^[11]	USA	84	F	NR	NR	0.94		NR	NR	TAVR	M	Well	
Akutagawa <i>et al.</i> ^[12]	Japan	67	F	NR	NR	0.8	Duodenum, jejunum	61	↓	SAVR	B	Well	6 months
Balbo <i>et al.</i> ^[13]	Brazil	81	M	35	22	NR	colon	64	NR	TAVR	M	Well	
Giovannini <i>et al.</i> ^[14]	Italy	70	M	162	103	0.4	Jejunum	NR	NR	SAVR	B	Die	10 months
Iquille <i>et al.</i> ^[15]	France	77	M	77	48	0.8		50	NR	SAVR	B	Well	1 year
Shibamoto <i>et al.</i> ^[16]	Japan	87	M	NR	59	NR	Ileum	87	↓	SAVR	B	Well	20 months
Uchida <i>et al.</i> ^[17]	Japan	77	F	100	NR	0.6	Colon	61	↓	SAVR	B	Die	31 days
Huang <i>et al.</i> ^[18]	China	76	F	NR	NR	NR		65	↓	TAVR	M	Well	
Hudzik <i>et al.</i> ^[19]	Canada	82	M	NR	37	0.7		95	↓	TAVR	M	Well	12 months
Wang <i>et al.</i> ^[20]	China	84	F	89	57	NR		69	NR	TAVR	M	Well	4 months
Perry <i>et al.</i> ^[21]	USA	57	F	NR	NR	NR		NR	NR	SAVR	NR	Well	
Benton <i>et al.</i> ^[22]	USA	77	M	132	62	NR	Colon	NR	↓	TAVR	M	Well	10 months
İlkeli <i>et al.</i> ^[23]	Turkey	46	F	144	86	NR	Duodenum, ileum, cecum, colon	90	NR	SAVR	B	Well	1 year
Kapila <i>et al.</i> ^[24]	USA	70	M	NR	40	0.7	Duodenum, jejunum	NR	↓	SAVR	NR	Well	2 months
Dos Santos <i>et al.</i> ^[25]	Brazil	71	M	NR	NR	NR	Cecum	NR	NR	SAVR	B	Well	
Maor <i>et al.</i> ^[26]	Israel	76	F	68	NR	0.8	Duodenum	70	NR	SAVR	B	Well	42 months
Negi <i>et al.</i> ^[27]	USA	75	F	NR	NR	NR	Jejunum	NR	NR	TAVR	M	Well	
Saad <i>et al.</i> ^[28]	UK	76	F	72	45	0.9		50	NR	SAVR	B	Well	4 months
Godino <i>et al.</i> ^[29]	Italy	83	F	104	58	0.7	Duodenum	61	NR	TAVR	M	Well	6 months
Gül <i>et al.</i> ^[30]	Turkey	75	F	NR	52	0.6	Ileum, cecum	NR	NR	TAVR	M	Well	2 months
Michot <i>et al.</i> ^[31]	France	79	M	140	75	0.6	Duodenum, ileum, cecum, colon	54	NR	SAVR	B	Well	9 years
Van <i>et al.</i> ^[32]	Belgium	76	F	46	30	1		89	NR	SAVR	B	NR	
Abi-Akar <i>et al.</i> ^[33]	Lebanon	68	M	97	66	0.7	Duodenum, jejunum, colon	NR	NR	SAVR	B	Well	9 months

Continued Table 1. Characteristics of the included cases

Studies	Country	Age (yrs)	Sex	AS		AVA (cm ²)	Gastrointes- tinal bleed- ing site ^a	Hb (g/L)	vWF	Operation	Valve	Follow-up	
				PPG (mm Hg)	MPG (mm Hg)							Prognosis	Time
D'Souza <i>et al.</i> ^[34]	Canada	79	M	NR	84	0.94	Jejunum	NR	↓	SAVR	B	Well	
Takahashi <i>et al.</i> ^[35]	Japan	82	F	139	NR	0.6	Colon	78	↓	SAVR	B	Well	4 years
Undas <i>et al.</i> ^[36]	Poland	74	F	98	68	0.6	Jejunum	NR	Normal	SAVR	B	Well	7 months
Morishima <i>et al.</i> ^[37]	Japan	78	F	127	NR	NR	Jejunum, colon	93	↓	SAVR	B	Well	
Singh <i>et al.</i> ^[38]	USA	71	F	100	65	NR		NR	NR	SAVR	B	Well	1 year
Soran <i>et al.</i> ^[39]	UK	80	M	100	NR	NR	Ileum	94	NR	SAVR	B	Well	1 year
Warkentin <i>et al.</i> ^[40]	Canada	65	F	NR	NR	0.67	Jejunum	NR	↓	SAVR	B	Well	10 years
Warkentin <i>et al.</i> ^[40]	Canada	70	F	NR	NR	0.5	Colon	NR	↓	SAVR	B	Well	10 years
Anderson <i>et al.</i> ^[41]	Australia	62	F	NR	NR	0.5	Cecum	NR	NR	SAVR	NR	Well	6 months
Anderson <i>et al.</i> ^[41]	Australia	77	F	NR	NR	0.5	Ileum	NR	NR	SAVR	NR	Well	6 months
Olearchyk <i>et al.</i> ^[42]	USA	69	M	>50	NR	0.6	Cecum	NR	NR	SAVR	M	Well	
Apostolakis <i>et al.</i> ^[43]	Germany	67	NR	NR	NR	NR	Colon	117	NR	SAVR	B	Not Well*	
Apostolakis <i>et al.</i> ^[43]	Germany	68	F	NR	NR	NR	Colon	60	NR	SAVR	B	Not Well [#]	
Casson <i>et al.</i> ^[44]	Canada	68	F	NR	60	NR		100	NR	SAVR	NR	Well	
Baciewicz <i>et al.</i> ^[45]	USA	56	M	32	NR	0.8	Ileum, cecum, colon	43	NR	SAVR	B	Well	8 months
Cappell <i>et al.</i> ^[46]	USA	66	F	120	NR	0.6	Duodenum	NR	NR	SAVR	B	Well	15 months
Cappell <i>et al.</i> ^[46]	USA	69	F	110	NR	0.5	Colon	NR	NR	SAVR	B	Well	18 months
Scheffer <i>et al.</i> ^[47]	USA	78	F	NR	65	NR		87	NR	SAVR	B	Well	9 months
Marcus <i>et al.</i> ^[48]	RSA	66	F	NR	NR	NR		NR	NR	SAVR	B	Well	42 months
Marcus <i>et al.</i> ^[48]	RSA	78	M	NR	NR	NR		NR	NR	SAVR	B	Well	42 months
Marcus <i>et al.</i> ^[48]	RSA	72	F	NR	NR	NR		NR	NR	SAVR	NR	Well	12 months
Marcus <i>et al.</i> ^[48]	RSA	66	F	NR	NR	NR		NR	NR	SAVR	NR	Well	12 months
Danilewitz <i>et al.</i> ^[49]	USA	66	F	115	NR	NR		NR	NR	SAVR	B	Well	12 months
Love <i>et al.</i> ^[50]	NR	72	F	100	NR	NR		NR	NR	SAVR	B	Well	
Love <i>et al.</i> ^[50]	NR	61	F	160	NR	NR		NR	NR	SAVR	B	Well	10 months

AS: aortic stenosis; PPG: peak pressure gradient; MPG: mean pressure gradient; AVA: aortic valve area; Hb: hemoglobin; vWF: von Willebrand factor; F: female; M: male; NR: not reported; SAVR: surgical aortic valve replacement; TAVR: transcatheter aortic valve replacement; B: bioprothetic valve; M: mechanical valve; *GI bleeding recurred one year after SAVR; [#]Melena occurred on postoperative day 13 after SAVR; ^aGastrointestinal bleeding was present in all the cases, but it was not easy to be detected, so we only described the bleeding at a specific site on the above table.

AVR, HS patients might experience bleeding due to warfarin administration. Therefore, some have suggested the use of the bioprosthetic valve instead to minimize anticoagulation-related bleeding post-AVR.

For critically-ill patients at high surgical risk and those unsuitable for thoracotomy, minimally invasive valve replacement therapy or TAVR is an alternative option.^[13] In the current study, 12 patients underwent TAVR and no bleeding reoccurred during follow-up. In view of the active gastrointestinal bleeding, clopidogrel monotherapy was chosen over dual antiplatelet therapy post-interventionally.^[8]

Uchida *et al.*^[17] reported a case of death following AVR in a patient who had preoperative acute left heart failure with exertional dyspnea, with a peak aortic pressure gradient of 100 mm Hg. The woman died of life-threatening acute subdural hematoma 31 days postoperatively. However, recurrent gastrointestinal bleeding was not observed during the postoperative period. Giovannini *et al.*^[14] reported a patient with repeated bleeding and massive blood transfusion and multiple comorbidities. With all conservative treatments failing to stop bleeding, the patient underwent SAVR and had no bleeding within 10 months. However, 10 months after the operation, the patient underwent intestinal resection due to intestinal bleeding and then anastomotic leakage, and finally died of multiple organ failure. A case of HS after SAVR due to a mismatched biological prosthesis was reported by Iquille *et al.*^[15] The patient recovered after reoperation and various physiological parameters returned to normal.

In recent years, with TAVR technology constantly progressing, choosing SAVR or TAVR has become a major problem that plagues clinicians. In a retrospective study by Desai *et al.*,^[56] there is no statistically difference in mortality and hospital cost between HS

patients undergoing SAVR and TAVR, but TAVR queues showed lower stroke incidence and blood transfusion incidence. Compared with SAVR, TAVR has fewer complications.

Due to the risk of bleeding, perioperative management of such patients is very challenging. However, in the above case reports, there is almost no mention of how to conduct perioperative management, so the summary is as follows.

Before surgery, if gastroenteroscopy suggested a clear bleeding point, endoscopic treatment should be performed prior to AVR. In most cases, it is difficult to assess the actual bleeding risk of the patient, and it is even more difficult to quantify the bleeding situation. Therefore, how to evaluate vWF activity to guide preventive treatment is particularly important. There are currently two kinds of tests. The first is a gel electrophoresis test, which can quantitatively analyze the level of HMW vWF multimer, but it is not suitable for the perioperative period because it is expensive and time-consuming. Another test is to use a platelet function analyzer to analyze platelet function, which can indirectly indicate vWF abnormal and is suitable for the entire perioperative period. Once vWF is reduced, desmopressin can be given for treatment, and vWF/FVIII concentrate, platelets, plasma and cryoprecipitate can be given if necessary.^[57] Hemostatic medications available for application during the perioperative period are shown in **Table 2**.

Due to the defects of HS and the influence of cardiopulmonary bypass, the risk of bleeding of HS patients during operation is greater. Tranexamic acid is recommended to use to enhance fibrin formation. In addition, be gentle when performing laryngoscopy, tracheal intubation, indwelling gastric tube, and placing transesophageal echocardiogram probe.

Table 2. Hemostatic interventions for HS

Products	Mechanism of action	Others
Desmopressin	Selective V2 agonist; Releases endogenous VWF, FVIII	Expensive; 0.3 µg/kg, iv, over 30 minutes
Alphanate/Fanhdii/Humate P	VWF/FVIII concentrates	Expensive; Increased chance of thromboembolism; Need to check the vWF and FVIII level
Vonvendi	Recombinant VWF	Expensive; Less thromboembolic risk as does not increase FVIII level;
Tranexamic acid	Antifibrinolytic agents	Can be administered along with desmopressin and vWF/FVIII concentrate
Platelet	Platelet activation and aggregation	Transmission of infection
Fresh frozen plasma	Contains VWF	Transmission of infection
Cryoprecipitate	Contains VWF	Transmission of infection

VWF: von Willebrand factor; FVIII: factor VIII.

After operation, as stenosis of aortic valve is corrected, vWF factor recovers quickly, risk of gastrointestinal bleeding is reduced.^[57]

HS is a rare syndrome unknown to many physicians. Therefore, clinicians should be mindful of HS as a potential differential diagnosis in patients with a systolic murmur and presenting with repeated episodes of gastrointestinal bleeding. Moreover, capsule endoscopy can be useful for diagnosing this disease. Although currently HS is not an indication for AS treatment, reports suggested that, the outcome of severe HS patients with AVR was good, and intestinal resection delayed to determine if bleeding stops after AVR.

Appendix Table 1 is available on the *CMSJ* website at doi: 10.24920/003881.

Declaration of Conflicts of Interest

The authors declare that they have no competing interests for this manuscript.

REFERENCES

- Heyde EC. Gastrointestinal bleeding in aortic stenosis (letter). *N Engl J Med* 1958; 259:196.
- Supino PG, Borer JS, Preibisz J, et al. The epidemiology of valvular heart disease: a growing public health problem. *Heart Fail Clin* 2006; 2(4):379-93. doi: 10.1016/j.hfc.2006.09.010.
- Vincentelli A, Susen S, Le Tourneau T, et al. Acquired von Willebrand syndrome in aortic stenosis. *N Engl J Med* 2003; 349(4):343-9. doi: 10.1056/NEJMoa022831.
- Pate GE, Mulligan A. An epidemiological study of Heyde's syndrome: an association between aortic stenosis and gastrointestinal bleeding. *J Heart Valve Dis* 2004; 13(5):713-6.
- Tsuchiya S, Matsumoto Y, Doman T, et al. Disappearance of angiodysplasia following transcatheter aortic valve implantation in a patient with Heyde's Syndrome: a case report and review of the literature. *J Atheroscler Thromb* 2020; 27(3):271-7. doi: 10.5551/jat.49239.
- Garcia LR, Garzesi AM, Tripoli G, et al. Heyde Syndrome treated by conventional aortic valve replacement. *Braz J Cardiovasc Surg* 2019; 34(5):630-2. doi: 10.21470/1678-9741-2019-0004.
- Mahboobi SK. Heyde's syndrome and postoperative bleeding after aortic valve replacement—is there a role for prophylactic desmopressin? *J Clin Anesth* 2019 ; 56:142. doi: 10.1016/j.jclinane.2019.02.002.
- Mirna M, Lichtenauer M, Theurl T, et al. Transcatheter aortic valve implantation in a patient with suspected hereditary von Willebrand disease and severe gastrointestinal bleeding—a case report. *Scott Med J* 2019; 64(4):142-7. doi: 10.1177/0036933019862155.
- Alshuwaykh O, Krier MJ. A case of Heyde Syndrome with resolution of gastrointestinal bleeding two weeks after aortic valve replacement. *Am J Case Rep* 2018; 19:924-6. doi: 10.12659/AJCR.911298.
- Iijima M, Itoh N, Murase R, et al. A surgical case of aortic stenosis with recurrent gastrointestinal bleeding: Heyde syndrome. *Int J Surg Case Rep* 2018; 53:281-4. doi: 10.1016/j.ijscr.2018.10.078.
- Achamallah N, Hardwick A. Tavr and heyde: the role of trans-catheter aortic valve replacement in aortic stenosis associated gastrointestinal bleeding. *J General Intern Med* 2017; 32:2.
- Akutagawa T, Shindo T, Yamanouchi K, et al. Persistent gastrointestinal angiodysplasia in Heyde's Syndrome after aortic valve replacement. *Intern Med* 2017; 56(18):2431-3. doi: 10.2169/internalmedicine.8603-16.
- Balbo CP, Seabra LP, Galoro VG, et al. Heyde's Syndrome and transcatheter aortic valve implantation. *Arq Bras Cardiol* 2017; 108(4):378-80. doi: 10.5935/abc.20160193.
- Giovannini I, Chiarla C, Murazio M, et al. An extreme case of Heyde syndrome. *Dig Surg* 2006; 23(5-6):387-8. doi: 10.1159/000098457.
- Iquille J, Nader J, Colpart E, et al. Unusual gastrointestinal bleeding after sutureless aortic valve replacement: a word of caution. *Ann Thorac Surg* 2017; 103(3):e225-e6. doi: 10.1016/j.athoracsurg.2016.08.073.
- Shibamoto A, Kawaratani H, Kubo T, et al. Aortic valve replacement for the management of Heyde Syndrome: a case report. *J Nippon Med Sch* 2017; 84(4):193-7. doi: 10.1272/jnms.84.193.
- Uchida T, Hamasaki A, Ohba E, et al. Life-threatening subdural hematoma after aortic valve replacement in a patient with Heyde syndrome: a case report. *J Cardiothorac Surg* 2017; 12(1):65. doi: 10.1186/s13019-017-0629-x.
- Huang DJ, Kong LJ, Wu Z. Transcatheter aortic valve implantation in the treatment of Hyde syndrome: a case report. *Chin J Intern Med* 2016; 55(9):721-2. doi: 10.3760/cma.j.issn.0578-1426.2016.09.013.

19. Hudzik B, Wilczek K, Gasior M. Heyde syndrome: gastrointestinal bleeding and aortic stenosis. *CMAJ* 2016; 188(2):135-8. doi: 10.1503/cmaj.150194.
20. Wang B, Wang Y, Su ML. Case report of a patient with Heyde syndrome treated by transcatheter aortic valve implantation. *Zhonghua Xin Xue Guan Bing Za Zhi* 2016; 44(3):269-70. doi:10.3760/cma.j.issn.0253-3758.2016.03.016.
21. Perry PA, Atkins BZ, Amsterdam EA. Gastrointestinal bleeding, aortic stenosis, and the hiding culprit. *Am J Med* 2015; 128(8):e5-6. doi: 10.1016/j.amjmed.2015.02.015.
22. Benton SM Jr, Kumar A, Crenshaw M, et al. Effect of transcatheter aortic valve implantation on the Heyde's syndrome. *Am J Cardiol* 2014; 114(6):953-4. doi: 10.1016/j.amjcard.2014.06.025.
23. İlkeli E, Albeyoğlu Ş, Çiloğlu U, et al. Heyde's syndrome. *Asian Cardiovasc Thorac Ann* 2014; 22(5):592-4. doi: 10.1177/0218492313478433.
24. Kapila A, Chhabra L, Khanna A. Valvular aortic stenosis causing angiodysplasia and acquired von Willebrand's disease: Heyde's syndrome. *BMJ Case Rep* 2014; 2014:bcr2013201890. doi: 10.1136/bcr-2013-201890.
25. Dos Santos VM, Dos Santos LA, Modesto AA, et al. Heyde syndrome in a 71-year-old man who underwent chest radiotherapy at young age. *An Sist Sanit Navar* 2013; 36(2):339-45. doi: 10.4321/s1137-66272013000200020.
26. Maor NR. Heyde syndrome: resolution of anemia after aortic valve surgery. *Isr Med Assoc J* 2013; 15(7):387-9.
27. Negi SI, Xu R, Anand A. Medical image. A murmur and a bleed: the Heyde syndrome. *N Z Med J* 2013; 126(1371):95-7.
28. Saad RA, Lwaleed BA, Kazmi RS. Gastrointestinal bleeding and aortic stenosis (Heyde syndrome): the role of aortic valve replacement. *J Card Surg* 2013; 28(4):414-6. doi: 10.1111/jocs.12131.
29. Godino C, Pavon AG, Mangieri A, et al. Aortic valvuloplasty as bridging for TAVI in high-risk patients with Heyde's Syndrome: a case report. *Case Rep Med* 2012; 2012:946764. doi: 10.1155/2012/946764.
30. Gül M, Sürçit Ö, Özal E, et al. Treatment of aortic valve stenosis and gastrointestinal bleeding by transcatheter aortic valve implantation in Heyde syndrome. *Anadolu Kardiyol Derg* 2012; 12(8):691-3. doi: 10.5152/akd.2012.224.
31. Michot JM, Treton X, Brink C, et al. Severe gastrointestinal angiodysplasia in context of Heyde's syndrome durably cured after aortic valve replacement. *Presse Med* 2012; 41(7-8):763-6. doi: 10.1016/j.lpm.2011.10.012.
32. van Hul IJ, Schwagten V, Vrelust I. Association of aortic valve stenosis and recurrent gastrointestinal bleeding. *Eur J Emerg Med* 2012; 19(2):122-4. doi: 10.1097/MEJ.0b013e32834991a0.
33. Abi-Akar R, El-Rassi I, Karam N, et al. Treatment of Heyde's Syndrome by aortic valve replacement. *Curr Cardiol Rev* 2011; 7(1):47-9. doi: 10.2174/157340311795677699.
34. D'Souza PM, Blostein MD. Diagnosis of Heyde's syndrome by abnormal closure times despite normal von Willebrand's activity. *Blood Coagul Fibrinolysis* 2011; 22(7):622-3. doi: 10.1097/MBC.0b013e32834a04d3.
35. Takahashi N, Tanabe K, Yoshitomi H, et al. Successful endoscopic clipping for bleeding from colonic angiodysplasia in a case of Heyde syndrome. *Med Sci Monit* 2010; 16(9):CS107-9.
36. Undas A, Windyga J, Bykowska K, et al. Heyde's syndrome without a decrease in large von Willebrand factor multimers: a case of intestinal bleedings reversed by valve replacement in a patient with aortic stenosis. *Thromb Haemost* 2009; 101(4):773-4.
37. Morishima A, Marui A, Shimamoto T, et al. Successful aortic valve replacement for Heyde syndrome with confirmed hematologic recovery. *Ann Thorac Surg* 2007; 83(1):287-8. doi: 10.1016/j.athoracsurg.2006.05.082.
38. Singh P, Scoyni R, Pooran N, et al. Aortic valve replacement: a last resort for aortic stenosis-associated refractory GI bleeding. *Gastrointest Endosc* 2002; 56(1):139-41. doi: 10.1067/mge.2002.125108.
39. Soran H, Lewis M, Whorwell PJ. Bleeding angiodysplasia: should we concentrate more on the aortic valve than on the bowel?. *Int J Clin Pract* 2002; 56(2):155-6.
40. Warkentin TE, Moore JC, Morgan DG. Gastrointestinal angiodysplasia and aortic stenosis. *N Engl J Med* 2002; 347(11):858-9. doi: 10.1056/NEJM200209123471122.
41. Anderson RP, McGrath K, Street A. Reversal of aortic stenosis, bleeding gastrointestinal angiodysplasia, and von Willebrand syndrome by aortic valve replacement. *Lancet* 1996; 347(9002):689-90. doi: 10.1016/s0140-6736(96)91240-4.
42. Olearchyk AS. Heyde's syndrome. *J Thorac Cardiovasc Surg* 1992; 103(4):823-4.
43. Apostolakis E, Doering C, Kantartzis M, et al. Calcific

- aortic-valve stenosis and angiodysplasia of the colon: Heyde's syndrome—report of two cases. *Thorac Cardiovasc Surg* 1990; 38(6):374-6. doi: 10.1055/s-2007-1014055.
44. Casson AG, McKenzie NN. Heyde's syndrome. *Chest* 1988; 94(4):891-2. doi: 10.1378/chest.94.4.891.
45. Baciewicz FA Jr, Davis JT. Heyde's syndrome: failure of a mechanical prosthesis and the possibility of a coagulation defect. *Ann Thorac Surg* 1987; 44(5):554-5. doi: 10.1016/s0003-4975(10)62126-1.
46. Cappell MS, Lebowitz O. Cessation of recurrent bleeding from gastrointestinal angiodysplasias after aortic valve replacement. *Ann Intern Med* 1986; 105(1):54-7. doi: 10.7326/0003-4819-105-1-54.
47. Scheffer SM, Leatherman LL. Resolution of Heyde's syndrome of aortic stenosis and gastrointestinal bleeding after aortic valve replacement. *Ann Thorac Surg* 1986; 42(4):477-80. doi: 10.1016/s0003-4975(10)60563-2.
48. Marcus RH, Kinsley RH. Aortic stenosis and unexplained gastro-intestinal haemorrhage. Case reports. *S Afr Med J* 1982; 61(15):559-61.
49. Danilewitz M, McKibbin J, Derman D. Cessation of gastrointestinal bleeding after valve replacement of aortic stenosis. *Am Heart J* 1981; 101(5):686. doi: 10.1016/0002-8703(81)90245-3.
50. Love JW, Jahnke EJ, Zacharias D, et al. Calcific aortic stenosis and gastrointestinal bleeding. *N Engl J Med* 1980; 302(17):968. doi: 10.1056/NEJM198004243021716.
51. Pate GE, Chandavimol M, Naiman SC, et al. Heyde's syndrome: a review. *J Heart Valve Dis* 2004; 13(5):701-12.
52. Massyn MW, Khan SA. Heyde syndrome: a common diagnosis in older patients with severe aortic stenosis. *Age Ageing* 2009; 38(3):267-70; discussion 251. doi: 10.1093/ageing/afp019.
53. Gelfand ML, Cohen T, Ackert JJ, et al. Gastrointestinal bleeding in aortic stenosis. *Am J Gastroenterol* 1979; 71(1):30-8.
54. Ogano M, Iwasaki YK, Takano H, et al. Successful colectomy for the treatment of repetitive bleeding from colonic angiodysplasia in a patient with Heyde syndrome. *Intern Med* 2006; 45(6):355-8. doi: 10.2169/internalmedicine.45.1526.
55. Warkentin TE, Moore JC, Morgan DG. Aortic stenosis and bleeding gastrointestinal angiodysplasia: is acquired von Willebrand's disease the link? *Lancet* 1992; 340(8810):35-7. doi: 10.1016/0140-6736(92)92434-h.
56. Desai R, Parekh T, Singh S, et al. Alarming increasing trends in hospitalizations and mortality with Heyde's Syndrome: a nationwide inpatient perspective (2007 to 2014). *Am J Cardiol* 2019; 123(7):1149-55. doi: 10.1016/j.amjcard.2018.12.043.
57. Mondal S, Hollander KN, Ibekwe SO, et al. Heyde Syndrome—pathophysiology and perioperative implications. *J Cardiothorac Vasc Anesth* 2020; S1053-0770(20)31098-3. doi: 10.1053/j.jvca.2020.10.003.

中文摘要

海德综合征与主动脉瓣置换术：已发表文献回顾

李慧利¹，林陪双²，姚允泰^{3*}

¹ 河北医科大学第一医院麻醉科，石家庄 050000

² 福建医科大学附属泉州第一医院麻醉科，泉州 362000

³ 中国医学科学院北京协和医学院，国家心血管病中心阜外医院麻醉科，北京 100037

摘要

目的 探讨采用主动脉瓣置换术治疗的海德综合征患者的临床特点及结局。

方法 使用检索词“Heyde syndrome”、“gastrointestinal bleeding”、“aortic stenosis”和“surgery”，检索PubMed、Embase、Ovid、WANFANG、VIP和CNKI等数据库，提取纳入病例报道中的患者生物学特征、主动脉瓣狭窄程度、胃肠道出血部位、手术治疗和预后等临床资料。

结果 纳入的46篇病例报道共包括55例患者，年龄46~87岁。主动脉瓣轻度狭窄1例，中度狭窄1例，重度狭窄42例（余未提及）。胃肠道出血部位分别为结肠（ $n=8$ ）、空肠（ $n=6$ ）、回肠（ $n=4$ ）、盲肠（ $n=3$ ）、十二指肠（ $n=3$ ）和多个部位（ $n=8$ ）；未发现特定出血部位的患者为23例。术前血红蛋白水平为43~117 g/L。16例患者接受高分子量 von Willebrand 因子检测，结果显示1例正常，其余病例均下降。在55例患者中，43例接受经胸主动脉瓣置换术，12例接受经导管主动脉瓣置换术；14例完成机械瓣置换，33例完成生物瓣置换。5例患者出现并发症：1例术后出现瓣周漏和消化道出血，并行二次主动脉瓣置换术；2例患者反复出现胃肠道出血。2例分别因急性硬脑膜下血肿和多器官衰竭死亡。在随访期内，其余患者状况良好。

结论 海德综合征是一种以主动脉瓣狭窄和消化道出血为特征的综合征。主动脉瓣置换术是海德综合征的一种有效治疗方法。

关键词：海德综合征；胃肠道出血；主动脉瓣狭窄