REVIEW

Biomarkers with Potential Predictive Value for Cardiotoxicity in Anticancer Treatments

Wei Yang, Mei Zhang*

The Key Laboratory of Cardiovascular Remodeling and Function Research of Chinese Ministry of Education and Chinese Ministry of Health, The State and Shandong Province Joint Key Laboratory of Translational Cardiovascular Medicine, Department of Cardiology, Qilu Hospital, Cheeloo College of Medicine, Shandong University, Jinan 250012, China

Key words: biomarkers; cardiotoxicity; anticancer treatments; review

Abstract Rapid development of anticancer treatments in recent years has greatly improved prognosis of cancer patients. However, with extension of survival time of cancer patients, various short-term and long-term side effects brought about by anticancer treatments, especially cardiotoxicity, have become increasingly prominent. Nonetheless, at present, there is few diagnostic methods with extremely high sensitivity and specificity to detect and accurately predict whether patients with anticancer treatment will experience cardiovascular complications. Inflammation, fibrosis and oxidative stress are considered to be important mechanisms involved in cardiotoxicity anticancer treatments. The cardiovascular biomarkers having the ability to predict and detect cardiovascular dysfunction earlier than clinical symptoms as well as left ventricular ejection fraction monitored by echocardiography, are of great value to timely treatment adjustment and prognosis evaluation. Cardiac troponin T/I and brain natriuretic peptide/ N-terminal prohormone of brain natriuretic peptide have been routinely used in clinical practice to monitor cardiotoxicity, and some new biomarkers such as soluble suppression of tumorigenicity-2, myeloperoxidase, growth differentiation factor-15, galectin-3, endothelin-1, have potential in this area. In the future, larger-scale experimental studies are needed to provide sufficient evidences, and how to detect them quickly and at low cost is also a problem to be dealt with.

MALIGNANT tumors and heart diseases have become the most common diseases and leading causes of death in the world, and more than 70% of patients

die from them.^[1] Nowadays, anticancer treatments have progressed great, and developed into a comprehensive treatment regimen, which has significantly improved prognosis of cancer patients. However, with survival rate improvement of patients with malignant tumors, side effects of anticancer treatments gradually emerged, increasing the morbidity and mortality of complications.^[2, 3] The cardiovascular disease caused

Received June 7, 2020. Published online October 8, 2021.

*Corresponding author Tel: 86-18560086629, E-mail: daixh@vip.sina.com

by cardiotoxicity is the primary complication of anticancer treatments.^[4] In the 2016 practical guideline published by European Society of Cardiology (ESC), cardiovascular diseases induced by anticancer therapy area classified into 9 categories: cardiac insufficiency and heart failure, coronary artery disease, valvular disease, arrhythmia (especially prolonged QT interval), hypertension, pulmonary hypertension, thromboembolic disease, peripheral vascular disease and stroke, and other cardiovascular diseases such as pericardial disease.^[5]

However, at present, there is no diagnostic method with high sensitivity and specificity to early detect and accurately predict cardiovascular complications in patients with malignant tumors. When abnormal changes in left ventricular ejection fraction (LVEF) detected by echocardiography or related clinical symptoms occur, cardiac function of patients has usually deteriorated to such an extent that they need to stop anticancer treatments and be treated with cardioprotective drugs (such as angiotensin-converting enzyme inhibitors, β -blockers, etc.) to alleviate the symptoms, which has an adverse effect on prognosis of patients to a great extent. Subtle cardiovascular dysfunction caused by anticancer therapy can be detected by myocardial strain and related cardiovascular biomarkers earlier than by LVEF, which is helpful to delay progress of cardiac complications and improve prognosis by adjusting anticancer therapy or adding cardioprotective therapy.^[6] Some studies have revealed that anthracyclines can induce cardiotoxicity on cardiovascular system through oxidative stress, inflammatory response and vascular remodeling.^[7-9] Moreover, studies have shown that multi-target tyrosine kinase growth and angiogenesis inhibitors exert cardiotoxic effects by inhibiting vascular endothelial growth factor and vascular endothelial growth factor receptor tyrosine kinase to damage vascular endothelial cells and destroy cardiac contractility and vasodilator function.^[10, 11] Therefore, through identifying potential biomarkers that can predict cardiotoxicity, cardiotoxicity maybe expected to be detected early, thus guiding treatment and improving prognosis of patients with anticancer treatment.

BIOMARKERS WITH POTENTIAL PREDICTIVE VALUE FOR CARDIOTOXICITY

Cardiac troponin

Cardiac troponin (cTn) is a protein complex that exists

only in cardiomyocytes and is related to myocardial contraction. In the case of myocardial infarction, acute and chronic myocardial injury, cTn is released into the blood and the concentration in serum is increased. Therefore, detection of serum cTn has become the first choice for diagnosis of acute coronary syndrome. Detecting high-sensitivity cTn (hs-cTn) can shorten diagnosis time of acute myocardial infarction, improve accuracy, and screen patients with asymptomatic acute and chronic myocardial injury early.^[12, 13] Moreover, elevated serum troponin concentration caused by cardiomyocyte necrosis induced by direct or indirect stimulation, can increase left ventricular filling pressure, excessive oxidative stress or catecholamine, proinflammatory factors and other stimulating factors, so cTn will be elevated in many other cardiovascular diseases, such as myocarditis, heart failure, cardiomyopathy (including Takotsubo cardiomyopathy), aortic valve stenosis, tachyarrhythmia, pulmonary embolism, aortic dissection, and so on (**Table 1**).^[14]

In the field of cardio-oncology, cardiotoxicity caused by cancer itself and anticancer treatments can also trigger abnormal cTn expression through cell damage, oxidative stress, fibrosis and other pathways. A recent Meta analysis has shown that anticancer treatments can elevate serum cTn level, and in patients receiving anticancer treatments, increased cTn is related to systolic dysfunction, suggesting its value for predicting left ventricular dysfunction, which deserves to be studied further. In addition, reduction of cardiotoxicity through β -blockers and angiotensin-converting enzyme inhibitors during anticancer treatments is associated with decrease of serum cTn concentration.^[15] Another study on cardiotoxicity of anthracyclines in patients with breast cancer has also shown that anthracyclines can increase concentration of hs-cTnT, and increase in hs-cTnT levels at the end of anthracycline treatment may indicate subsequent cardiotoxicity.^[16] It means that cTn has a potential value in risk stratification for patients receiving anticancer therapy and prognosis assessment for these patients. In the previous International CardioOncology Society-One Trial (ICOS-ONE), researchers found that through prospective follow-up, chemotherapy regimens containing anthracyclines increased troponin level even with low cumulative doses of anthracyclines in adult patients with low cardiovascular risk.^[17] Because anthracycline drugs have potent clinical therapeutic effects on different solid cancers (especially breast cancer) and

lymphomas, anthracycline drugs are the main components of anticancer chemotherapy.^[18] It also reflects the value of cTn in the early detection of cardiotoxicity caused by anticancer therapy. In recent years, troponin has been widely used in clinical practice due to convenience of detection by only taking a tube of blood, low cost and good diagnostic value. Routinely monitoring troponin for early detection of cardiotoxicity in patients receiving anticancer therapy has become a tendency. The latest European Society of Medical Oncology (ESMO) consensus also recommends that hs-cTnI/T baseline measurements be considered for high-risk patients (previously significant cardiovascular diseases) and those patients receiving high doses of cardiotoxic chemotherapy (such as anthracyclines), at the same time, and regularly monitoring hs-cTnI/T for patients receiving anticancer treatments.^[19]

Brain natriuretic peptide and N-terminal prohormone of brain natriuretic peptide

Brain natriuretic peptide (BNP) and N-terminal prohormone of brain natriuretic peptide (NT-proBNP) are hormones secreted by cardiomyocytes under stimulating by factors such as cardiomyocyte stretch, neurohormone activation and myocardial hypoxia, among which cardiomyocyte stretch plays a leading role in the process. BNP/NT-proBNP acts on distant tissues and has effects of diuresis, vasodilation, and regulating body's water-sodium balance.^[20] In contrast, NT-proBNP has a longer half-life period than BNP.^[20] Both of them are widely used in clinic as markers of heart failure. Yang *et al.*^[21] have confirmed that lower NT-proBNP levels indicate worse endothelial function and can be used as a sign of cardiac decompensation. It also has value for early detection of cardiotoxicity of anticancer treatments (**Table 1**).

However, BNP and NT-proBNP as biomarkers to early detecting cardiotoxicity caused by anticancer treatments are still controversial issues. A study has shown that there is a consistent temporal correlation between NT-proBNP and cardiotoxicity during long-term follow-up,^[16] which indicating that NT-proBNP has a certain significance in predicting cardiac toxicity. But a previous study on longitudinal changes of cardiac markers in breast cancer patients receiving anticancer treatments got an opposite result. In the study, even if the enrolled patients were followed up for a maximum of 15 months, no a clear correlation between NT-proBNP and cardiotoxicity have been found.^[22]

Therefore, the predictive and diagnostic value of BNP and NT-proBNP for cardiotoxicity induced by anticancer therapy still needs further studies to verify. Nevertheless, considering their advantages of low cost and convenience, BNP/NT-proBNP can be promoted as a biomarker of myocardial toxicity in patients with anticancer treatments, so as to monitor cardiac function. Guidelines from the Canadian Cardiovascular Society (CCS) recommend routine BNP monitoring for early detection of cardiac dysfunction in patients receiving cardiotoxic drugs.^[23] The consensus of the European Society of Medical Oncology (ESMO) also recommends routine monitoring of BNP and NT-proBNP for high-risk patients (previously significant cardiovascular diseases) and ones receiving high-dose cardiotoxic chemotherapy (such as anthracyclines) to detect cardiotoxic effects, so as to treat them in time and delay their progress.^[19]

Soluble suppression of tumorigenicity-2

Suppression of tumorigenicity-2 (ST2) is a member of interleukin-1 (IL-1) receptor family. There are four subtypes of ST2, which are soluble ST2 (sST2), transmembrane ST2 (ST2L), and two splicing variants (ST2V and ST2LV). Among them, sST2 and ST2L are mainly related to cardiovascular system. IL-33 is considered to be a specific functional ligand of ST2. IL-33/ST2 is a kind of fibroblast-cardiomyocyte intercellular signaling system, which participates in many processes such as antigen/allergen reaction, autoimmunity, organ fibrosis and heart injury, and plays a cardioprotective role in reducing fibrosis, reducing myocardial hypertrophy and myocardial compensation and so on. In this signaling system, sST2 acts as a decoy receptor that competes with ST2L to bind IL-33 to block its cardioprotective effect, and is associated with poor prognosis of patients with heart diseases (**Table 1**).^[24]

As a biomarker of inflammation, fibrosis, and myocardial stress, more and more studies have been conducted on diagnostic and prognostic value of sST2 in heart failure and myocardial infarction. Recently, Morin *et al.*^[25] have conducted an one-year follow-up study on 196 patients with LVEF $\leq 35\%$ and primary prophylaxis implanted cardioverter defibrillator, and monitoring results from baseline and every 3 months showed that sST2 has significantly independent predictive value for all-cause mortality. In addition, periodically repeat monitoring does not provide a more accurate risk stratification than baseline measurements alone. In management of chronic diseases, expensive

medical costs make many patients give up halfway. This finding illustrates that monitoring cardiac insufficiency with sTS2 may save medical costs. Another large sample analysis of 5301 patients with chronic heart failure found that sST2 was less affected by age than NT-proBNP or hs-TnT, helping to achieve more accurate risk stratification and prognosis management of heart failure.^[26] Another study of patients with myocardial infarction also showed that sST2 has predictive value for cardiovascular adverse events.^[27] These studies bring more consideration on potential value of sST2 in diagnosis of cardiac dysfunction of patients with anticancer treatments and predicting prognosis of them.

A previous study showed that sST2 levels increased in breast cancer patients during and after treatment with anthracyclines, however, the study did not specify whether the elevated levels of sST2 were caused by breast cancer itself or cardiotoxicity caused by anthracyclines.^[28] Another study followed up breast cancer patients who received radiotherapy and found that there was a negative correlation between sST2 levels and cardiac systolic function, and this study suggested that sST2 could also predict cardiotoxicity induced by radiotherapy.^[29] However, sample size of these studies is small, and accuracy of the results needs to be further verified. Clinical studies with larger sample size, longer follow-up time and better trial design are needed in the future to verify the efficacy of SSTZ monitoring. Moreover, rapid and routine detection of sST2 in clinic is also a problem to be solved.

Myeloperoxidase

Myeloperoxidase (MPO), a member of the heme peroxidase superfamily, is secreted by activated polymorphonuclear leukocytes, and catalyzes the transformation of active substances including hypochlorous acid and other reactive oxidant species, which posttranslationally modify the target proteins and contribute to pathogenesis of systemic inflammatory diseases. It is considered as a biomarker of oxidative stress and inflammatory response.^[30-33] Previous study has shown that MPO levels in blood are negatively correlated with the prognosis of a variety of cardiovascular diseases and positively correlated with risk of cardiovascular death.^[34] MPO contributes to risk stratification in a variety of cardiovascular diseases, such as acute coronary syndrome,^[35] atherosclerosis, and heart failure^[30] (**Table 1**).

Since oxidative stress is considered to be the core mechanism of cardiotoxicity induced by anthra-

cyclines,^[36] attention has been extensively paid to the potential value of MPO in monitoring cardiotoxicity in anticancer treatments. A study of breast cancer patients treated with anthracyclines and trastuzumab found an association between elevated MPO levels and cardiotoxicity.^[22] A recent study of breast cancer patients who were also treated with anthracycline and trastuzumab also found that patients with higher baseline levels of MPO were more likely to develop cardiovascular adverse events.^[16] These findings indicate that MPO may be a predictor of cardiotoxicity, and baseline measurement of MPO may contribute to early risk stratification. However, for HER-2-negative breast cancer patients who received non-anthracycline (bevacizumab combined with docetaxel and cyclophosphamide), MPO did not show a significant correlation with cardio-toxicity even if cardiotoxicity was found during follow-up.^[37] Therefore, more in-depth studies are needed on application of MPO in patients with cardiotoxicity caused by non-anthracycline therapy, even for patients treated with anthracycline. However, as a new biomarker in the field of cardio-oncology, MPO has a broad research prospect.

Growth differentiation factor-15

Growth differentiation factor-15 (GDF-15) is a member of transforming growth factor- β superfamily,^[38] which is secreted by cardiac myocytes stimulated by mechanical stretch, ischemia, injury, inflammation, and oxidative/nitric stress.^[39] As a biomarker of myocardial remodeling and pro-inflammation, GDF-15 is closely related to cardiovascular diseases.^[40] GDF-15 is independently associated with cardiovascular adverse events in patients of chronic heart failure with reduced ejection fraction (HFrEF).^[41] Higher GDF-15 level indicates poor cardiac systolic function, which is considered to be an independent predictor of heart failure.^[42] In addition, in patients with acute coronary syndrome, GDF-15 levels were positively correlated with risk of cardiovascular adverse events (including bleeding, spontaneous myocardial infarction, stroke) and total mortality.^[43] Subsequent study showed that higher GDF-15 levels predicted a three-fold increased risk of death from heart failure and a one-fold increased risk of sudden cardiac death/arrhythmia after acute coronary syndrome events. GDF-15 is seen as a powerful predictor of all-cause death (**Table 1**).^[44]

In recent years, GDF-15, a biomarker of pro-

inflammation, has also been studied for monitoring cardiac dysfunction associated with anticancer treatments. Tromp *et al.*^[45] found that GDF-15 level was associated with cardiotoxicity induced by anti-breast cancer drugs and had independent predictive value for the event of decreased LVEF. There is the evidence that GDF-15 levels continued to increase during a 15-month follow-up in patients treated with anti-breast cancer drugs (adriamycin, taxane, trastuzumab), supporting that GDF-15 is associated with cardio-toxicity.^[22] However, related study has not found evidence of GDF-15 for early prediction of cardiotoxicity caused by anti-breast cancer drugs.^[46] Considering limitation of sample size and follow-up time, and influence of potential confounding factors, the existing results may not exactly reveal what role GDF-15 plays in cardiotoxicity. Therefore, GDF-15 cannot be clinically used.

Galectin-3

Galectin-3 (Gal-3) is a kind of stromal cell polysaccharide binding protein, which is mainly secreted by macrophages under inflammatory reaction, stress, tissue injury and other stimulating factors. Gal-3 is considered as a biomarker reflecting myocardial fibrosis by promoting activation of fibroblasts and participating in myocardial fibrosis and remodeling.^[47] Gal-3 has important predictive value for heart failure, cardiovascular diseases, and all-cause mortality, and has diagnostic and prognostic value for both ejection fraction preserved heart failure (HFpEF) and HFrEF.^[48-50] Another study has shown that Gal-3 contributes to risk stratification of cardiovascular diseases and directs treatment, especially in older adults (**Table 1**).^[51]

Animal studies have found that doxorubicin can induce expression of Gal-3 through stress-activated protein kinase/c-Jun-NH2 terminal kinase signal, promoting myocardial fibrosis and resulting in cardiac toxicity.^[52] Gal-3 is considered to be a novel biomarker for monitoring cardiotoxicity induced by anticancer treatments. Previous studies found that Gal-3 levels were abnormal in some breast cancer patients who received TAC (docetaxel, doxorubicin and cyclophosphamide) chemotherapy regimen one year after the end of chemotherapy.^[53] However, due to limitation of sample size and follow time, the results do not show that Gal-3 can predict cardiotoxicity in its early stage. In addition, radiotherapy can also cause cardiotoxicity through mechanism of fibrosis.^[54] Abnormally high levels of Gal-3 can be induced by radiotherapy.^[55] There-

fore, for patients with malignant tumors who receive radiotherapy (especially in the mediastinum and left chest), more studies are needed to explore the link between Gal-3 and cardiotoxicity caused by radiotherapy in the future.

Endothelin-1

Endothelin-1 (ET-1) is a potent vasoconstrictor peptide synthesized and secreted mainly by cardiomyocytes,^[56, 57] which can cause vasoconstriction by acting on vascular smooth muscle cells and exert positive muscle strength effect by acting on cardiomyocytes (**Table 1**).^[58] Anticancer drug doxorubicin can promote production of reactive oxygen species, destroy function of endothelium-based cardiomyocytes, cause secretion of ET-1, thus leading cardiotoxicity.^[59] ET-1 has two functional receptors: A receptor and B receptor. Some researchers have found that blocking ET-1 receptor can reduce cardiotoxicity of doxorubicin and play a cardioprotective role.^[60] Researchers should take a step further by verifying whether ET-1 can diagnose and detect cardiotoxicity caused by anticancer treatments earlier.

CLINICAL SETTING OF BIOMARKERS FOR CARDIOTOXICITY

In clinical practice, baseline measurements and regular routine monitoring of cTnT/I and BNP/NT-proBNP are recommended for high-risk patients (previously significant cardiovascular disease) and those treated with cardiotoxic drugs on the basis of current guidelines and consensus.^[19, 23] Periodic routine monitoring of GDF-15 can be performed at the same time with consent of cancer patients if possible, which contribute to assess severity of cardiotoxicity in patients receiving anticancer drugs, and then adjust anticancer treatment regimens or add cardioprotective therapy in time. Monitoring of sST2 and Gal-3 is beneficial for risk stratification and guiding treatment in elderly patients. In addition, due to the widespread use of anthracyclines in the treatment of different solid tumors and lymphomas,^[18] periodic monitoring of MPO and ET-1 to assess cardiac and vascular toxicity has a certain value in evaluating prognosis of tumor patients treated with anthracyclines. However, more in-depth studies and larger clinical trials are needed to explore their mechanism and specific applicable population before these biomarkers (sST2, MPO, GDF-15, Gal-3, ET-1, etc.) are widely used in clinical practice.

In conclusion, rapid development of comprehensive anticancer treatments not only prolongs survival time of patients, but also brings cardiotoxicity and a variety of cardiovascular complications. Biomarkers may be of great value for early detection, treatment

guidance and prognosis evaluation of cardiotoxicity. Although immature, cTnT/I and BNP/NT-proBNP have been recommended for routine clinical use in monitoring cardiotoxicity.^[19, 23] Some novel biomarkers (sST2, MPO, GDF-15, Gal-3, ET-1, etc.) have been considered

Table 1. Features of the biomarkers for cardiotoxicity in anticancer treatments

Biomarkers	Physiopathologic mechanism	Clinical value for cardiotoxicity	Correlations with CVD	Advantages
cTn	Involving in acute and chronic myocardial injury and cell death	Prediction of left ventricular dysfunction; Evaluation of efficacy of cardiac protective therapy; Evaluation of prognosis	Myocarditis, HF, cardiomyopathy (including Takotsubo cardiomyopathy), aortic valve stenosis, tachyarrhythmia, pulmonary embolism, aortic dissection	Convenience of detection; Low cost Superior diagnostic value
BNP/NT-proBNP	Diuresis, vasodilation and regulation of human water-sodium balance; Related to endothelial function	Consistent temporal correlation between NT-proBNP and cardiotoxicity	Acute and chronic cardiac dysfunction and even HF	Convenience of detection; Low cost
sST2	Act as a decoy receptor that competes with ST2L to bind IL-33 to block its cardioprotective effect (reducing fibrosis, reducing myocardial hypertrophy, myocardial compensation and so on)	Prediction of cardiac systolic dysfunction; Risk stratification; Evaluation of prognosis	Acute and chronic cardiac dysfunction and even HF, MI	Independent predictive value for all-cause mortality; Less affected by age; Early risk stratification may be achieved at the baseline level
MPO	Play an anti-inflammatory role after catalytic reaction; Reflective of oxidative stress and inflammatory response	Early risk stratification of cardiotoxicity induced by anthracyclines	CVD such as ACS, atherosclerosis, and HF	Early risk stratification of cardiotoxicity induced by anthracyclines may be achieved at the baseline level
GDF-15	Involving in myocardial remodeling and pro-inflammation	Prediction of left ventricular dysfunction; Correlation between GDF-15 level and severity of cardiotoxicity	HFrEF, ACS, bleeding, spontaneous MI, stroke, sudden cardiac death, arrhythmia	Independent predictive value for all-cause mortality; Related to severity of cardiotoxicity
Gal-3	Involving in myocardial fibrosis	Risk stratification; Evaluation of prognosis	HFpEF and HFrEF	Independent predictive value for all-cause mortality; Contribute to risk stratification and guide treatment in the elderly
ET-1	Vasoconstriction; Positive muscle strength effect; Involving in endothelial function	Blocking ET-1 receptor can reduce cardiotoxicity of doxorubicin and play a cardioprotective role	Incident CVD	Related to cardiotoxicity induced by anthracyclines

CVD: cardiovascular disease; cTn: cardiac troponin; HF: heart failure; BNP: brain natriuretic peptide; NT-proBNP: N-terminal prohormone of brain natriuretic peptide; sST2: soluble suppression of tumorigenicity-2; ST2L: transmembrane suppression of tumorigenicity ST2; IL-33: interleukin-33; MI: myocardial infarction; MPO: myeloperoxidase; ACS: acute coronary syndrome; GDF-15: growth differentiation factor-15; HFrEF: heart failure with reduced ejection fraction; Gal-3: galectin-3; HFpEF: heart failure with preserved ejection fraction; ET-1: endothelin-1.

to be concerned with cardiotoxicity in preceding studies. More studies are needed to explore their application values in cardiotoxicity management of patients with malignant tumors receiving anticancer treatments.

Conflicts of Interest Statement

All authors have no competing interests to declare.

REFERENCES

1. Jones DS, Podolsky SH, Greene JA. The burden of disease and the changing task of medicine. *N Engl J Med* 2012; 366(25):2333-8. doi: 10.1056/NEJMp1113569.
2. Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer* 2013; 49(6):1374-403. doi: 10.1016/j.ejca.2012.12.027.
3. Siegel R, DeSantis C, Virgo K, et al. Cancer treatment and survivorship statistics, 2012. *CA Cancer J Clin* 2012; 62(4):220-41. doi: 10.3322/caac.21149.
4. Ewer MS, Ewer SM. Cardiotoxicity of anticancer treatments. *Nat Rev Cardiol* 2015; 12(9):547-58. doi: 10.1038/nrcardio.2015.65.
5. Zamorano JL, Lancellotti P, Rodriguez MD, et al. 2016 ESC Position Paper on cancer treatments and cardiovascular toxicity developed under the auspices of the ESC Committee for Practice Guidelines: The Task Force for cancer treatments and cardiovascular toxicity of the European Society of Cardiology (ESC). *Eur Heart J* 2016; 37(36):2768-801. doi: 10.1093/eurheartj/ehw211.
6. Sawaya H, Sebag IA, Plana JC, et al. Early detection and prediction of cardiotoxicity in chemotherapy-treated patients. *Am J Cardiol* 2011; 107(9):1375-80. doi: 10.1016/j.amjcard.2011.01.006.
7. Ky B, Vejpongsa P, Yeh ET, et al. Emerging paradigms in cardiomyopathies associated with cancer therapies. *Circ Res* 2013; 113(6):754-64. doi: 10.1161/CIRCRESAHA.113.300218.
8. Octavia Y, Tocchetti CG, Gabrielson KL, et al. Doxorubicin-induced cardiomyopathy: from molecular mechanisms to therapeutic strategies. *J Mol Cell Cardiol* 2012; 52(6):1213-25. doi: 10.1016/j.yjmcc.2012.03.006.
9. Xu MF, Tang PL, Qian ZM, et al. Effects by doxorubicin on the myocardium are mediated by oxygen free radicals. *Life Sci* 2001; 68(8):889-901. doi: 10.1016/s0024-3205(00)00990-5.
10. Lenneman CG, Sawyer DB. Cardio-oncology: an update on cardiotoxicity of cancer-related treatment. *Circ Res* 2016; 118(6):1008-20. doi: 10.1161/CIRCRESAHA.115.303633.
11. Ylä-Herttuala S, Rissanen TT, Vajanto I, et al. Vascular endothelial growth factors: biology and current status of clinical applications in cardiovascular medicine. *J Am Coll Cardiol* 2007; 49(10):1015-26. doi: 10.1016/j.jacc.2006.09.053.
12. Morrow DA. The Fourth Universal Definition of Myocardial Infarction and the Emerging Importance of Myocardial Injury. *Circulation* 2020; 141(3):172-5. doi: 10.1161/CIRCULATIONAHA.119.044125.
13. Shah KS, Yang EH, Maisel AS, et al. The role of biomarkers in detection of cardio-toxicity. *Curr Oncol Rep* 2017; 19(6):42. doi: 10.1007/s11912-017-0602-9.
14. Eggers KM, Lindahl B. Application of cardiac troponin in cardiovascular diseases other than acute coronary syndrome. *Clin Chem* 2017; 63(1):223-35. doi: 10.1373/clinchem.2016.261495.
15. Michel L, Mincu RI, Mahabadi AA, et al. Troponins and brain natriuretic peptides for the prediction of cardiotoxicity in cancer patients: a meta-analysis. *Eur J Heart Fail* 2020; 22(2):350-61. doi: 10.1002/ehf.1631.
16. Demissei BG, Hubbard RA, Zhang L, et al. Changes in cardiovascular biomarkers with breast cancer therapy and associations with cardiac dysfunction. *J Am Heart Assoc* 2020; 9(2):e014708. doi: 10.1161/JAHA.119.014708.
17. Cardinale D, Ciceri F, Latini R, et al. Anthracycline-induced cardiotoxicity: a multicenter randomised trial comparing two strategies for guiding prevention with enalapril: The International CardioOncology Society-One trial. *Eur J Cancer* 2018; 94:126-37. doi: 10.1016/j.ejca.2018.02.005.
18. Curigliano G, Cardinale D, Dent S, et al. Cardiotoxicity of anticancer treatments: epidemiology, detection, and management. *CA Cancer J Clin* 2016; 66(4):309-25. doi: 10.3322/caac.21341.
19. Curigliano G, Lenihan D, Fradley M, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. *Ann Oncol* 2020; 31(2):171-90. doi: 10.1016/j.annonc.2019.10.023.
20. Hall C. NT-ProBNP: the mechanism behind the marker. *J Card Fail* 2005; 11(5 Suppl):S81-3. doi: 10.1016/j.cardfail.2005.04.019.
21. Yang S, Huang S, Daniels LB, et al. NT-proBNP, race

- and endothelial function in the Multi-Ethnic Study of Atherosclerosis. *Heart* 2019; 105(20):1590-6. doi: 10.1136/heartjnl-2019-314707.
22. Putt M, Hahn VS, Januzzi JL, et al. Longitudinal changes in multiple biomarkers are associated with cardiotoxicity in breast cancer patients treated with doxorubicin, taxanes, and trastuzumab. *Clin Chem* 2015; 61(9):1164-72. doi: 10.1373/clinchem.2015.241232.
23. Virani SA, Dent S, Brezden-Masley C, et al. Canadian Cardiovascular Society Guidelines for evaluation and management of cardiovascular complications of cancer therapy. *Can J Cardiol* 2016; 32(7):831-41. doi: 10.1016/j.cjca.2016.02.078.
24. Kakkar R, Lee RT. The IL-33/ST2 pathway: therapeutic target and novel biomarker. *Nat Rev Drug Discov* 2008; 7(10):827-40. doi: 10.1038/nrd2660.
25. Morin DP, Chong-Yik R, Thihalolipavan S, et al. Utility of serial measurement of biomarkers of cardiovascular stress and inflammation in systolic dysfunction. *Eurpace* 2020; 22(7):1044-53. doi: 10.1093/europace/ euaa075.
26. Aimo A, Januzzi JL, Vergaro G, et al. Circulating levels and prognostic value of soluble ST2 in heart failure are less influenced by age than N-terminal pro-B-type natriuretic peptide and high-sensitivity troponin T. *Eur J Heart Fail* 2020; 22(11):2078-88. doi: 10.1002/ ejhf.1701.
27. Zagidullin N, Motloch LJ, Gareeva D, et al. Combining novel biomarkers for risk stratification of two-year cardiovascular mortality in patients with ST-elevation myocardial infarction. *J Clin Med* 2020; 9(2):550. doi: 10.3390/jcm9020550.
28. Frères P, Bouznad N, Servais L, et al. Variations of circulating cardiac biomarkers during and after anthracycline-containing chemotherapy in breast cancer patients. *BMC Cancer* 2018; 18(1):102. doi: 10.1186/s12885-018-4015-4.
29. Aula H, Skyttä T, Tuohinen S, et al. ST2 levels increased and were associated with changes in left ventricular systolic function during a three-year follow-up after adjuvant radiotherapy for breast cancer. *Breast* 2020; 49:183-6. doi: 10.1016/j.breast.2019.12.001.
30. Tang WH, Tong W, Troughton RW, et al. Prognostic value and echocardiographic determinants of plasma myeloperoxidase levels in chronic heart failure. *J Am Coll Cardiol* 2007; 49(24):2364-70. doi: 10.1016/j.jacc.2007.02.053.
31. Odobasic D, Kitching AR, Holdsworth SR. Neutrophil-mediated regulation of innate and adaptive immunity: the role of myeloperoxidase. *J Immunol Res* 2016; 2016:2349817. doi: 10.1155/2016/2349817.
32. Abdo AI, Rayner BS, van Reyk DM, et al. Low-density lipoprotein modified by myeloperoxidase oxidants induces endothelial dysfunction. *Redox Biol* 2017; 13:623-32. doi: 10.1016/j.redox.2017.08.004.
33. Love DT, Barrett TJ, White MY, et al. Cellular targets of the myeloperoxidase-derived oxidant hypothiocyanous acid (HOSCN) and its role in the inhibition of glycolysis in macrophages. *Free Radic Biol Med* 2016; 94:88-98. doi: 10.1016/j.freeradbiomed.2016.02.016.
34. Ndrepepa G. Myeloperoxidase—a bridge linking inflammation and oxidative stress with cardiovascular disease. *Clin Chim Acta* 2019; 493:36-51. doi: 10.1016/j.cca.2019.02.022.
35. Baldus S, Heeschen C, Meinertz T, et al. Myeloperoxidase serum levels predict risk in patients with acute coronary syndromes. *Circulation* 2003; 108(12):1440-5. doi: 10.1161/01.CIR.0000090690.67322.51.
36. Finkelman BS, Putt M, Wang T, et al. Arginine-nitric oxide metabolites and cardiac dysfunction in patients with breast cancer. *J Am Coll Cardiol* 2017; 70(2):152-62. doi: 10.1016/j.jacc.2017.05.019.
37. Gullo G, J EA, Canonici A, et al. Pilot study of bevacizumab in combination with docetaxel and cyclophosphamide as adjuvant treatment for patients with early stage HER-2 negative breast cancer, including analysis of candidate circulating markers of cardiac toxicity: ICORG 08-10 trial. *Ther Adv Med Oncol* 2019; 11:1758835919864236. doi: 10.1177/1758835919864236.
38. Bootcov MR, Bauskin AR, Valenzuela SM, et al. MIC-1, a novel macrophage inhibitory cytokine, is a divergent member of the TGF-beta superfamily. *Proc Natl Acad Sci USA* 1997; 94(21):11514-9. doi: 10.1073/pnas.94.21.11514.
39. Xu XY, Nie Y, Wang FF, et al. Growth differentiation factor (GDF)-15 blocks norepinephrine-induced myocardial hypertrophy via a novel pathway involving inhibition of epidermal growth factor receptor transactivation. *J Biol Chem* 2014; 289(14):10084-94. doi: 10.1074/jbc.M113.516278.
40. Wollert KC, Kempf T, Wallentin L. Growth differentiation factor 15 as a biomarker in cardiovascular disease. *Clin Chem* 2017; 63(1):140-51. doi: 10.1373/clinchem.2016.255174.
41. Bouabdallaoui N, Claggett B, Zile MR, et al. Growth differentiation factor-15 is not modified by sacubitril/valsartan and is an independent marker of risk in

- patients with heart failure and reduced ejection fraction: the PARADIGM-HF trial. *Eur J Heart Fail* 2018; 20(12):1701-9. doi: 10.1002/ejhf.1301.
42. Stenemo M, Nowak C, Byberg L, et al. Circulating proteins as predictors of incident heart failure in the elderly. *Eur J Heart Fail* 2018; 20(1):55-62. doi: 10.1002/ejhf.980.
 43. Hagström E, James SK, Bertilsson M, et al. Growth differentiation factor-15 level predicts major bleeding and cardiovascular events in patients with acute coronary syndromes: results from the PLATO study. *Eur Heart J* 2016; 37(16):1325-33. doi: 10.1093/eurheartj/ehv491.
 44. Lindholm D, James SK, Gabrys K, et al. Association of multiple biomarkers with risk of all-cause and cause-specific mortality after acute coronary syndromes: a secondary analysis of the PLATO biomarker study. *JAMA Cardiol* 2018; 3(12):1160-6. doi: 10.1001/jamacardio.2018.3811.
 45. Tromp J, Boerman LM, Sama IE, et al. Long-term survivors of early breast cancer treated with chemotherapy are characterized by a pro-inflammatory biomarker profile compared to matched controls. *Eur J Heart Fail* 2020; 22(7):1239-46. doi: 10.1002/ejhf.1758.
 46. Ky B, Putt M, Sawaya H, et al. Early increases in multiple biomarkers predict subsequent cardiotoxicity in patients with breast cancer treated with doxorubicin, taxanes, and trastuzumab. *J Am Coll Cardiol* 2014; 63(8):809-16. doi: 10.1016/j.jacc.2013.10.061.
 47. Suthahar N, Meijers WC, HHW S, et al. From inflammation to fibrosis-molecular and cellular mechanisms of myocardial tissue remodelling and perspectives on differential treatment opportunities. *Curr Heart Fail Rep* 2017; 14(4):235-50. doi: 10.1007/s11897-017-0343-y.
 48. Zile MR, O'Meara E, Claggett B, et al. Effects of sacubitril/valsartan on biomarkers of extracellular matrix regulation in patients with HFrEF. *J Am Coll Cardiol* 2019; 73(7):795-806. doi: 10.1016/j.jacc.2018.11.042.
 49. Ghorbani A, Bhamhani V, Christenson RH, et al. Longitudinal change in galectin-3 and incident cardiovascular outcomes. *J Am Coll Cardiol* 2018; 72(25):3246-54. doi: 10.1016/j.jacc.2018.09.076.
 50. Kosmala W, Przewlocka-Kosmala M, Rojek A, et al. Comparison of the diastolic stress test with a combined resting echocardiography and biomarker approach to patients with exertional dyspnea: diagnostic and prognostic implications. *JACC Cardiovasc Imaging* 2019; 12(5):771-80. doi: 10.1016/j.jcmg.2017.10.008.
 51. Mortensen MB, Fuster V, Muntendam P, et al. Negative risk markers for cardiovascular events in the elderly. *J Am Coll Cardiol* 2019; 74(1):1-11. doi: 10.1016/j.jacc.2019.04.049.
 52. Tanaka R, Umemura M, Narikawa M, et al. Reactive fibrosis precedes doxorubicin-induced heart failure through sterile inflammation. *ESC Heart Fail* 2020; 7(2):588-603. doi: 10.1002/ehf2.12616.
 53. van Boxtel W, Bulten BF, Mavinkurve-Groothuis AM, et al. New biomarkers for early detection of cardiotoxicity after treatment with docetaxel, doxorubicin and cyclophosphamide. *Biomarkers* 2015; 20(2):143-8. doi: 10.3109/1354750X.2015.1040839.
 54. Groarke JD, Nguyen PL, Nohria A, et al. Cardiovascular complications of radiation therapy for thoracic malignancies: the role for non-invasive imaging for detection of cardiovascular disease. *Eur Heart J* 2014; 35(10):612-23. doi: 10.1093/eurheartj/ehf114.
 55. Lenihan DJ, Cuculich P. Cardioprotection during therapeutic radiation treatment. *Circ Heart Fail* 2018; 11(8):e005294. doi: 10.1161/CIRCHEARTFAILURE.118.005294.
 56. Yanagisawa M, Kurihara H, Kimura S, et al. A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* 1988; 332(6163):411-5. doi: 10.1038/332411a0.
 57. Suzuki T, Kumazaki T, Mitsui Y. Endothelin-1 is produced and secreted by neonatal rat cardiac myocytes *in vitro*. *Biochem Biophys Res Commun* 1993; 191(3):823-30. doi: 10.1006/bbrc.1993.1291.
 58. Ishikawa T, Yanagisawa M, Kimura S, et al. Positive inotropic action of novel vasoconstrictor peptide endothelin on guinea pig atria. *Am J Physiol* 1988; 255(4 Pt 2):H970-3. doi: 10.1152/ajpheart.1988.255.4.H970.
 59. Luu AZ, Chowdhury B, Al-Omran M, et al. Role of endothelium in doxorubicin-induced cardiomyopathy. *JACC Basic Transl Sci* 2018; 3(6):861-70. doi: 10.1016/j.jacbts.2018.06.005.
 60. Schwebe M, Ameling S, Hammer E, et al. Protective effects of endothelin receptor A and B inhibitors against doxorubicin-induced cardiomyopathy. *Biochem Pharmacol* 2015; 94(2):109-29. doi: 10.1016/j.bcp.2015.01.014.

中文摘要

抗癌治疗中对心脏毒性效应具有潜在预测价值的生物标志物

杨威，张梅

山东大学齐鲁医院心内科，山东 济南 250012 中国

摘要

抗癌治疗在延长肿瘤患者生存期的同时，抗癌治疗引起的各种近期和远期副作用，尤其是心脏毒性显得日益突出。然而，对于恶性肿瘤患者是否会经历心血管并发症，尚无敏感度及特异度极高的早期发现和准确预测手段。炎症、纤维化和氧化应激被认为是抗癌治疗引发心脏毒性的重要机制。相对于临床症状和左室射血分数，相关的心血管生物标志物可能具有更早发现心血管功能紊乱的能力，对于及时肿瘤调整治疗方案以及患者预后评估具有更大价值。临床上，心肌肌钙蛋白 T/I 和脑钠肽 /N- 末端脑钠肽前体已被推荐用于常规心脏毒性监测，一些新的生物标志物，如可溶性肿瘤抑制因子 -2、髓过氧化物酶、生长分化因子 -15、半乳糖凝集素 -3、内皮素 -1 具有潜在的研究价值。在应用于临床之前，需要大样本的试验证实其有效性，同时提高易操作性、降低成本。

关键词：生物标志物；心脏毒性；抗癌治疗；综述