REVIEW

Review of Neuromyelitis Optica Spectrum Disorder with Pain-Depression Comorbidity

Xue Zhang¹, Yan Xu², Lijian Pei^{1*}

¹Department of Anesthesiology, ²Department of Neurology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100730, China

Key words: neuromyelitis optica spectrum disorder; pain; depression; comorbidity

Abstract Neuromyelitis optica spectrum disorder (NMOSD) is an inflammatory disorder of the central nervous system predominantly targeting optic nerves and the spinal cord. The prevalence of the disease is much higher in Asia than in other parts of the world. Pain can be detected in more than 80% of NMOSD patients, with evoked pain mostly being caused by painful tonic muscle spasms and neuropathic pain as the most characteristic types. Depression is often comorbid with pain, and their comorbidity can severely influence quality of life. In recent years, studies have found considerable overlaps between the mechanisms of pain and depression; however, their association remains unclear. This article reviews the epidemiology, mechanism, evaluation and treatment of pain-depression comorbidity in NMOSD patients.

NEUROMYELITIS optica spectrum disorder (NMOSD) is an inflammatory disorder that targets the central nervous system. It is characterized by severe inflammation-mediated demyelination, mainly influencing the optic nerve and spinal cord. The prevalence of NMOSD is 0.52-10/100 000^[1-4] in different studies, with a median patient age of 32 to 41 years. The incidence of NMOSD in the Asian population is 3 times higher than that in other areas of the world.^[3]

Acute bilateral or consecutive optic neuritis leading to severe loss of vision or transverse myelitis

resulting in limb weakness, hypoesthesia or bladder dysfunction are the main characteristics of NMOSD, typically with recurrence. Symptoms of the disease usually appear within a couple of days, and most of the patients can recover to different degrees in the following weeks or months. The detailed etiology of NMOSD remains unclear; however, the important role of aquaporin-4 (AQP4) in the pathophysiology of the disease has been widely acknowledged. AQP4 is the target of the neuromyelitis optica-IgG antibody, and it is distributed in the gray matter of the spinal cord, periaqueductal midbrain and periventricular areas. It is also highly aggregated in the foot processes of astrocytes in the blood-brain barrier. It has been confirmed that neuromyelitis optica-IgG antibody (anti-AQP4 antibody) is a highly specific biomarker in NMOSD patients^[5, 6] that plays a direct role in the pathogenesis of the disease.^[7, 8]

Huang and colleagues had just investigated

Received November 30, 2020. Published online November 8, 2021.

*Corresponding author Tel: 86-18612672127, E-mail: hazelbeijing@vip.163.com

Supported by the Fundamental Research Funds for the Central Universities (3332021015).

NMOSD in China. They found that NMOSD had a great negative impact on the life quality of Chinese patients, including both physical and emotional health. Non-specific oral immunosuppressants were still the most common preventive treatments, and 55.7% of patients were not satisfied with their current treatment option.^[9] Pain and depression always comorbid in NMOSD patients, but were often neglected and remain untreated. This article reviews the epidemiology, mechanism, evaluation and treatment of pain-depression comorbidity in NMOSD patients, and hopefully more attention could be paid to this comorbidity.

NMOSD AND PAIN

Epidemiology

The number of patients in cohort studies in the literature is small (11 to 210 patients) due to the low incidence rate of the disease.^[9-22] The cohort with the largest sample size comes from the United States. Eaneff and colleagues collected 522 patient data points from an online patient community called PatientsLikeMe.^[23] Nevertheless, the data in this study were from voluntary self-reports by patients instead of professional evaluations by physicians, so the reliability of the data was challenged.

Pain is a common symptom of NMOSD. The prevalence of pain in NMOSD patients was as high as 80% in retrospective studies.^[10, 15, 21, 22] Qian and colleagues had proved that current pain was more common in NMOSD patients vs. multiple sclerosis patients (86.2% vs. 40.9%), and more severe on a 10-point scale (5.38 vs. 1.85). They also noticed that medication for multiple sclerosis patients to relieve pain was far more effective than for NMOSD patients.^[21] Two different types of pain were the most common in NMOSD patients: induced pain caused by muscle spasm and neuropathic pain (NeP). Painful tonic spasms were observed in one fourth of NMOSD cases, and were experienced in unilateral or bilateral limbs.^[24] It always emerges during recovery convalescence without accompanying with new neurological deficits or MRI lesions. Muscle spasm could be treated with sodium channel-blocking antiepileptic agents, muscle relaxants, antiparkinsonian drugs and physiotherapy.^[25] NeP constitutes 41.6% to 90.9% of pain in NMOSD patients and mostly appears in the thoracolumbar, leg or back areas.^[10, 12, 22] NeP in NMOSD patients always manifests as pain accompanied by abnormal pain or thermal sensation, such as

allodynia, hyperalgesia, hyperthermesthesia, numbness, and paradoxical heat sensation.

Pain severely influences quality of life in NMOSD patients.^[11, 12, 26] Beekman *et al.*^[27] found that NMOSD had a strong negative impact on physical functioning according to Short Form-36 (SF-36) evaluation. Pain imposed the greatest negative physical impact on overall quality of life, whereas the impact of pain on emotional well-being was controversial.^[9, 27] Studies also found that dissatisfaction with pain treatment options correlated inversely with quality of life.^[27]

Mechanism

Disorder of excitatory glutamate metabolism At the initial stage of disease, the AQP4 antibody combined with AQP4 destroys astrocytes by activating complement or initiating the interaction of effector cells and astrocytes.^[28] Damaged astrocytes recruit T cells, and activated macrophages, microglia and granulocytes create and release a series of cytokines, such as interleukin-1 β (IL-1 β), IL-6, IL-17 and tumor necrosis factor.^[25, 29] IL-1 β and tumor necrosis factor are able to enhance the signal transduction of glutamate, which can further increase the synaptic transmission between nociceptive C fibers in zones I and II of the spinal lamina and neurons. The opening of the N-methyl-D-aspartate (NMDA) receptor on glutamate-controlled calcium channels results in long-term potentiation on the synapses of C fibers, which is an important mechanism of hyperalgesia.^[30, 31]

Damage to AQP4 leads to water balance disorder. At the same time, the expression of other molecules co-expressed with AQP4 in the lipid raft domain, including excitatory amino acid transporter 2 (EAAT2), is also decreased. EAAT2 plays an important role in glutamate metabolism in astrocytes. A decrease in EAAT2 expression leads to a delay in extracellular elimination of glutamate and results in an elevation of extracellular glutamate concentration in areas damaged by NMOSD. Consequently, abnormal neuronal excitation and excitotoxic tissue injury occur.^[32, 33]

α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors are highly expressed in some inhibitory neurons, and these neurons are very sensitive to high glutamate concentrations. Therefore, these neurons are influenced by the excitotoxicity of glutamate much earlier and more strongly than other neurons.^[34] Such damage could destroy the balance between excitatory and inhibitory conduction of

noxious stimulation, decrease the excitation threshold, increase neuroexcitability and neurotoxicity, and lead to pain symptoms such as spontaneous pain, hyperalgesia, and allodynia.^[25]

Disorder of inhibitory γ -aminobutyric acid and glycine metabolism Pain signaling in the spinal cord includes not only bidirectional excitatory conduction but also a strong inhibitory effect to inhibit pain signals and relieve pain. γ -aminobutyric acid (GABA) and glycine are the two most important inhibitory neurotransmitters in the nociceptive stimulus system.^[35] GABA and glycine can prevent the formation of spontaneous pain and allodynia, as well as the regional generalization of hyperalgesia. They can also prevent long-term potentiation formation on the synapses of C fibers to maintain an appropriate reaction for pain stimulation, which plays a significant role in the inhibitory pathway for nociceptive stimuli. After neuronal injury, decreased K^+-Cl^- exchange channel activity and increased $Na^+K^+-Cl^-$ exchange channel activity can both influence chloride exchange to increase intracellular chloride concentrations and result in enhanced apoptosis of inhibitory interneurons in the spinal cord.^[36, 37]

In the central nervous system, glycine is transformed into glutamine and released by astrocytes. Glutamine is then taken up through active transport by GABAergic neurons and transformed back to glycine. Glycine is further transformed into GABA after decarboxylation to take part in the inhibitory process.^[38] When neurons are injured, astrocyte death destroys the stabilization of the glycine-glutamine-GABA pathway; as a result, the inhibitory pathway is damaged.

Neurons in the inhibitory pathway are damaged by anti-AQP4 antibody Inhibitory neurons are mainly distributed in the periaqueductal gray.^[39] The anti-AQP4 antibody can bind to the AQP4 protein in the periaqueductal gray to directly destroy cells in the inhibitory pathway. Therefore, pain inhibition is decreased, and pain symptoms are aggravated.

NMOSD AND PAIN-DEPRESSION COMORBIDITY

Epidemiology

Pain and depression always coexist, and the concurrence of the two diseases is called pain-depression comorbidity. The prevalence of pain-depression comor-

bidity is not clear because of the inconsistency among different studies. Rayner and colleagues reported in their research after the investigation of 1204 patients with chronic pain that 60.8% of patients met the criteria for a diagnosis of depression. Moreover, 33.8% of patients even met the diagnostic criteria for severe depression.^[40] Consequently, depression should be evaluated in patients with chronic pain. The number of studies on NMOSD accompanied by pain-depression comorbidity is limited due to the low prevalence of the disease. In a study that recruited 71 NMOSD patients, 48% had accompanying NeP, and 28% had moderate to severe depression. Researchers also found that the severity of pain and depression were significantly correlated with each other.^[14]

Mechanism

There is no research on the mechanism of pain-depression comorbidity in NMOSD available in the literature. As mentioned above, glutamate and GABA play important roles in the mechanism of pain in NMOSD patients. It is noteworthy that some studies have reviewed whether glutamatergic system hyperfunction^[41, 42] and GABAergic system hypofunction^[43, 44] are both related to depression.

In animal models, signal enhancement of the excitatory glutamate receptor complex was correlated with allodynia and hyperalgesia, including NMDA receptors and AMPA receptors.^[45] Goffer and colleagues indicated that glutamate receptors in the nucleus accumbens are related to chronic pain-related depressive behavior.^[46] They found that the GluA1 subunit of the AMPA receptor in the nucleus accumbens was increased, so more AMPA receptors were calcium permeable. Interestingly, this type of receptor could relieve depressive symptoms without reducing pain. As a result, they speculated that this might be a compensatory mechanism of pain-related depression. Consequently, calcium-permeable AMPA receptors might become a new therapeutic target in the future for the treatment of pain-depression comorbidity.

GABAergic system hypofunction was also related to pain-depression comorbidity; however, research in this area was also limited. Salat and colleagues found in their study that drugs inhibiting the GABA reuptake channel GABA transporter 1 could relieve symptoms of pain and depression in mice. This revealed that the elevation of GABA levels might be effective for the treatment of comorbidities.^[47]

Diagnosis, evaluation and treatment

International consensus diagnostic criteria for NMOSD were published in *Neurology* magazine in 2015 and have been widely used in clinical practice.^[48] The classification and severity of pain-depression comorbidity accompanied by NMOSD are always evaluated by multiple assessment scales. These scales are divided into three types: pain assessment scales, depression assessment scales, and quality of life assessment scales. Pain assessment scales can be used to evaluate the type, symptoms, nature, location, and severity of pain and the degree of relief after treatment. Depression assessment scales are used to evaluate the severity of depression. Physicians use quality of life assessment scales to tell how much the disease influences a patient's quality of life. Here, we summarize the most widely used assessment scales for pain-depression comorbidity evaluation in NMOSD patients, as shown in **Table 1**.

Steroids and immunosuppressive drugs have been widely used for NMOSD treatment. Pulsed steroid therapy such as methylprednisolone at a dose of 1 g given intravenously for 3 to 5 days and plasma exchange for 5 to 7 procedures every other day have been proved

to be effective in treating severe NMOSD relapse.^[49, 50] Intravenous immunoglobulins might also be benefit in NMOSD relapse treatment, but data was limited.^[51] With improved understanding of the mechanism of NMOSD, many new therapeutic strategies have recently been evaluated such as Satralizumab and Tocilizumab interfering with IL-6 receptor signalling, Eculizumab preventing complement activation, and Inebilizumab targeting on AQP-4 antibody producing cells, providing realistic hope that NMOSD patients could be treated more effectively.^[52]

Other than the treatment of the primary disease, NMOSD, analgesia and depression treatment are the most important ways of controlling symptoms of pain and depression. As one of the symptoms that most severely influences patient quality of life, only a few patients with pain receive analgesic treatment, and the effect of medication is far from satisfactory.^[11, 13, 20, 26] It has been proven that, compared with multiple sclerosis patients, NMOSD patients have more accompanying pain symptoms and need higher doses of analgesic drugs but are less sensitive to treatment.^[21] Pain-depression comorbidity influences patient quality of life even more severely but attracts insufficient attention.

Table 1. Assessment scales for pain-depression comorbidity evaluation in NMOSD patients and their characteristics

Name of scales	Characteristics
Pain assessment scales	
NRS	Pain severity evaluation
DN4	Screening NeP
LANSS	Self-reported by patients, screening NeP
BPI	Overall assessment of pain, including severity, location, nature, and relief degree after treatment
NPSI	NeP symptom evaluation
Depression assessment scales	
BDI	Most widely used depression evaluation scale
PHQ-9	High sensitivity and specificity in screening depression in patients with chronic pain
SDS-Zung	Can be finished within 5 min, easy to use
HADS	Used for the evaluation of anxiety and depression
Quality of life assessment scales	
SF-36	Most widely used evaluation of HRQoL
MSQoL	Specific evaluation scale for HRQoL in MS and NMOSD patients
GJCF-NMOSD	Specific evaluation scale for HRQoL in MS and NMOSD patients

NMOSD: neuromyelitis optica spectrum disorder; NRS: Numerical Rating Scale; DN4: Douleur Neuropathique 4 Questions (Neuropathic Pain Four Questions in French); NeP: Neuropathic Pain; LANSS: Leeds Assessment of Neuropathic Symptoms and Signs Pain Scale; BPI: Brief Pain Inventory; NPSI: Neuropathic Pain Symptom Inventory; BDI: Beck's Depression Inventory; PHQ: Patient Health Questionnaire; SDS-Zung: Zung Self-rating Depression Scale; HADS: Hospital Anxiety and Depression Scale; SF: Social Functioning; HRQoL: Health-Related Quality of Life; MSQoL: Multiple Sclerosis Quality of Life; MS: multiple sclerosis; GJSF-NMOSD: Guthy-Jackson Charitable Foundation-Neuromyelitis Optica Spectrum Disorder.

Chavarro and colleagues reported that only 40% of patients with moderate to severe depression received anti-depression treatment among NMOSD patients with comorbidities, and only 50% of the treatment was effective.^[14]

Studies suggested that pain and depression were highly intertwined so that some antidepressants also had analgesic effects such as serotonin and norepinephrine antidepressants.^[53] Lots of drugs were available to control pain-depression comorbid symptoms including anticonvulsants (gabapentin, pregabalin, carbamazepine, oxcarbazepine), antispasticity medications (baclofen, tizanidine, diazepam), opioids (fentanyl, oxycodone), tricyclic antidepressants (amitriptyline), selective serotonin norepinephrine reuptake inhibitor (SNRI, duloxetine), selective serotonin reuptake inhibitors (SSRI, citalopram), and atypical antidepressants (mirtazapine), *et al.*^[14, 21] Non-drug treatment was also important for pain-depression comorbidity treatment in NMOSD patients. Recently, Mealy and his colleagues proved that Scrambler therapy for 10 consecutive weekdays could significantly release pain and depression symptom safely.^[54] The neglect of pain-depression comorbidity in NMOSD patients leads to late treatment for the symptoms. If pain intervention is not provided as early as possible, the effect of drugs will be influenced, especially when the severity of pain and depression reach a certain degree. If preventive intervention is provided for the early treatment of NMOSD, the therapeutic effect will be greatly improved.

CONCLUSIONS

NMOSD patients always suffer from pain, mainly NeP. A large proportion of these patients have pain-depression comorbidity, which influences patients' quality of life to a great extent and deserves attention. Limited proportion of patients with pain-depression comorbidity receive treatment, and the effectiveness of the current strategy is far from satisfactory. Psychotropic and behavioral therapies usually aimed at treating depression might also be useful for releasing pain symptoms, more drugs targeting specific effective site have been evaluated to better control comorbidity symptoms. What's more, non-drug therapies also have caught our attention. More studies on the mechanism and therapeutic method of pain-depression comorbidity in NMOSD are required. Early detection, early in-

tervention, and early treatment should be provided, all of which will be significantly important for improving quality of life in NMOSD patients.

Conflicts of Interest Statement

We have no conflicts of interest to declare.

REFERENCES

1. Cabrera-Gómez JA, Kurtzke JF, González-Quevedo A, et al. An epidemiological study of neuromyelitis optica in Cuba. *J Neurol* 2009; 256(1):35-44. doi: 10.1007/s00415-009-0009-0.
2. Asgari N, Lillevang ST, Skejoe HP, et al. A population-based study of neuromyelitis optica in Caucasians. *Neurology* 2011; 76(18):1589-95. doi: 10.1212/WNL.0b013e3182190f74.
3. Flanagan EP, Cabre P, Weinshenker BG, et al. Epidemiology of aquaporin-4 autoimmunity and neuromyelitis optica spectrum. *Ann Neurol* 2016; 79(5):775-83. doi: 10.1002/ana.24617.
4. Bukhari W, Prain KM, Waters P, et al. Incidence and prevalence of NMOSD in Australia and New Zealand. *J Neurol Neurosurg Psychiatry* 2017; 88(8):632. doi: 10.1136/jnnp-2016-314839.
5. Lennon VA, Wingerchuk DM, Kryzer TJ, et al. A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis. *Lancet* 2004; 364(9451):2106-12. doi: 10.1016/S0140-6736(04)17551-X.
6. Jarius S1, Wildemann B. Aquaporin-4 antibodies (NMO-IgG) as a serological marker of neuromyelitis optica: a critical review of the literature. *Brain Pathol* 2013; 23(6):661-83. doi: 10.1111/bpa.12084.
7. Bennett JL, Lam C, Kalluri SR, et al. Intrathecal pathogenic anti-aquaporin-4 antibodies in early neuromyelitis optica. *Ann Neurol* 2009; 66(5):617-29. doi: 10.1002/ana.21802.
8. Bradl M, Misu T, Takahashi T, et al. Neuromyelitis optica: pathogenicity of patient immunoglobulin *in vivo*. *Ann Neurol* 2009; 66(5):630-43. doi: 10.1002/ana.21837.
9. Huang W, Zhang Bao J, Chang X, et al. Neuromyelitis optica spectrum disorder in China: quality of life and medical care experience. *Mult Scler Relat Disord* 2020; 46:102542. doi: 10.1016/j.msard.2020.102542.
10. Pellkofer HL, Havla J, Hauer D, et al. The major brain endocannabinoid 2-AG controls neuropathic pain and mechanical hyperalgesia in patients with neuromyelitis

- optica. *PLoS One* 2013; 8(8):e71500. doi: 10.1371/journal.pone.0071500.
11. Asseyer S, Schmidt F, Chien C, et al. Pain in AQP4-IgG-positive and MOG-IgG-positive neuromyelitis optica spectrum disorders. *Mult Scler J Exp Transl Clin* 2018; 4(3):2055217318796684. doi: 10.1177/2055217318796684.
 12. Zhao S, Mutch K, Elson L, et al. Neuropathic pain in neuromyelitis optica affects activities of daily living and quality of life. *Mult Scler* 2014; 20(12):1658-61. doi: 10.1177/1352458514522103.
 13. Kong Y, Okoruwa H, Revis J, et al. Pain in patients with transverse myelitis and its relationship to aquaporin 4 antibody status. *J Neurol Sci* 2016; 368:84-8. doi: 10.1016/j.jns.2016.06.041.
 14. Chavarro VS, Mealy MA, Simpson A, et al. Insufficient treatment of severe depression in neuromyelitis optica spectrum disorder. *Neurol Neuroimmunol Neuroinflamm* 2016; 3(6):e286. doi: 10.1212/NXI.0000000000000286.
 15. Brody J, Hellmann MA, Marignier R, et al. Neuromyelitis optica spectrum disorder: disease course and long-term visual outcome. *J Neuropthalmol* 2016; 36(4):356-62. doi: 10.1097/WNO.0000000000000403.
 16. Jarius S, Ruprecht K, Kleiter I, et al. MOG-IgG in NMO and related disorders: a multicenter study of 50 patients. Part 2: epidemiology, clinical presentation, radiological and laboratory features, treatment responses, and long-term outcome. *J Neuroinflammation* 2016; 13(1):280. doi: 10.1186/s12974-016-0718-0.
 17. Masuda H, Mori M, Uzawa A, et al. Validation of the Modified Fatigue Impact Scale and the relationships among fatigue, pain and serum interleukin-6 levels in patients with neuromyelitis optica spectrum disorder. *J Neurol Sci* 2018; 385:64-8. doi: 10.1016/j.jns.2017.11.041.
 18. Mealy MA, Boscoe A, Caro J, et al. Assessment of patients with neuromyelitis optica spectrum disorder using the EQ-5D. *Int J MS Care* 2019; 21(3):129-34. doi: 10.7224/1537-2073.2017-076.
 19. Chanson JB, Zéphir H, Collongues N, et al. Evaluation of health-related quality of life, fatigue and depression in neuromyelitis optica. *Eur J Neurol* 2011; 18(6):836-41. doi: 10.1111/j.1468-1331.2010.03252.x.
 20. Seok JM, Choi M, Cho EB, et al. Fatigue in patients with neuromyelitis optica spectrum disorder and its impact on quality of life. *PLoS One* 2017; 12(5):e0177230. doi: 10.1371/journal.pone.0177230.
 21. Qian P, Lancia S, Alvarez E, et al. Association of neuromyelitis optica with severe and intractable pain. *Arch Neurol* 2012; 69(11):1482-7. doi: 10.1001/archneurol.2012.768.
 22. Kanamori Y, Nakashima I, Takai Y, et al. Pain in neuromyelitis optica and its effect on quality of life: a cross-sectional study. *Neurology* 2011; 77(7):652-8.
 23. Eaneff S, Wang V, Hanger M, et al. Patient perspectives on neuromyelitis optica spectrum disorders: data from the PatientsLikeMe online community. *Mult Scler Relat Disord* 2017; 17:116-22. doi: 10.1212/WNL.0b013e318229e694.
 24. Kim SM, Go MJ, Sung JJ, et al. Painful tonic spasm in neuromyelitis optica: incidence, diagnostic utility, and clinical characteristics. *Arch Neurol* 2012; 69(8):1026-31. doi: 10.1001/archneurol.2012.112.
 25. Bradl M, Kanamori Y, Nakashima I, et al. Pain in neuromyelitis optica-prevalence, pathogenesis and therapy. *Nat Rev Neurol* 2014; 10(9):529-36. doi: 10.1038/nrneurol.2014.129.
 26. Mealy MA, Kozachik SL, Levy M. Review of treatment for central spinal neuropathic pain and its effect on quality of life: implications for neuromyelitis optica spectrum disorder. *Pain Manag Nurs* 2019; 20(6):580-91. doi: 10.1016/j.pmn.2019.03.003.
 27. Beekman J, Keisler A, Pedraza O, et al. Neuromyelitis optica spectrum disorder: patient experience and quality of life. *Neurol Neuroinflamm* 2019; 6(4):e580. doi: 10.1212/NXI.0000000000000580.
 28. Misu T, Höftberger R, Fujihara K, et al. Presence of six different lesion types suggests diverse mechanisms of tissue injury in neuromyelitis optica. *Acta Neuropathol* 2013; 125(6):815-27. doi: 10.1007/s00401-013-1116-7.
 29. Lucchinetti CF, Mandler RN, McGavern D, et al. A role for humoral mechanisms in the pathogenesis of Devic's neuromyelitis optica. *Brain* 2002; 125(Pt 7):1450-61. doi: 10.1093/brain/awf151.
 30. Ikeda H, Stark J, Fischer H, et al. Synaptic amplifier of inflammatory pain in the spinal dorsal horn. *Science* 2006; 312(5780):1659-62. doi: 10.1126/science.1127233.
 31. Drdla R, Gassner M, Gingl E, et al. Induction of synaptic long-term potentiation after opioid withdrawal. *Science* 2009; 325(5937):207-10.
 32. Hinson SR, Roemer SF, Lucchinetti CF, et al. Aquaporin-4-binding autoantibodies in patients with neuromyelitis optica impair glutamate transport by down-regulating EAAT2. *J Exp Med* 2008;

- 205(11):2473-81. doi: 10.1126/science.1171759.
33. Marignier R, Nicolle A, Watrin C, et al. Oligodendrocytes are damaged by neuromyelitis optica immunoglobulin G *via* astrocyte injury. *Brain* 2010; 133(9):2578-91. doi: 10.1093/brain/awq177.
34. Moga D, Hof PR, Vissavajjhala P, et al. Parvalbumin-containing interneurons in rat hippocampus have an AMPA receptor profile suggestive of vulnerability to excitotoxicity. *J Chem Neuroanat* 2002; 23(4):249-53. doi: 10.1016/s0891-0618(02)00012-1.
35. Zeilhofer HU, Wildner H, Yévenes GE. Fast synaptic inhibition in spinal sensory processing and pain control. *Physiol Rev* 2012; 92(1):193-235. doi: 10.1152/physrev.00043.2010.
36. Janssen SP, Truin M, Van Kleef M, et al. Differential GABAergic disinhibition during the development of painful peripheral neuropathy. *Neuroscience* 2011; 184:183-94. doi: 10.1016/j.neuroscience.2011.03.060.
37. Moore KA, Kohno T, Karchewski LA, et al. Partial peripheral nerve injury promotes a selective loss of GABAergic inhibition in the superficial dorsal horn of the spinal cord. *J Neurosci* 2002; 22(15):6724-31. doi: 10.1523/JNEUROSCI.22-15-06724.2002.
38. Albrecht J, Sidoryk-Węgrzynowicz M, Zielińska M, et al. Roles of glutamine in neurotransmission. *Neuron Glia Biol* 2010; 6(4):263-76. doi: 10.1017/S1740925X11000093.
39. Basbaum AI, Bautista DM, Scherrer G, et al. Cellular and molecular mechanisms of pain. *Cell* 2009; 139(2):267-84. doi: 10.1016/j.cell.2009.09.028.
40. Rayner L, Hotopf M, Petkova H, et al. Depression in patients with chronic pain attending a specialised pain treatment centre: prevalence and impact on health care costs. *Pain* 2016; 157(7):1472-9. doi: 10.1097/j.pain.0000000000000542.
41. Mitchell ND, Baker GB. An update on the role of glutamate in the pathophysiology of depression. *Acta Psychiatr Scand* 2010; 122(3):192-210. doi: 10.1111/j.1600-0447.2009.01529.x.
42. Hashimoto K. The role of glutamate on the action of antidepressants. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; 35(7):1558-68. doi: 10.1016/j.pnpbp.2010.06.013.
43. Pehrson AL, Sanchez C. Altered γ -aminobutyric acid neurotransmission in major depressive disorder: a critical review of the supporting evidence and the influence of serotonergic antidepressants. *Drug Des Devel Ther* 2015; 9:603-24. doi: 10.2147/DDDT.S62912.
44. Kaner S, Colquhoun H, Gunduz-Bruce H, et al. Brexanolone (SAGE-547 injection) in post-partum depression: a randomised controlled trial. *Lancet* 2017; 390(10093):480-9. doi: 10.1016/S0140-6736(17)31264-3.
45. Benson C, Mifflin K, Kerr B, et al. Biogenic amines and the amino acids GABA and glutamate: relationships with pain and depression. *Mod Trends Pharmacopsychiatry* 2015; 30:67-79. doi: 10.1159/000435933.
46. Goffer Y, Xu D, Eberle SE, et al. Calcium-permeable AMPA receptors in the nucleus accumbens regulate depression-like behaviors in the chronic neuropathic pain state. *J Neurosci* 2013; 33(48):19034-44. doi: 10.1523/JNEUROSCI.2454-13.2013.
47. Sałat K, Podkowa A, Malikowska N, et al. Novel, highly potent and *in vivo* active inhibitor of GABA transporter subtype 1 with anticonvulsant, anxiolytic, antidepressant and antinociceptive properties. *Neuropharmacology* 2017; 113(Pt A):331-42. doi: 10.1016/j.neuropharm.2016.10.019.
48. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 2015; 85(2):177-89. doi: 10.1212/WNL.0000000000001729.
49. Abboud H, Petrak A, Mealy M, et al. Treatment of acute relapses in neuromyelitis optica: steroids alone *versus* steroids plus plasma exchange. *Mult Scler* 2016; 22(2):185-92. doi: 10.1177/1352458515581438.
50. Watanabe S, Nakashima I, Misu T, et al. Therapeutic efficacy of plasma exchange in NMO-IgG-positive patients with neuromyelitis optica. *Mult Scler* 2007; 13(1):128-32. doi: 10.1177/1352458506071174.
51. Elson L, Panicker J, Mutch K, et al. Role of intravenous immunoglobulin in the treatment of acute relapses of neuromyelitis optica: experience in 10 patients. *Mult Scler* 2014; 20(4):501-4. doi: 10.1177/1352458513495938.
52. Selma J, Selma J. Novel emerging treatments for NMOSD. *Neurol Neurochir Pol* 2019; 53(3):317-26. doi: 10.5603/PJNNS.a2019.0049.
53. Butler S. The triumvirate of co-morbid chronic pain, depression, and cognitive impairment: attacking this "chicken-and-egg" in novel ways. *Scand J Pain* 2017; 15:148-9. doi: 10.1016/j.sjpain.2017.03.004.
54. Mealy MA, Kozachik SL, Cook LJ, et al. Scrambler therapy improves pain in neuromyelitis optica: a randomized controlled trial. *Neurology* 2020; 94(18):e1900-7. doi: 10.1212/WNL.0000000000009370.

中文摘要

视神经脊髓炎谱系疾病合并疼痛抑郁共病综述

张雪¹，徐雁²，裴丽坚^{1*}

¹ 麻醉科，² 神经内科，中国医学科学院北京协和医院，北京，100730，中国

摘要

视神经脊髓炎谱系疾病是一种中枢神经系统炎性疾病，主要累及视神经及脊髓，在亚洲人群中发病率显著高于世界其他地区。80%以上视神经脊髓炎谱系疾病患者合并疼痛症状，其中痛性痉挛及神经病理性疼痛最具特点。抑郁常与疼痛共存，严重影响患者生活质量。近年来，很多研究发现疼痛与抑郁在机制上存在重叠，然而两者之间的联系仍不明确。本文对视神经脊髓炎谱系疾病患者合并疼痛抑郁共病的流行病学特征、发病机制及诊断治疗进行综述。

关键词：视神经脊髓炎谱系疾病；疼痛；抑郁；共病