ORIGINAL ARTICLE

Reliability and Validity of the Chinese Version of the Lymphedema Quality of Life Questionnaire

Zhengyun Liang¹, Xiao Long^{1*}, Elan Yang¹, Yunzhu Li¹, Zhujun Li¹,
Min Xu², Bifen Zhang², Nanze Yu¹, Jiuzuo Huang¹

¹Department of Plastic Surgery and Reconstructive Surgery, Peking Union
Medical College Hospital, Chinese Academy of Medical Sciences &
Peking Union Medical College, Beijing 100005, China

²Shenzhen Yuanwei Medical Equipment Co. Ltd.,
Shenzhen 518116, China

Key words: lymphedema quality of life questionnaire; reliability; validity

Objective To study the reliability and validity of the Chinese version of the Lymphedema Quality of Life Questionnaire (LYMQOL) in lymphedema patients.

Methods LYMQOL was translated into Chinese. The Chinese version of the LYMQOL was distributed with the official Wechat account “Lymphedema Channel” to lymphedema patients who were recruited from October 28th, 2020 to February 23rd, 2021. Patients with upper limb lymphedema and lower limb lymphedema completed the LYMQOL-ARM subscale and the LYMQOL-LEG subscale separately, at enrollment, 1 week later, and 1 month later. Reliability, validity, feasibility, responsiveness and average time required for completing the questionnaire were assessed.

Results A total of 195 patients participated in the study. The Chinese questionnaire showed high reliability with Cronbach's α coefficients of 0.849-0.902 for the LYMQOL-ARM and intraclass correlation coefficient (ICC) of 0.848-0.884 and Cronbach's α coefficients of 0.726-0.902 for the LYMQOL-LEG and ICC of 0.863-0.900. The LYMQOL showed moderate to good correlations with the EQ-5D ($0.4 < r < 0.6$, $P < 0.01$; for the LYMQOL-ARM, $n=95$, for the LYMQOL-LEG, $n=102$). Responsiveness analysis suggested that quality of life of upper limb lymphedema patients decreased after 1 month ($P < 0.05$). The average time of patients to finish the questionnaire was approximately 12 minutes.

Conclusions The Chinese version of the LYMQOL is easy to answer, comprehensive and appropriate in length, and has good reliability and validity. It may be utilized to assist treatment decision-making and track changes in clinical setting or research for lymphedema.

LYMPHEDEMA is a pathological condition involving protein-rich fluid buildup in the interstitial space, leading to chronic swelling of tissues. There are primary and secondary lymphedema depending on the underlying causes to induce inadequate drainage of lymph fluid.^[1] Primary lymphedema is usually a congenital condition of abnormal lymphatic system. Secondary lymphedema is caused by damage to lymphatic system that is mostly due to treatments for breast cancer, gynecological cancer and other malignant tumors, etc. It is reported that about 300 million people worldwide suffer from lymphedema with the incidence of primary and secondary lymphedema being 1:100 000 and 1:1000, respectively.^[2] With distressing symptoms and complications, lymphedema develops into a condition that has a serious impact on daily activity, social ability, psychological state, and quality of life (QoL).

According to document released by Centers for Disease Control and Prevention (CDC), health-related QoL is defined as an individual's or a group's perceived physical and mental health over time, usually including physical functioning, social and psychological wellbeing, etc.^[3] The negative impact of currently incurable lymphedema on QoL has been largely underestimated in the past, but now increasing attention has been focused on patient-reported outcome instruments for lymphedema. A lymphedema-specific self-reported questionnaire, no doubt, can help to fully understand the condition, assisting in clinical decision-making and evaluation of effectiveness as well as cost-utility of interventions.^[4, 5]

The Lymphedema Quality of Life Questionnaire (LYMQOL) is a comprehensive and compact questionnaire specifically designed and developed to evaluate function, body image, symptom, mood and overall QoL of patients with lymphedema.^[6, 7] Its English,^[6] Dutch^[8] and Turkish^[9-11] versions have been validated in terms of psychometric properties. Because in China patient-reported outcome instruments available and suitable for evaluating QoL of lymphedema patients are rather scarce, in this study we translated the LYMQOL into the Chinese version by referring to the cross-cultural adaptation guidelines of questionnaires and further validated psychometric properties of the questionnaires in Chinese lymphedema patients.

MATERIALS AND METHODS

Study design

This study was designed as a two-phase cross-section-

al survey. In Phase 1, we translated the LYMQOL into a Chinese draft and revised it if necessary to develop the final Chinese version of the LYMQOL. In Phase 2, we validated the reliability and validity of the translated version.

This study has been approved by the Ethics Board of Peking Union Medical College Hospital, Chinese Academy of Medical Sciences (IRB number: S-K1544). Patients involved in the study were fully aware of the purpose and risks of the study, and only those who gave their informed consents were formally enrolled.

The Lymphedema Quality of Life Questionnaire instruments

The Lymphedema quality of Life questionnaire (LYMQOL) was originally designed in 2004 by Keeley *et al.*^[7] and fully developed into the final version in 2010.^[6] With a structure similar to that of the EORTC QLQ-C30 (EORTC: The European Organization for Research and Treatment of Cancers Core Quality of Life Questionnaire 30),^[12] the content of the LYMQOL was based on interviews with 22 lymphedema patients and the opinions of lymphedema experts and physicians. Finally, some questions were deleted or revised to come into the final LYMQOL, adjusting to the feedback of 90 patients in Keeley *et al.*'s survey. The questionnaire comprises four domains: Function, Body image/Appearance, Symptoms and Mood. Each item is scored on a 4-point Likert scale ranging from "not at all" to "a lot", with values of 1 to 4". The higher the score, the worse the QoL. The total score of each domain is the sum of all item scores belonging to the domain divided by the number of questions answered. Unanswered questions are scored as 0, and the domain will be scored as 0 if more than 50% of questions are left unanswered in the domain. The overall QoL score, the last question of the questionnaire, ranges from 0 to 10. Higher overall QoL scores indicate better QoL ratings.

The LYMQOL consists of two subscales for the upper and lower limb lymphedema, respectively. The subscale LYMQOL-ARM, consisting of 21 items (Function aspect: items 1a-h, 2, 3, Body image/Appearance aspect: items 4-8, Symptoms aspect: items 9-14, Mood aspect: items 15-20, and the overall QoL: item 21), is used to assess QoL of patients with upper limb lymphedema. The subscale LYMQOL-LEG, containing 22 items (Function aspect: items 1a-f, 2, 3, Body image/Appearance aspect: items 4-10, Symptoms aspect: items 11-15, Mood aspect: items 16-21, and the

overall QoL: item 22), is applied to lymphedema involving lower limb extremity.

Keeley *et al.*^[6] has verified the reliability and validity of the LYMQOL in a group of 209 lymphedema patients, with Cronbach's α coefficients of domains of the LYMQOL-ARM and the LYMQOL-LEG being greater than 0.8 and a good correlation with the EORTCQLQ-C30 (The European Organization for Research and Treatment of Cancers Core Quality of Life Questionnaire 30). Most of the patients agreed that the questionnaire with an appropriate length was reasonable, clear and easy to complete.

The European Quality of Life-5 Dimensions Questionnaire (EQ-5D), a general health-related QoL questionnaire, has been widely used worldwide.^[13] The Chinese version of this questionnaire has been proved reliable and valid in cardiovascular diseases and chronic kidney diseases, etc.^[14, 15] The scale composes of five dimensions: Mobility, Self-Care, Usual Activities, Pain/Discomfort and Anxiety/Depression, in each dimension where has three questions. Because both China and Japan are located in the East Asia, the EQ-5D index in this study is scored by applying the Japanese Integral Conversion Table.^[16]

Changes of the affected limb circumference in lymphedema patients usually indicate progression of the disease,^[17] thus limb circumference measurements would be indispensable to lymphedema monitoring. Patients are encouraged to measure their affected limbs with a non-stretch tape at following points: wrist, 10 cm above wrist, 10 cm below elbow, elbow, and 10 cm above elbow for upper limb lymphedema; ankle, 10 cm above ankle, 10 cm below knee, and 10 cm above knee for lower limb lymphedema. By applying the truncated cone mathematical formula,^[18] the measured values are then converted into volumes, which can evaluate the concurrent validity of the LYMQOL.

Translation of the LYMQOL

Two lymphedema experts and two other native English speakers from the Department of Plastic Surgery of Peking Union Medical College Hospital were involved. Following the guideline for cross-cultural translation of QoL questionnaire,^[19, 20] two experts in lymphedema translated the questionnaire into Chinese; and then, the bilingual speakers who had no access to the original LYMQOL translated the draft back into English text. Finally, the authors (Zhengyun Liang, Xiao Long, Zhu-jun Li) and the translators involved together discussed

any discrepancy and revised the draft into the final Chinese version (**Appendix Tables 1, 2**). **Appendix Table 3** is the LYMQOL response scale.

Study population

We utilized the official Wechat account "Lymphedema Channel" to distribute the Chinese version of the questionnaire to patients who were clinically diagnosed with primary or secondary lymphedema from October 28th, 2020 to February 23rd, 2021. The patients should be at least 18 years old and able to read in Chinese. Those who presented with mental illness, acute stage of diseases, ongoing radiotherapy or chemotherapy, illiteracy or other diseases that might lead to chronic edema (such as heart or kidney failure) were excluded.

Basic information, relevant demographics (age, sex, body mass index, etc.) and clinical characteristics of lymphedema (affected side, site of lymphedema, duration of lymphedema, previous lymphedema treatment, cause of lymphedema, etc.) at the time enrolling in the study were collected. Patients with upper limb lymphedema completed the LYMQOL-ARM and EQ-5D simultaneously, and those with lower limb lymphedema filled in the LYMQOL-LEG and EQ-5D at the same time. If lymphedema affected both upper and lower limb, patients would fill out the LYMQOL-ARM and the LYMQOL-LEG concurrently. One week and one month after enrollment, the LYMQOL and the LYMQOL response scale were filled out by the participants again.

Statistical analysis

Internal consistency reliability The Cronbach's α coefficients of the domains were used to test homogeneity.^[21] Commonly accepted values were as follows: >0.9 excellent, >0.7 good, >0.6 acceptable.^[22] By comparing the first and second half, the odd and even half of each domain, the split-half reliability was tested.^[21] The Spearman-Brown coefficient values were categorized as: >0.8 excellent, >0.7 good, >0.6 acceptable.^[21]

Test-retest reliability The correlation between the 1st and the 2nd follow-up of the LYMQOL measurements were displayed as intra-class correlation coefficient (ICC)^[23] of domain scores and the overall QoL score of the LYMQOL to examine the test-retest reliability. ICC >0.75 was considered to be excellent.^[24, 25] One week was considered to be an appropriate time interval to examine test-retest reliability.^[26]

Content and face validity By utilizing the LYMQOL responses scale to obtain patients' opinions about

whether they think any questions were redundant or significant questions were missed out in the LYMQOL, the content and face validity were examined subjectively.

Criterion validity is determined by comparing the domain scores and the total score of the LYMQOL with that of the EQ-5D. It is hypothesized that the EQ-5D domain Mobility and Usual Activities correlate with the LYMQOL domain Function, the EQ-5D domain Self-Care correlates with the LYMQOL domain Body image/Appearance, the EQ-5D domain Pain/Discomfort correlates with the LYMQOL domain Symptoms and the EQ-5D domain Anxiety/Depression correlates with the LYMQOL domain Mood. The *Pearson's* correlation coefficient values were categorized as: 0.21-0.40 fair, 0.41-0.60 moderate, 0.61-0.80 good, >0.80 excellent^[27] and at least 75% of the hypotheses were confirmed.

Concurrent validity Correlations between the LYMQOL scores and the limb volume, as well as the LYMQOL scores and the lymphedema stage were conducted to test the concurrent validity. The *Pearson's correlation coefficient/the Spearman's* correlation coefficient values were categorized as forementioned.^[27] Lymphedema was staged according to the International Society of Lymphology (ISL) classification of lymphedema.

Construct validity An exploratory factor analysis (EFA) with a principal-axis factor analysis was conducted to analyze if the factor structure of the LYMQOL was the best fitting.^[28]

Feasibility was evaluated by the overall response rate (%) in the three follow-ups.

Responsiveness represents whether the LYMQOL is sensitive to change. It was assessed by measuring the changes in the LYMQOL scores with paired *t*-test between the 1st and the 3rd follow-up.

In addition, the average time required to finish the LYMQOL was calculated. Statistical analyses were performed with IBM SPSS 22.0 software. $P < 0.05$ was considered to be statistically significant.

RESULTS

General information

We enrolled a total of 195 patients diagnosed with lymphedema, of whom 93 with upper limb lymphedema, 100 with lower limb lymphedema, 2 cases of lymphedema affecting both upper limb and lower limb. The female accounted for 97.95%. Only 1.03% (2/195)

patients had primary lymphedema. The average age was 52.8 (23-81) years old. 46.15% (90/195) of the patients had a disease course of 1 to 5 years. 34.87% of lymphedema was not properly treated. The demographics and clinical characteristics of the participants are shown in **Table 1**.

Table 1. Demographics and clinical characteristics of the participants ($n=195$)

Items	Results
Gender [n (%)]	
Male	4 (2.05)
Female	191 (97.95)
Age [yrs, mean $\pm$ SD (range)]	52 $\pm$ 9 (23-81)
BMI [kg/m ² , mean $\pm$ SD (range)]	24.2 $\pm$ 3.1 (17.5-33.5)
Causes [n (%)]	
Endometrium cancer	24 (12.30)
Ovary cancer	14 (7.18)
Cervical cancer	52 (26.67)
Breast cancer	94 (48.21)
Other cancer	2 (1.03)
Lymphatics abnormality	2 (1.03)
Trauma	2 (1.03)
Others	5 (2.55)
Lymphedema site [n (%)]	
Upper limb	93 (47.69)
Lower Limb	100 (51.28)
Upper & lower limb	2 (1.03)
Affected site [n (%)]	
Left	96 (49.23)
Right	77 (39.49)
Bilateral	22 (11.28)
Duration [yrs, mean $\pm$ SD (range)]	3.7 $\pm$ 4.9 (0.1-37.6)
<1 [n (%)]	62 (31.79)
1-5 [including 1 yrs, n (%)]	90 (46.15)
5-10 [including 5 yrs, n (%)]	26 (13.33)
10-20 [including 10 yrs, n (%)]	12 (6.15)
≥ 20 [n (%)]	5 (2.56)
ISL staging [n (%)]	
Stage I	116 (59.49)
Stage II	35 (17.95)
Stage III	44 (22.56)
Treatment [n (%)]	
No previous treatment	68 (34.87)
Conservative therapy	114 (58.46)
Surgery	2 (1.05)
Both treatment	11 (5.64)

BMI: Body Mass Index; ISL: International Society of Lymphology.

Internal consistency reliability

The Cronbach's α coefficients for all the domains were above 0.7, which indicated a high internal reliability of the Chinese questionnaire. Except that Spearman-Brown coefficient of Symptoms for the LYMQOL-LEG was only 0.638, the coefficients for other domains were above 0.7. (Table 2)

Test-retest reliability

ICC revealed a good consistency in LYMQOL score between 1 week and 1 month after enrollment (ICC>0.800, $P<0.001$, Table 3).

Content and face validity

According to response of the patients to our questionnaire, we found over 90% patients considered that the LYMQOL was comprehensive, succinct, and easy to answer (Table 4). Moreover, most of participants regarded the issues concerning sex life, cost and curative

effect, etc. should be included.

Criterion validity

The EQ-5D index scores positively correlated overall QoL of the LYMQOL ($0.4<r<0.6$, $P<0.01$). Comparison between the EQ-5D index score and the LYMQOL domain scores demonstrated moderate to good correlation, except for the LYMQOL Appearance domain. 80% of the forementioned hypotheses were confirmed. Thus, criterion validity was acceptable between the EQ-5D and the LYMQOL (Table 5).

Concurrent validity

For the LYMQOL-ARM, no correlation was found between the LYMQOL domains and ISL staging or limb volume (for ISL staging, $n=95$, $r=0.062$, $P>0.5$ and for limb volume, $n=84$, $r=-0.034$, $P>0.5$). For the LYMQOL-LEG, ISL staging positively correlated with Symptoms domain ($n=102$, $r=0.297$, $P<0.01$), and

Table 2. Internal consistency reliability of the LYMQOL

Domains	The LYMQOL-ARM ($n=95$)			The LYMQOL-LEG ($n=102$)		
	α	Spearman-Brown coefficient		α	Spearman-Brown coefficient	
		1 st /2 nd half	Odd/Even		1 st /2 nd half	Odd/Even
Function	0.902	0.925	0.948	0.861	0.840	0.866
Appearance	0.889	0.843	0.910	0.902	0.882	0.935
Symptoms	0.849	0.811	0.888	0.726	0.638	0.733
Mood	0.877	0.849	0.904	0.834	0.777	0.879

LYMQOL: Lymphedema Quality of Life Questionnaire.

Table 3. Test-retest reliability of the LYMQOL

Domains	The LYMQOL-ARM ($n=74$)		The LYMQOL-LEG ($n=83$)	
	ICC	P	ICC	P
Function	0.848	<0.001	0.900	<0.001
Appearance	0.884	<0.001	0.886	<0.001
Symptoms	0.866	<0.001	0.863	<0.001
Mood	0.874	<0.001	0.894	<0.001
Overall QoL	0.871	<0.001	0.868	<0.001

ICC: intra-class correlation coefficient; QoL: quality of life.

Table 4. Content and face validity of the LYMQOL

Items	The LYMQOL-ARM ($n=77$)		The LYMQOL-LEG ($n=83$)	
	Yes (%)	No (%)	Yes (%)	No (%)
Is it easy to complete?	94.8	5.2	96.4	3.6
Is it too long?	14.3	85.7	7.2	92.8
Are all the questions clear?	92.2	7.8	91.6	8.4
Are there any important areas not covered?	15.6	84.4	26.5	73.5
Are there any questions redundant?	3.9	96.1	4.8	95.2

Table 5. Criterion validity of the LYMQOL (Pearson's correlation coefficient)

The EQ-5D	The LYMQOL-ARM (n=95)					The LYMQOL-LEG (n=102)				
	Function	Appearance	Symptoms	Mood	Overall QoL	Function	Appearance	Symptoms	Mood	Overall QoL
Mobility	0.198					0.471**				
Self-Care		0.333**					-0.002			
Usual activities	0.498**					0.630**				
Discomfort			0.626**					0.508**		
Depression				0.680**					0.438**	
EQ-5D index	-0.617**	-0.377**	-0.727**	-0.633**	0.552**	-0.662**	-0.267**	-0.511**	-0.507**	0.555**

EQ-5D: European Quality of Life-5 Dimensions Questionnaire. ** $P < 0.01$

limb volume positively correlated with Appearance domain ($n=87$, $r=0.313$, $P < 0.01$).

Construct validity

Since the Kaiser-Meyer-Olkin (KMO) was above 0.8 and in Bartlett's test P was below 0.05, which means the sample size is sufficient,^[29] an EFA was conducted. All of the factor loading values were above 0.4, indicating no item should be deleted from the LYMQOL. We

identified a four-factor structure in the Chinese questionnaire, similar to that of the original version, with the same items in the same factors. These four factors explained 65.282% and 59.984% of the total variance in the LYMQOL-ARM and the LYMQOL-LEG, each accounting for 19.756%, 17.597%, 14.257%, 13.672% and 17.198%, 16.738%, 14.291%, 11.756%, respectively. Factor loadings and factor names are shown in

Table 6.

Table 6. Factor loadings of the LYMQOL-ARM and of the LYMQOL-LEG (KMO)

Items	LYMQOL-ARM				LYMQOL-LEG			
	Factor 1	Factor 2	Factor 3	Factor 4	Factor 1	Factor 2	Factor 3	Factor 4
1a	0.527	0.452	0.296	0.163	0.061	0.024	0.822	0.307
1b	0.545	0.497	0.130	0.176	0.063	0.127	0.279	0.658
1c	0.183	0.215	0.120	0.789	0.004	0.089	0.592	0.477
1d	0.475	0.302	0.076	0.420	0.065	0.197	0.458	0.447
1e	0.285	0.158	0.161	0.793	0.330	0.000	0.424	0.554
1f	0.021	0.154	0.108	0.906	0.179	-0.016	0.677	0.429
1g	0.414	0.506	0.155	0.272				
1h	0.100	0.194	0.281	0.839				
2	0.618	0.406	0.227	0.215	0.368	0.431	0.565	0.002
3	0.549	0.419	0.201	0.207	0.531	0.166	0.196	0.324
4	0.758	0.111	0.269	0.138	0.324	0.786	0.217	-0.074
5	0.873	0.039	-0.003	0.044	0.216	0.841	-0.013	0.092
6	0.883	-0.031	0.013	0.124	0.186	0.885	0.017	0.096
7	0.775	0.025	0.381	0.071	0.020	0.729	0.109	0.196
8	0.525	0.356	0.387	0.059	0.203	0.767	0.075	0.178
9	-0.020	0.681	0.176	0.043	0.579	0.548	0.083	0.087
10	0.137	0.659	0.154	0.166	0.547	0.459	0.201	0.004
11	0.047	0.768	0.088	0.019	0.152	0.102	0.082	0.721
12	0.184	0.725	0.043	0.375	0.288	-0.001	0.472	0.228
13	0.357	0.589	0.142	0.309	0.076	0.096	0.238	0.657
14	0.139	0.658	0.301	0.186	0.349	-0.018	0.551	0.317
15	0.044	0.515	0.497	0.146	0.204	0.332	0.669	-0.067
16	0.223	0.405	0.583	0.260	0.499	0.065	0.019	0.419
17	0.342	0.107	0.742	0.150	0.649	0.153	-0.015	0.320
18	0.423	0.197	0.717	0.140	0.740	0.083	0.257	0.125
19	0.052	0.172	0.819	0.078	0.728	0.236	0.299	0.017
20	0.099	0.179	0.793	0.189	0.682	0.278	0.027	0.057
21					0.742	0.191	0.312	-0.013

Bold font: highest value for each item. KMO: the Kaiser-Meyer-Olkin value.

For LYMQOL-ARM, KMO=0.876, Bartlett's test: $P=0.000$. For LYMQOL-LEG, KMO=0.870, Bartlett's test: $P=0.000$.

Feasibility

A total of 195 patients (100%) completed the questionnaires of the first survey. The response rates of the second and third surveys were 77.9% (the LYMQOL-ARM), 81.4% (the LYMQOL-LEG) and 74.7% (the LYMQOL-ARM), 76.4% (the LYMQOL-LEG).

Responsiveness

Paired *t*-test showed that for the LYMQOL-ARM, compared with the the first survey, all domains scores of the third survey increased significantly, except for overall QoL ($P < 0.05$, **Table 7**); for the LYMQOL-LEG, however, no difference was detected in all domains scores compared with the first survey ($P > 0.05$, **Table 7**).

Time to finish the questionnaire

The average time required for patients to finish the LYMQOL-ARM and the LYMQOL-LEG was 746 (range: 94-3940) s and 774 (range: 122-7023) s, respectively.

DISCUSSION

Currently, there are two main types of health related QoL measures for lymphedema: one is the general QoL, and the other one is condition-specific QoL questionnaires. The latter consists of cancer-specific and lymphedema-specific. Commonly used general health questionnaires include the Medical Outcome Study-36 Item Short Form (SF-36),^[30] the Nottingham Health Profile part 1 (NHP-1),^[31] and the EQ-5D scale,^[13] etc. These types of questionnaires are very effective in evaluating the diseases' overall negative impact on QoL, but not as accurate, sensitive or informative as a condition-specific tool especially when it comes to monitoring or detecting subtle changes over time.^[4, 32] Cancer-specific instruments encompass the Functional Assessment of Cancer Therapy-Breast can-

cer questionnaire (FACT-B)^[33] for breast cancer survivor assessment and the Functional Living Index-Cancer (FLIC),^[34] etc. These types of questionnaires are restricted for QoL assessment of patients with secondary lymphedema caused by malignant tumors, and are not used for primary lymphedema nor for non-cancer related secondary lymphedema. Additionally, some of the questions focus on the tumor instead of lymphedema, which is redundant. Lymphedema-specific instruments include Lymphoedema Functioning, Disability and Health questionnaire (Lymph-ICF),^[35] the Lymphedema Quality of Life Inventory (LyQLI),^[36] the LYMQOL,^[6] and etc., which all have been verified to be reliable and effective. Patient-reported outcome measures provide a new dimension for the overall evaluation of lymphedema apart from objective measures such as imaging examination and limb circumference. Supplementary information may assist clinical decision-making and intervention effectiveness monitoring. The LYMQOL has been localized and validated in terms of psychometric properties in the United Kingdom,^[6] Turkey,^[9-11, 37, 38] the Netherlands,^[8] Ireland,^[39] Italy,^[40, 41] Japan^[42] and Sweden,^[43] with their own language versions, respectively.

This study aims to introduce the LYMQOL, translate it into Chinese with reference to the cross-cultural adaptation guidelines of questionnaires and validate the psychometric properties of the LYMQOL among lymphedema patients in China. The results from 195 participants show that the LYMQOL owns very satisfactory overall reliability and validity.

In the study, the Cronbach's *a* coefficients of the LYMQOL-ARM were 0.849-0.902 with ICC being 0.848-0.884. While in the translated Turkish version, the Cronbach's *a* coefficients were 0.70-0.94, with ICC being 0.98-0.99.^[9] Reported domain Cronbach's *a* coefficients range from 0.79 to 0.93, with ICC ranging from 0.53 to 0.87 in the validation study of the Swedish version of the LYMQOL.^[43] Similarly, the Cronbach's *a* coefficients of the LYMQOL-LEG were 0.726-0.902, with ICC being 0.863-0.900 in our study. While good internal consistency and test-retest reliability were also observed in the Turkish version (the Cronbach's *a* coefficients: 0.74-0.94, ICC: 0.83-0.95),^[10] the Swedish version (the Cronbach's *a* coefficients: 0.88-0.94, ICC: 0.78-0.91)^[43] and the Dutch version (the Cronbach's *a* coefficient: 0.89, ICC: 0.73-0.91)^[44] of the LYMQOL-LEG. Based on these studies, it might be safe to say that both the LYMQOL-ARM and the LYMQOL-LEG

Table 7. Comparisons of the domain scores between the first and the third survey for the LYMQOL-ARM and LYMQOL-LEG

Domains	LYMQOL-ARM		LYMQOL-LEG	
	<i>t</i> value	<i>P</i> value	<i>t</i> value	<i>P</i> value
Function	-5.071	0.000	-1.901	0.063
Appearance	-3.302	0.002	0.348	0.729
Symptoms	-4.000	0.000	-1.378	0.175
Mood	-2.694	0.010	-0.898	0.374
Overall QoL	1.480	0.145	1.600	0.116

are stable and excellent in internal consistency and test-retest reliability.

The EQ-5D scale is utilized to assess the criterion validity of the Chinese version of the LYMQOL in our study. The fact that the EQ-5D scale is a general QoL questionnaire and does not comprise questions concerning appearance, may explain why the weakest correlation is detected between the EQ-5D index and the LYMQOL Appearance domain. This methodology of using a generic questionnaire as “the gold standard” is also seen in other researches.^[6, 8-11, 37, 38] Wang *et al.* translate and validate the Lymph-ICF-Lower Limb subscale (the Lymph-ICF-LL) among the Chinese lower limb lymphedema patients in 2018.^[45] Generally speaking, a lymphedema-specific tool is more ideal to be used as the criterion than a generic measure. The study do not choose Lymph-ICF-LL as the criterion due to: 1. The study includes both upper and lower limb lymphedema patients, thus the Lymph-ICF-LL translated by Wang *et al.* might not be as sufficient as the criterion; 2. The Lymph-ICF-LL consists of 28 items in total, the increase in number of questionnaires items might lead to patient fatigue, lower patient compliance and response rate; 3. The study aims to verify the reliability and validity of the Chinese version of the LYMQOL, thus using the EQ-5D scale may be sufficient. Further research may focus on the strengths and weaknesses analysis between the Lymph-ICF and the LYMQOL in terms of content, questionnaire length, response rate, and etc.

Positive correlations are detected only between the Symptoms domain and the ISL staging, as well as between the Appearance domain and the limb volume for the LYMQOL-LEG. The findings are reasonable because the ISL staging is mainly based on lymphedema symptoms and the larger the limb volume, the greater the affected body image. Other than that, no significant correlations exist between the ISL staging/the limb volume and the LYMQOL. It is consistent with the findings of Keeley *et al.*^[6] One of the plausible reasons is that, patients' QoL is already damaged in early stages of lymphedema, however, limb volume has not increased accordingly. As for the ISL staging, it stratifies patients chiefly according to the symptoms, however, lymphedema affects both the physiological and psychological aspects of the patients. Therefore, disease-specific QoL questionnaires might provide additional information in the severity assessment of lymphedema apart from clinical staging and limb volumes.

When Keeley *et al.* evaluated the responsiveness of the LYMQOL, no significant differences are observed between one week and one month data.^[6] Same results are found in our study for the LYMQOL-LEG. For the LYMQOL-ARM, however, significant increase in all four domain scores (lymphedema worsens) is detected, partially confirming the responsiveness of the LYMQOL. Moreover, it suggests that the LYMQOL might be more sensitive in tracking changes for upper limb lymphedema than for lower limb lymphedema. Larger sample size and longer follow-up terms are in need to further assess the responsiveness of the LYMQOL.

The LYMQOL is divided into two subscales: the LYMQOL-ARM and the LYMQOL-LEG, respectively designed for the evaluation of upper and lower limb lymphedema. Since these two subscales share the same rating rule with similar questionnaire structure, it might facilitate data comparison between the two lymphedema groups. The representative of questionnaires specifically designed for upper limb lymphedema is the Upper Limb Lymphedema 27 (ULL27, 27 items in total),^[46] which has been validated in Turkey^[47] and France^[48]. The Italian IDI-ILA scale (27 items in total),^[49] is one of the very few lower limb lymphedema-specific questionnaires, but has only been used in lymphedema secondary to melanoma with unknown data on its reliability and validity. Compared with these two types of questionnaires, the LYMQOL can be used for both upper and lower limb lymphedema assessment and can serve as more convenient in clinical setting. There are other scales like the LYMQOL, such as the Lymph-ICF (Upper limb subscale: 28 items; Lower limb subscale: 28 items),^[50, 51] the LSIDS (Upper limb subscale: 30 items; Lower limb subscale: 31 items).^[52, 53] These scales, including LYMQOL are not perfect: especially when assessing lymphedema manifesting in the perineum and trunk.

As for lymphedema patients with both upper and lower limb affected, they are required to finish the LYMQOL-ARM and the LYMQOL-LEG. Undoubtedly, this gives rise to certain bias for arm lymphedema affecting the overall QoL of the LYMQOL-LEG or vice versa. Until now, there are still no specific questionnaires targeting this type of patients.

The LYMQOL assesses both the overall QoL and also four important domains for the affected arm or leg, thus we deem it is effective in assessing patients with both upper and lower limb lymphedema. Plus, there are only 2 patients who fit in this category in our

study, so it is included in our data analysis. Of note, the most appropriate way to approach this type of patients still remain to be discussed for future study.

It is believed that the length of the self-reported questionnaires negatively correlates with the response rate,^[54] as the time required for completion increases for lengthy questionnaires. Therefore, consideration must be given to the length and the time required for completion for patient-reported outcome measures. The LYMQOL might be superior to the Lymph-ICF and the LSIDS, etc. when taking the length of the questionnaire into account. Plus, the majority of the participants agree that the content of the LYMQOL is comprehensive and closely related to their lymphedema troubles in daily life. The response rate is also high (>75%), with the average time required for patients to finish the LYMQOL-ARM and the LYMQOL-LEG being 746 s (range: 94-3940 s) and 774 s (range: 122-7023 s), respectively. In the previous studies, the estimated average time for completion is 5 minutes,^[10] which is shorter than that in our study. The reason may be due to the use of a semi-automated electronic platform to distribute the LYMQOL, which is unlike filling out the questionnaires on the spot. In other words, patients are more likely to be distracted by other affairs or disturbed by terrible network connection leading to an unexpected disruption of the process and prolonged answer time. Up to the time of our study, there is no specific data regarding the actual time required for completion from other studies, thus making statistic comparison difficult. However, given the fact that lymphedema is a chronic progressive disease in nature, which requires long-term management and multiple follow-ups, the time required for completion should be an important aspect to consider when choosing QoL tool. As patients become more familiar with the LYMQOL, the completion time shortens in our study. Therefore, the LYMQOL is potentially useful for long-term assessment of patients with lymphedema in terms of questionnaire length and time.

As of 2020, the number of Wechat monthly active users has reached 1.213 billion. As a functioning social application in our mobile phone, Wechat provides its users with easy access to large quantities of scientific, educational and medical information. The official account and Mini Program are two of the most commonly used functions for Wechat users. This study uses Wechat, an electronic mobile social platform for questionnaire follow-up survey, and the response rate

of with lymphedema patients is found to be high (the 1st follow-up: 100%, the 2nd follow-up: 77%-81%, the 3rd follow-up: 74%-76%), suggesting the feasibility of lymphedema follow-up on electronic platform. For chronic refractory diseases such as lymphedema, electronic patient-reported outcome measures enable health care team to track the changes of the patients' condition regardless of time and space, and minimize unnecessary clinic visits, freeing patients from the troublesome arrangement for work and transportation. Eventually it leads to more efficient medical data collection and reduced medical burden.^[55, 56] The comparison of the cost-effectiveness with the conventional paper questionnaire is yet to be explored.

In conclusion, the Chinese version of the LYMQOL is easy to fill out, appropriate in length and comprehensive in content with good reliability and validity when used to evaluate the QoL of Chinese patients with lymphedema. Therefore, it might be safe to conclude that the Chinese LYMQOL is a validated patient-reported outcome measure tool, which can be utilized to assist treatment selections and track changes in clinical setting or facilitate future study on lymphedema.

The Supporting Information is available on the CMSJ website at doi: 10.24920/003918.

Appendix Table 1. Chinese version of the LYMQOL-ARM.

Appendix Table 2. Chinese version of the LYMQOL-LEG.

Appendix Table 3. Chinese version of the LYMQOL responses scale.

Conflict of Interest Disclosures

We have no conflicts of interest to declare.

REFERENCES

1. Liang ZY, Long X, Yu NZ, et al. Diagnostic workup of lymphedema. *Plast Aesthet Res* 2019; 6:23. doi: 10.20517/2347-9264.2019.33.
2. Grada AA, Phillips TJ. Lymphedema: pathophysiology and clinical manifestations. *J Am Acad Dermatol* 2017; 77(6):1009-20. doi: 10.1016/j.jaad.2017.03.022.
3. Bowling A, Iliffe S. Psychological approach to successful aging predicts future quality of life in older adults. *Health Qual Life Outcomes* 2011; 9:13. doi: 10.1186/1477-7525-9-13.

4. Keeley V. Quality of life assessment tools in chronic oedema. *Br J Community Nurs* 2008; 13(10):S22-7. doi: 10.12968/bjcn.2008.13.Sup5.31193.
5. Higginson I, Carr AJ. Measuring quality of life: using quality of life measures in the clinical setting. *BMJ* 2001; 322(7297):1297-300. doi: 10.1136/bmj.322.7297.1297.
6. Keeley VN, Crooks S, Locke J, et al. A quality of life measure for limb lymphoedema (LYMQOL). *J Lymphoedema* 2010; 5(1):26-37.
7. Keeley V, Veigas D, Crooks S. The development of a condition-specific quality of life measure for lymphoedema (LYMQOL). *Eur J Lymphol* 2004; 12:36.
8. van de Pas C, Biemans A, Boonen R, et al. Validation of the Lymphoedema Quality-of-Life Questionnaire (LYMQOL) in Dutch patients diagnosed with lymphoedema of the lower limbs. *Phlebology* 2016; 31(4):257-63. doi: 10.1177/0268355515586312.
9. Bakar Y, Tuğral A, Özdemir Ö, et al. Translation and validation of the Turkish version of lymphedema quality of life tool (LYMQOL) in patients with breast cancer related lymphedema. *Eur J Breast Health* 2017; 13(3):123-8. doi: 10.5152/ejbh.2017.3522.
10. Bakar Y, Tuğral A. Translation, reliability, and validation of the Turkish version of the Lymphedema Quality-of-Life tool in Turkish-speaking patients with lower limb lymphedema. *J Vasc Nurs* 2019; 37(1):11-7. doi: 10.1016/j.jvn.2018.11.005.
11. Karayurt Ö, Deveci Z, Eyigör S, et al. Adaptation of quality of life measure for limb lymphedema-arm in Turkish women with breast cancer-related lymphedema. *Cancer Nurs* 2021; 44(1):45-52. doi: 10.1097/NCC.0000000000000741.
12. Sprangers MA, Cull A, Groenvold M, et al. The European Organization for Research and Treatment of Cancer approach to developing questionnaire modules: an update and overview. *EORTC Quality of Life Study Group. Qual Life Res* 1998; 7(4):291-300. doi: 10.1023/a:1024977728719.
13. Devlin NJ, Brooks R. EQ-5D and the EuroQol Group: Past, Present and Future. *Applied health economics and health policy. Appl Health Econ Health Policy* 2017; 15(2):127-37. doi: 10.1007/s40258-017-0310-5.
14. Chang TJ, Tarn YH, Hsieh CL, et al. Taiwanese version of the EQ-5D: validation in a representative sample of the Taiwanese population. *J Formos Med Assoc* 2007; 106(12):1023-31. doi: 10.1016/S0929-6646(08)60078-9.
15. Wang H, Kindig DA, Mullahy J. Variation in Chinese population health related quality of life: results from a EuroQol study in Beijing, China. *Qual Life Res* 2005; 14(1):119-32. doi: 10.1007/s11136-004-0612-6.
16. Tsuchiya A, Ikeda S, Ikegami N, et al. Estimating an EQ-5D population value set: the case of Japan. *Health Econ* 2002; 11(4):341-53. doi: 10.1002/hec.673.
17. Spillane A, Saw R, Tucker M, et al. Defining lower limb lymphedema after inguinal or ilio-inguinal dissection in patients with melanoma using classification and regression tree analysis. *Ann Surg* 2008; 248(2):286-93. doi: 10.1097/SLA.0b013e31817ed7c3.
18. Hidding JT, Viehoff PB, Beurskens CH, et al. Measurement properties of instruments for measuring of lymphedema: systematic review. *Phys Ther* 2016; 96(12):1965-81. doi: 10.2522/ptj.20150412.
19. Donnally C, Lugo-Pico J, Bondar K, et al. Characteristics and trends of the most cited spine publications. *Spine (Phila Pa 1976)* 2021; 46(11):765-71. doi: 10.1097/BRS.0000000000003890.
20. Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol* 1993; 46(12):1417-32. doi: 10.1016/0895-4356(93)90142-n.
21. Eisinga R, Grotenhuis M, Pelzer B. The reliability of a two-item scale: Pearson, Cronbach, or Spearman-Brown? *Int J Public Health* 2013; 58(4):637-42. doi: 10.1007/s00038-012-0416-3.
22. Bhatnagar R, Kim J, Many JE. Candidate surveys on program evaluation: examining instrument reliability, validity and program effectiveness. *Am Educ Res J* 2014; 51(2):683-90. doi: 10.12691/education-2-8-18.
23. Fleiss JL, Cohen J. The equivalence of weighted kappa and the intraclass correlation coefficient as measures of reliability. *Educ Psychol Measurement* 1973; 33(3):613-9.
24. Shamliyan T, Kane RL, Dickinson S. A systematic review of tools used to assess the quality of observational studies that examine incidence or prevalence and risk factors for diseases. *J Clin Epidemiol* 2010; 63(10):1061-70. doi: 10.1016/j.jclinepi.2010.04.014.
25. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med* 2016; 15(2):155-63. doi: 10.1016/j.jcm.2016.02.012.
26. Marx RG, Menezes A, Horovitz L, et al. A comparison of two time intervals for test-retest reliability of health

- status instruments. *J Clin Epidemiol* 2003; 56(8):730-5. doi: 10.1016/s0895-4356(03)00084-2.
27. Feise RJ, Michael MJ. Functional rating index: a new valid and reliable instrument to measure the magnitude of clinical change in spinal conditions. *Spine Spine (Phila Pa 1976)* 2001; 26(1):78-86; discussion 87. doi: 10.1097/00007632-200101010-00015.
28. Kaiser HF. The varimax criterion for analytic rotation in factor analysis. *Edu Psychol Measurement* 1958; 23:770-3.
29. Costello AB, Osborne JW. Best practices in exploratory factor analysis: four recommendations for getting the most from your analysis. *Practical Assessment Research Evaluation* 2005; 10(7):1-9.
30. Ware JE Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992; 30(6):473-83.
31. Hunt SM, McEwen J. The development of a subjective health indicator. *Sociol Health Illn* 1980; 2(3):231-46. doi: 10.1111/1467-9566.ep11340686.
32. Pusic A, Cemal Y, Albornoz C, et al. Quality of life among breast cancer patients with lymphedema: a systematic review of patient-reported outcome instruments and outcomes. *J Cancer Surviv* 2013; 7(1):83-92. doi: 10.1007/s11764-012-0247-5.
33. Brady MJ, Cella DF, Mo F, et al. Reliability and validity of the Functional Assessment of Cancer Therapy-Breast quality-of-life instrument. *J Clin Oncol* 1997; 15(3):974-86. doi: 10.1200/JCO.1997.15.3.974.
34. Schipper H, Clinch J, McMurray A, et al. Measuring the quality of life of cancer patients: the Functional Living Index-Cancer: development and validation. *J Clin Oncol* 1984; 2(5):472-83. doi: 10.1200/JCO.1984.2.5.472.
35. Devoogdt N, Van K M, Geraerts I, et al. Lymphoedema Functioning, Disability and Health questionnaire (Lymph-ICF): reliability and validity. *Phys Ther* 2011; 91(6):944-57. doi: 10.2522/ptj.20100087.
36. Klernas P, Johnsson A, Horstmann V, et al. Lymphedema Quality of Life Inventory (LyQLI)-Development and investigation of validity and reliability. *Qual Life Res* 2015; 24(2):427-39. doi: 10.1007/s11136-014-0783-8.
37. Borman P, Yaman A, Denizli M, et al. The reliability and validity of Lymphedema Quality of Life Questionnaire-Arm in Turkish patients with upper limb lymphedema related with breast cancer. *Turk J Phys Med Rehabil* 2018; 64(3):205-12. doi: 10.5606/tftrd.2018.2843.
38. Borman P, Yaman A, Denizli M, et al. The reliability and validity of Lymphedema Quality of Life Questionnaire-Leg in Turkish patients with lower limb lymphedema. *Lymphat Res Biol* 2020; 18(1):42-8. doi: 10.1089/lrb.2018.0048.
39. Greene A, Meskell P. The impact of lower limb chronic oedema on patients' quality of life. *Int Wound J* 2017; 14(3):561-8. doi: 10.1111/iwj.12648.
40. Salgarello M, Mangialardi M, Pino V, et al. A prospective evaluation of health-related quality of life following lymphaticovenular anastomosis for upper and lower extremities lymphedema. *J Reconstr Microsurg* 2018; 34(9):701-7. doi: 10.1055/s-0038-1642623.
41. Monticone M, Ferriero G, Keeley V, et al. Lymphedema quality of life questionnaire (LYMQOL): cross-cultural adaptation and validation in Italian women with upper limb lymphedema after breast cancer. *Disabil Rehabil* 2021;1-6. doi: 10.1080/09638288.2021.1890842.
42. Iuchi T, Dai M, Sanada H, et al. Associations between the treatments and outcomes of patients with upper and lower lymphoedema in Japan: a cross-sectional observational study. *Int J Nurs Stud* 2015; 52(5):913-9. doi: 10.1016/j.ijnurstu.2015.01.011.
43. Wedin M, Fredrikson M, Ahlner E, et al. Validation of the Lymphoedema Quality of Life Questionnaire (LYMQOL) in Swedish cancer patients. *Acta Oncol* 2020; 59(3):365-71. doi: 10.1080/0284186X.2019.1701199.
44. van de Pas CB, Biemans AA, Boonen RS, et al. Validation of the Lymphoedema Quality-of-Life Questionnaire (LYMQOL) in Dutch patients diagnosed with lymphoedema of the lower limbs. *Phlebology* 2016; 31(4):257-63. doi: 10.1177/0268355515586312.
45. Wang C, Lee S, Hsu K, et al. Psychometric evaluation of a Chinese version of Lymphoedema Functioning, Disability and Health Questionnaire for Lower Limb Lymphoedema in women with gynaecological cancer surgery. *Eur J Cancer Care (Engl)* 2018; 27(6):e12940. doi: 10.1111/ecc.12940.
46. Launois R, Alliot F. Quality of Life Scale in Upper Limb Lymphedema—a validation study. *Lymphology* 2000; 33:226-74.
47. Kayali VA, Yavuzşen T, Karadibak D. The reliability and validity of Quality of Life Questionnaire Upper Limb Lymphedema (ULL-27) Turkish patient with breast cancer related lymphedema. *Front Oncol* 2020; 10:455. doi: 10.3389/fonc.2020.00455.
48. Williams A, Rapport F, Russell I, et al. Psychometric development of the Upper Limb Lymphedema Quality of Life Questionnaire demonstrated the patient-reported

- outcome measure to be a robust measure for breast cancer-related lymphedema. *J Clin Epidemiol* 2018; 100:61-70. doi: 10.1016/j.jclinepi.2018.04.014.
49. Carmeli E, Bartoletti R. Retrospective trial of complete decongestive physical therapy for lower extremity secondary lymphedema in melanoma patients. *Support Care Cancer* 2011; 19(1):141-7. doi: 10.1007/s00520-009-0803-3.
50. Devoogdt N, De Groef A, Hendrickx A, et al. Lymphoedema Functioning, Disability and Health Questionnaire for Lower Limb Lymphoedema (Lymph-ICF-LL): reliability and validity. *Phys Ther* 2014; 94(5):705-21. doi: 10.2522/ptj.20130285.
51. Devoogdt N, Van Kampen M, Geraerts I, et al. Lymphoedema Functioning, Disability and Health questionnaire (Lymph-ICF): reliability and validity. *Phys Ther* 2011; 91(6):944-57. doi: 10.2522/ptj.20100087.
52. Ridner SH, Dietrich MS. Development and validation of the Lymphedema Symptom and Intensity Survey-Arm. *Support Care Cancer* 2015; 23(10):3103-12. doi: 10.1007/s00520-015-2684-y.
53. Ridner SH, Doersam JK, Stollendorf DP. Development and validation of the lymphedema symptom intensity and distress survey-lower limb. *Lymphat Res Biol* 2018; 16(6):538-46. doi: 10.1089/lrb.2017.0069.
54. Edwards P, Roberts I, Clarke M, et al. Increasing response rates to postal questionnaires: systematic review. *BMJ* 2002; 324(7347):1183. doi: 10.1136/bmj.324.7347.1183.
55. Hoang N, Hwang W, Katz D, et al. Electronic patient-reported outcomes: semi-automated data collection in the Interventional Radiology Clinic. *J Am Coll Radiol* 2019; 16(4 Pt A):472-7. doi: 10.1016/j.jacr.2018.08.033.
56. Rogausch A, Sigle J, Seibert A, et al. Feasibility and acceptance of electronic quality of life assessment in general practice: an implementation study. *Health Qual Life Outcomes* 2009; 7:51. doi: 10.1186/1477-7525-7-51.

中文摘要

淋巴水肿生活质量问卷（LYMQOL）信效度研究

梁铮韵¹，龙笑^{1*}，杨伊兰¹，李云竹¹，李竹君¹，

徐敏²，张毕分²，俞楠泽¹，黄久佐¹

¹ 中国医学科学院 北京协和医学院 北京协和医院整形美容外科，北京 100005

² 深圳原位医疗设备有限公司，深圳 518116

摘要

目的 验证淋巴水肿生活质量问卷（LYMQOL）在中国淋巴水肿患者中的信度和效度。

方法 在2020年10月28日至2021年2月23日期间，我们将翻译成中文版的淋巴水肿生活质量问卷（LYMQOL）通过微信公众号“淋巴水肿频道”推送给纳入的淋巴水肿患者。上肢淋巴水肿和下肢淋巴水肿患者分别在纳入时、纳入后1周和1个月完成LYMQOL-ARM子量表和LYMQOL-LEG子量表。我们对问卷的信度、效度、可行性、应答性和完成问卷所需平均时间进行了评估。

结果 研究共纳入195例患者。中文版LYMQOL问卷具有较高的信度，LYMQOL-ARM的 α 系数为0.849~0.902，组内相关系数（ICC）为0.848~0.884，LYMQOL-LEG的 α 系数为0.726~0.902，ICC为0.863~0.900。LYMQOL与EQ-5D两者呈中度至良好相关（LYMQOL-ARM， $n=95$ ；LYMQOL-LEG， $n=102$ ）。应答性分析结果提示：术后1个月上肢淋巴水肿患者的生活质量明显下降（ $P<0.05$ ）。患者完成问卷的平均时间约为12分钟。

结论 中文版LYMQOL简明易懂、内容全面、篇幅适中，具有良好的信效度。在临床或科研中，或许可用于协助治疗决策和淋巴水肿疗效评估。

关键词：淋巴水肿生活质量问卷；淋巴水肿；生活质量；效度；信度