

CHINESE MEDICAL SCIENCES JOURNAL

ORIGINAL ARTICLE

Efficacy and Safety of Low Molecular Weight Heparin Prophylaxis for Venous Thromboembolism Following Lumbar Decompression Surgery

Zhi-jian Sun, Yu Zhao*, Giu-xing Qiu, Yi-peng Wang, Xi-sheng Weng, Hong Zhao, Jian-xiong Shen, Yu Jiang, Ye Li, and Xiang Li

Department of Orthopedics, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100730, China

Key words: venous thromboembolism; prophylaxis; low molecular weight heparin; lumbar decompression surgery

Objective To evaluate the efficacy and safety of low molecular weight heparin (LMWH) prophylaxis for venous thromboembolism (VTE) after lumbar decompression surgery.

Methods Patients at high or the highest risk of VTE who underwent lumbar spine surgery in Peking Union Medical College Hospital from January 2004 to April 2011 were included in the present study. All the patients received a half dose of LMWH 6 hours after surgery followed by a full dose LMWH once per day until discharge. We recorded incidences of deep venous thrombosis (DVT), pulmonary embolism (PE), bleeding complications, and medication side effects.

Results Seventy-eight consecutive patients were eligible and enrolled in this study. The mean hospital stay was 8.5 ± 4.5 days. No symptomatic DVT, PE, or major bleeding events were observed. One patient developed wound ecchymosis, another developed wound bleeding, four had mild hepatic aminotransferase level elevation, and one developed a suspicious allergic reaction.

Conclusion LMWH may be applied as an effective and safe prophylaxis for VTE in high-risk patients undergoing lumbar decompression surgery.

Chin Med Sci J 2011; 26(4):221-226

VENOUS thromboembolism (VTE) includes deep venous thrombosis (DVT) and pulmonary embolism (PE). Total knee arthroplasty, total hip arthroplasty, and hip fracture surgery are all associated with a high incidence of postoperative VTE,

resulting in high mortality.^{1,2} There is a consensus on standardised VTE prophylaxis for these operations.^{3,4} However, few reports have been published discussing the use of prophylactic anticoagulation following spine surgery. Glotzbecker et al⁵ surveyed 94 spine surgeons regarding their use of prophylactic anticoagulation, and found significant inconsistencies, with 63% of the surgeons replying that they relied on personal experience to manage anticoagulation. Owing to the lack of published data and evi-

dence, the American College of Chest Physicians (ACCP) guidelines for DVT prevention (8th edition) do not provide definitive guidelines for spine surgery patients, but suggest the use of either pharmacological or mechanical prophylaxis for VTE in patients with other risk factors.³ Pharmacological prophylaxis can be achieved with low dose unfractionated heparin (Grade 1B recommendation) or low molecular weight heparin (LMWH) (Grade 1B recommendation), which is strongly recommended by ACCP guidelines.

One reason for the uncertainty is that the incidence of VTE after elective spine surgery without prophylactic anticoagulation varies in different reports. Glotzbecker et al⁶ undertook an analysis of 25 articles and found a DVT range of 0.3%-31% following spine surgery, with an overall incidence of 2.1% (181/8618). Sansone et al⁷ reviewed 14 studies including 4383 patients, in whom the incidence of DVT was 1.09% and the incidence of PE was 0.06% following elective spine surgery. In patients with pre-existing VTE risk factors, the incidence of postoperative VTE is greatly increased.⁸ Some authors think the incidence is also related to the type of spine surgery, with major spinal reconstruction placing patients at greater risk.^{9,10}

Although the incidence of VTE after elective spine surgery varies and the debate regarding the need for prophylactic anticoagulation is going on, many patients with degenerative lumbar spine have concomitant cardiovascular disease for which postoperative anticoagulation treatment is required. It is therefore also necessary to evaluate the safety of anticoagulation treatment in the patients at high risk early after lumbar spine surgery.

Apart from insufficiency in published data, surgeons have concerns regarding the safety of anticoagulation treatment in the perioperative period, such as the risk of increasing bleeding complications (e.g. subdural haematoma). How to prevent VTE with anticoagulation safely following lumbar decompression surgery in normal clinical practice? To answer this question, a prospective observational study was carried out to evaluate the safety of LMWH administration for VTE prophylaxis following lumbar decompression surgery.

PATIENTS AND METHODS

Patients

The present study enrolled patients admitted to the Peking Union Medical College Hospital from January 2004 to April 2011 for lumbar decompression surgery, who were classified as having high risk or the highest risk of VTE according to the 6th edition of ACCP Venous Thromboembolism

Prevention Guideline and Levels of Thromboembolism Risk in Surgical Patients without Prophylaxis (Table 1).¹¹ Patients with haemorrhagic disease or other contraindications to LMWH were excluded.

The study protocol was approved by the Ethics Committee of Peking Union Medical College Hospital, and informed consent was obtained from all the enrolled patients.

VTE prevention

Following posterior lumbar spinal decompression under general anaesthesia, all patients received a half dose of LMWH (2050 U) administered subcutaneously 6 hours after surgery, followed by a full dose (4100 U) once per day until discharge. Ambulation was encouraged for discharged patient as further VTE prophylaxis.

Wound drain was removed when 24-hour drainage volume was less than 50 mL, and was scheduled 2 hours prior to LMWH administration (Fig. 1).

Table 1. Levels of thromboembolism risk in surgical patients without prophylaxis

Risk level	Level analysis
Low risk	Minor surgery in patients <40 years with no additional risk factors
Moderate risk	Minor surgery in patients with additional risk factors; non-major surgery in patients aged 40-60 years with no additional risk factors; major surgery in patients <40 years with no additional risk factors
High risk	Non-major surgery in patients >60 years or with additional risk factors; major surgery in patients >40 years or with additional risk factors
Highest risk	Major surgery in patients >40 years plus prior VTE, cancer, or molecular hypercoagulable state; hip or knee arthroplasty, hip fracture surgery; major trauma; spinal cord injury

VTE: venous thromboembolism.




Figure 1. Timing of drain removal in lumbar decompression patients receiving low molecular weight heparin (LMWH) treatment.

n: number of days of LMWH administration before drain removal

Efficacy of LMWH prophylaxis

We used the Wells' score to evaluate postoperative DVT and PE probability (Fig. 2).^{12,13} Patients with a DVT score ≥ 2 or a PE score ≥ 4 underwent a D-dimer test. If the D-dimer results were negative ($<500 \mu\text{g/L}$), DVT and PE were temporarily excluded, and patients were closely observed. If the D-dimer results were positive ($\geq 500 \mu\text{g/L}$), lower-limb venous compression ultrasonography (VCUS) was performed in patients with suspected DVT, and computed tomographic pulmonary angiography (CTPA) in patients with suspected PE. Definitive diagnosis of DVT or PE was made if these investigations showed positive results. In patients with negative D-dimer results but with high clinical suspicion of VTE, lower-limb VCUS or CTPA were also performed as a precaution (Fig. 2).

Safety of LMWH prophylaxis

Standard postoperative wound observations, wound drainage volume, and drain removal time were carefully recorded. Patient symptoms were noted, and lower limb sensation, muscle tone and strength were under regular examination. Computed tomography or magnetic resonance imaging was undertaken to rule out compression of the spinal cord by haematoma or a dural haematoma if there was a sudden deterioration in limb signs. Instances of bleeding and observations of LMWH administration sites including skin colour changes were recorded. Major haemorrhage was defined as acute bleeding fulfilling at

least one of the following conditions: decrease in haemoglobin level by 20 g/L or more within 24 hours, transfusion requirement of two units or more of red blood cells, bleeding at a crucial site (including intracranial, spinal canal, intraocular, pericardial, retroperitoneal), bleeding at the surgical site requiring intervention, or intramuscular haemorrhage causing compartment syndrome.¹⁴ All other types of bleeding were defined as minor.

Aminotransferase levels and signs of allergic reaction were recorded. Thrombocytopenia was defined as a postoperative platelet count $<100 \times 10^9/\text{L}$ if the preoperative count was $>150 \times 10^9/\text{L}$, or a 50% reduction after surgery if the preoperative count was $<150 \times 10^9/\text{L}$.¹⁵

RESULTS

General characteristics of the patients

Altogether 78 consecutive patients were eligible and enrolled, of which 10 cases had a diagnosis of lumbar spondylolisthesis, 55 cases had lumbar spinal stenosis, and 13 cases had lumbar disc herniation. The mean age was 62.4 ± 10.6 years, with 29 males (37.2%) and 49 females (62.8%). The average hospital stay was 24.9 ± 20.4 days. Lumbar decompression surgery was performed in 65 cases (83.3%), and lumbar laminectomy and discectomy in 13 cases (16.7%). All the patients received the operation under general anaesthesia, with the mean surgical time being 174.2 ± 62.8 minutes. The mean intraoperative blood




Figure 2. Diagnostic flowchart for DVT and PE following lumbar decompression surgery.

DVT: deep venous thrombosis; PE: pulmonary embolism; CTPA: computed tomographic pulmonary angiography.

*In cases with D-dimer less than $500 \mu\text{g/L}$ but high clinical suspicion of venous thromboembolism, further examination was still performed.

loss was 441 ± 364.7 mL, with 30 cases (38.5%) requiring intraoperative blood transfusion (autologous or allogeneic) and five cases (6.4%) requiring postoperative blood transfusion. Wound drainage tubes were placed in all except three patients with lumbar disc herniation. The mean total drainage volume was 319.5 ± 218.5 mL, and the time of drain removal was 108 ± 36 hours after surgery on average. The mean time to mobilisation after surgery was 5.5 ± 1.9 days (Table 2).

Efficacy and safety analysis results

Postoperative VTE risk assessments categorised 72 cases (92.3%) as having high risk and 6 (7.7%) as having the

highest risk. One of the highest risk patients already had a history of DVT prior to surgery. The average anticoagulation time was 8.5 ± 4.5 days.

In the 3 cases with a postoperative Wells' DVT score ≥ 2 , D-dimer tests were performed, followed by lower-limb VCUS in two cases. No thrombus formation was detected in any of these three cases. In the 4 cases with a postoperative Wells' PE score ≥ 4 , D-dimer tests were conducted, followed by CTPA in one case. PE was excluded in these 4 patients (Table 3). The patient with a DVT history was taking warfarin prior to admission and was switched to LMWH treatment upon hospitalisation; this patient did not experience any VTE recurrence during hospital stay.

Table 2. Baseline, surgical, and postoperative characteristics of the enrolled cases

Characteristics	Lumbar spondylolisthesis (n=10)	Lumbar spinal stenosis (n=55)	Lumbar disc herniation (n=13)	Total (n=78)
Basic information				
Age (years)	64.6 ± 10.7	63.7 ± 10.1	55.3 ± 10.2	62.4 ± 10.6
Gender				
Male	1(10%)	23(41.8%)	5(38.5%)	29(37.2%)
Female	9(90%)	32(58.2%)	8(61.5%)	49(62.8%)
Hospital stay (days)	23.2 ± 5.5	27.2 ± 23.6	16.7 ± 5.3	24.9 ± 20.4
Surgical information				
Surgical approach				
Lumbar decompression	10(100%)	55(100%)	0(0%)	65(83.3%)
Lumbar discectomy	0(0%)	0(0%)	13(100%)	13(16.7%)
General anaesthesia	10(100%)	55(100%)	13(100%)	78(100%)
Surgery time (min)	168.0 ± 38.4	188.9 ± 64.1	117.0 ± 34.0	174.2 ± 62.8
Intraoperative bleeding (mL)	305.0 ± 114.1	538.1 ± 382.8	134.6 ± 154.2	441.0 ± 364.7
Intraoperative blood transfusion	1(10%)	28(50.9%)	1(7.7%)	30(38.5%)
Postoperative information				
Postoperative blood transfusion	1(10%)	4(7.3%)	0(0%)	5(6.4%)
Postoperative drainage (mL)	329.1 ± 83.0	356.6 ± 225.9	$106.1 \pm 142.5^*$	319.5 ± 218.5
Time of drain removal (hours)	114.0 ± 18.0	114.0 ± 30.0	102.0 ± 42.0	108.0 ± 36.0
Time to mobilisation (days)	5.2 ± 2.3	$5.6 \pm 2.0^{\#}$	5.1 ± 1.5	5.5 ± 1.9

*Three cases with lumbar disc herniation did not have postoperative wound drains; #two cases with lumbar spinal stenosis complicated by hemiplegia did not mobilise prior to discharge.

Table 3. VTE risk level and anticoagulation time, efficacy, and safety

Assessment	Lumbar spondylolisthesis (n=10)	Lumbar spinal stenosis (n=55)	Lumbar disc Herniation (n=13)	Total (n=78)
VTE risk level				
High	8(80%)	52 (94.5%)	12 (92.3%)	72 (92.3%)
Highest	2(20%)	3 (5.5%)	1 (7.7%)	6 (7.7%)
Days of anticoagulation	7.5 ± 3.6	9.1 ± 5.0	6.8 ± 1.8	8.5 ± 4.5
Efficacy assessment				
Symptomatic DVT	0(0%)	0(0%)	0(0%)	0(0%)
Symptomatic PE	0(0%)	0(0%)	0(0%)	0(0%)
Safety assessment				
Major bleeding	0(0%)	0(0%)	0(0%)	0(0%)
Minor bleeding	0(0%)	2(3.6%)	0(0%)	2(2.6%)
Thrombocytopenia	0(0%)	0(0%)	0(0%)	0(0%)
Elevated aminotransferase	0(0%)	3(5.5%)	1(7.7%)	4(5.1%)
Suspicious allergic reaction	0(0%)	1(1.8%)	0(0%)	1(1.3%)

No major haemorrhagic events were encountered in this study. One case developed wound ecchymosis and another had wound oozing. Four cases showed mild elevation in aminotransferase level, but none had decreased platelet count or severe heparin-induced thrombocytopenia. One patient developed a suspicious allergic reaction, presenting with red skin rash over the abdomen and both lower limbs without other positive physical signs such as heart rate or blood pressure changes, which recovered with medication (Table 3). LMWH was continued in all the cases.

DISCUSSION

The risk of VTE increases in patients after lumbar decompression surgery, especially in those with additional risk factors.⁸ Chemical prophylaxis in these patients was considered appropriate. In spite of that, bleeding complications, especially subdural haematoma, was the major concern of spine surgeons during chemoprophylaxis due to the exposure of spinal dural mater during surgery.

Regarding complications after anticoagulation treatment following spine surgery, Cheng et al¹⁶ reported that the incidence of anticoagulation-induced haemorrhagic complications was low, with a 0%-4.3% incidence of major bleeding. Gerlach et al¹⁷ analysed 1949 post-spine-surgery patients treated with LMWH, and found only 8 cases of postoperative haematoma, with an incidence of 0.4%. In the present study, there was no occurrence of major haemorrhagic complications, and only one case of wound bruising and another of wound oozing, suggesting that administration of LMWH is quite safe. Previous reports have indicated that adverse reactions of LMWH are low,^{18,19} and no serious adverse reactions were observed in this study. Four cases had elevated aminotransferase level and one had a suspicious allergic reaction, but LMWH was not discontinued in any of these cases. For the patient with a suspicious allergic reaction, no further examinations were taken since the red skin rash disappeared after topical drug treatment. Care should be taken, however, to monitor patients for adverse drug reactions, as indicated by various LMWH pharmacology reports and articles.^{20,21}

This study included patients at high risk and the highest risk of VTE, who received regular LMWH administration along with adjunct mechanical therapy measures. The first dose of LMWH was administered 6 hours after surgery. Postoperative administration protects against interactions with anaesthetic drugs and against increased intraoperative bleeding. This 6-hour postoperative timing has been demonstrated to effectively prevent VTE following major lower limb procedures without increasing bleeding

risks.²² Drain removal was 2 hours prior to LMWH administration, when blood levels were at their lowest, thus reducing the risk of bleeding during the removal process. The 2-hour interval before LMWH administration allowed for clotting within the wound, thus decreasing internal wound bleeding and preventing haematoma formation and compression of the spinal cord.²³ None of the patients in this study bled during the drain removal process.

DVT and PE diagnosis during anticoagulation treatment were made strictly in accordance with Wells' scoring system and D-dimer test results, with suspicious cases undergoing lower-limb VCUS or CTPA. The studies by Wells et al^{12,13} demonstrated that the Wells' analysis together with D-dimer test could efficiently detect VTE, greatly reducing the costs by decreasing the requirements for more expensive VCUS and CTPA.

Voth et al²⁴ compared the efficacy of LMWH versus heparin for VTE prophylaxis in spine surgery. Postoperative treatment time was 8.0 days with LMWH and 7.6 days with heparin. There were only 2 cases (2/92) of symptomatic DVT in the heparin group, and 1 case (1/87) in the LMWH group, with no PE detected in either group. Otero-Fernández et al²⁵ reported the application of LMWH prophylaxis for VTE in 231 spine surgery patients, observing no cases of DVT or PE. The present study also indicates that LMWH is effective for VTE prophylaxis; there were no cases of DVT or PE among the 78 enrolled patients.

Apart from LMWH treatment, early ambulation was encouraged for all patients to prevent VTE in this study. The mean time of mobilization in this group of patients was 5.5±1.9 days. However, two cases with lumbar spinal stenosis complicated by hemiplegia did not mobilise prior to discharge.

The present study has limitations. For instance, we did not undertake routine test for VTE during the anticoagulation period, which may have allowed asymptomatic VTE to go undetected.^{6,7} In conclusion, the present study results show that LMWH could prevent VTE in patients at high risk and the highest risk who underwent lumbar spine surgery, with a low incidence of hemorrhagic complications.

REFERENCES

1. Liew NC, Moissinac K, Gul Y. Postoperative venous thromboembolism in Asia: a critical appraisal of its incidence. *Asian J Surg* 2003; 26:154-8.
2. Cordell-Smith JA, Williams SC, Harper WM, et al. Lower limb arthroplasty complicated by deep venous thrombosis: prevalence and subjective outcome. *J Bone Joint Surg Br*

- 2004; 86:99-101.
3. Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* 2008; 133:381S-453S.
 4. Haas SB, Barrack RL, Westrich G, et al. Venous thromboembolic disease after total hip and knee arthroplasty. *J Bone Joint Surg Am* 2008; 90:2764-80.
 5. Glotzbecker MP, Bono CM, Harris MB, et al. Surgeon practices regarding postoperative thromboembolic prophylaxis after high-risk spinal surgery. *Spine (Phila Pa 1976)* 2008; 33:2915-21.
 6. Glotzbecker MP, Bono CM, Wood KB, et al. Thromboembolic disease in spinal surgery: a systematic review. *Spine (Phila Pa 1976)* 2009; 34:291-303.
 7. Sansone JM, del Rio AM, Anderson PA. The prevalence of and specific risk factors for venous thromboembolic disease following elective spine surgery. *J Bone Joint Surg Am* 2010; 92:304-13.
 8. Anderson FA Jr, Spencer FA. Risk factors for venous thromboembolism. *Circulation* 2003; 107:19-16.
 9. Brambilla S, Ruosi C, La Maida GA, et al. Prevention of venous thromboembolism in spinal surgery. *Eur Spine J* 2004; 13:1-8.
 10. Samama CM, Albaladejo P, Benhamou D, et al. Venous thromboembolism prevention in surgery and obstetrics: clinical practice guidelines. *Eur J Anaesthesiol* 2006; 23: 95-116.
 11. Geerts WH, Heit JA, Clagett GP. Prevention of venous thromboembolism. *Chest* 2001; 119:132S-75S.
 12. Wells PS, Anderson DR, Rodger M, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000; 83: 416-20.
 13. Wells PS, Anderson DR, Rodger M, et al. Evaluation of D-dimer in the diagnosis of suspected deep-vein thrombosis. *N Engl J Med* 2003; 349:1227-35.
 14. Schulman S, Kearon C, Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost* 2005; 3:692-4.
 15. Lassen MR, Gallus A, Raskob GE, et al. Apixaban versus enoxaparin for thromboprophylaxis after hip replacement. *N Engl J Med*. 2010; 363: 2487-98.
 16. Cheng JS, Arnold PM, Anderson PA, et al. Anticoagulation risk in spine surgery. *Spine (Phila Pa 1976)* 2010; 35:S117-24.
 17. Gerlach R, Raabe A, Beck J, et al. Postoperative nadroparin administration for prophylaxis of thromboembolic events is not associated with an increased risk of hemorrhage after spinal surgery. *Eur Spine J* 2004; 13:9-13.
 18. Testroote M, Stigter W, de Visser DC, et al. Low molecular weight heparin for prevention of venous thromboembolism in patients with lower-leg immobilization. *Cochrane Database Syst Rev* 2008; (4):CD006681.
 19. Salazar CA, Malaga G, Malasquez G. Direct thrombin inhibitors versus vitamin K antagonists or low molecular weight heparins for prevention of venous thromboembolism following total hip or knee replacement. *Cochrane Database Syst Rev* 2010; (4):CD005981.
 20. Liebman HA. Heparin-induced thrombocytopenia: diagnosis and management. *Vascular* 2008; 16 Suppl 1:S71-6.
 21. Jappe U. Allergy to heparins and anticoagulants with a similar pharmacological profile: an update. *Blood Coagul Fibrinolysis* 2006; 17:605-13.
 22. Raskob GE, Hirsh J. Controversies in timing of the first dose of anticoagulant prophylaxis against venous thromboembolism after major orthopedic surgery. *Chest* 2003; 124:379S-85S.
 23. Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Third Edition). *Reg Anesth Pain Med* 2010; 35:64-101.
 24. Voth D, Schwarz M, Hahn K, et al. Prevention of deep vein thrombosis in neurosurgical patients: a prospective double-blind comparison of two prophylactic regimens. *Neurosurg Rev* 1992; 15:289-94.
 25. Otero-Fernández R, Gómez-Outes A, Martínez-González J, et al. Evaluation of the effectiveness and safety of bempiparin in a large population of orthopedic patients in a normal clinical practice. *Clin Appl Thromb Hemost* 2008; 14:75-83.